

The Equivalence Postulate of Quantum Mechanics

Alon E. Faraggi¹ and Marco Matone²

¹ *Department of Physics*

University of Minnesota, Minneapolis MN 55455, USA

e-mail: faraggi@mnhepo.hep.umn.edu

² *Department of Physics “G. Galilei” – Istituto Nazionale di Fisica Nucleare*

University of Padova, Via Marzolo, 8 – 35131 Padova, Italy

e-mail: matone@padova.infn.it

Abstract

The removal of the peculiar degeneration arising in the classical concepts of *rest frame* and *time parameterization* is at the heart of the recently formulated Equivalence Principle (EP). The latter, stating that all physical systems can be connected by a coordinate transformation to the free one with vanishing energy, univocally leads to the Quantum Stationary HJ Equation (QSHJE). This is a third-order non-linear differential equation which provides a trajectory representation of Quantum Mechanics (QM). The trajectories depend on the Planck length through hidden variables which arise as initial conditions. The formulation has manifest p - q duality, a consequence of the involutive nature of the Legendre transformation and of its recently observed relation with second-order linear differential equations. This reflects in an intrinsic ψ^D - ψ duality between linearly independent solutions of the Schrödinger equation. Unlike Bohm's theory, there is a non-trivial action even for bound states and no pilot-wave guide is present. A basic property of the formulation is that no use of any axiomatic interpretation of the wave-function is made. For example, tunnelling is a direct consequence of the quantum potential which differs from the Bohmian one and plays the role of particle's self-energy. Furthermore, the QSHJE is defined only if the ratio ψ^D/ψ is a local homeomorphism of the extended real line into itself. This is an important feature as the $L^2(\mathbb{R})$ condition, which in the Copenhagen formulation is a consequence of the axiomatic interpretation of the wave-function, directly follows as a basic theorem which only uses the geometrical gluing conditions of ψ^D/ψ at $q = \pm\infty$ as implied by the EP. As a result, the EP itself implies a dynamical equation that does not require any further assumption and reproduces both tunnelling and energy quantization. Several features of the formulation show how the Copenhagen interpretation hides the underlying nature of QM. Finally, the non-stationary higher dimensional quantum HJ equation and the relativistic extension are derived.

Contents

1	Introduction	1
2	Canonical and coordinate transformations	13
2.1	The classical HJ equation	13
2.2	Coordinate transformations and the distinguished frame	15
2.3	Time parameterization and space in Classical Mechanics	16
2.4	v -transformations	18
2.5	A geometrical picture	19
3	p-q duality and the Legendre transformation	20
3.1	Energy conservation and classical p - q duality	20
3.2	The dual reduced action	22
3.3	The dual HJ equation	23
4	Möbius symmetry and the canonical equation	25
4.1	$\mathcal{T}_0$ -transformation induced by v	25
4.2	The canonical equation	26
4.3	Legendre transformation as Möbius transformation	27
4.4	Möbius invariance of $\mathcal{U}$	27
4.5	Canonical equation, dynamics and initial conditions	28
4.6	Covariance	29
5	Legendre duality	31
5.1	u -transformations	31
5.2	The dual canonical equation	32
5.3	Legendre brackets	32
5.4	p and q as Legendre pair	33
6	Self-dual states	34
6.1	Where do the $\mathcal{S}_0$ - $\mathcal{T}_0$ pictures overlap?	34
6.2	Dilatations	36
6.3	Changing sign: $pq \longrightarrow p^u q_u = -pq = p_v q^v$	36
6.4	Self-dual states: $q^v = q_u, p^u = p_v$	37
7	The Equivalence Principle	39
7.1	Equivalence Principle and v -transformations	39
7.2	Classical Mechanics and the Equivalence Principle	40
7.3	Modifying the classical HJ equation	40
7.4	Covariance as consistency condition	41

8	Implementing the Equivalence Principle	41
8.1	$\mathcal{W}$ states as inhomogeneous terms	42
8.2	The cocycle condition	43
8.3	Cocycle condition and Möbius symmetry	44
8.4	Selecting a self-dual state	46
9	The quantum stationary HJ equation	46
9.1	Complex entries and real values of the Schwarzian derivative	47
9.2	$(q^a; q^b) = -\frac{\beta^2}{4m}\{q^a, q^b\}$	48
9.3	Uniqueness of the solution	52
10	The Schrödinger equation	53
10.1	QSHJE and Schrödinger equation	53
10.2	Wave-function and Copenhagen School	54
10.3	$\mathcal{W} \longleftrightarrow Q$ and the dual Schrödinger equation	56
10.4	$\mathcal{W}^{sd} = \mathcal{W}^0$ and existence of the Legendre transformation	57
10.5	The trivializing map	58
11	Quantum HJ equation and the reality condition	59
11.1	Quantum HJ equation and reality of R and $\hat{\mathcal{S}}_0$	59
11.2	The classical limit	61
11.3	The WKB approximation and the quantum HJ equation	63
12	The wave equation of Classical Mechanics	63
12.1	The wave-function of Classical Mechanics	64
12.2	Probabilistic interpretation of the classical wave-function	64
12.3	$\mathcal{W}$ states in Classical and Quantum Mechanics	65
13	The trivializing map and quantum transformations	66
13.1	On the quantum canonical transformations	68
13.2	Analogy with uniformization theory	68
13.3	Quantum transformations	72
13.4	Classical and quantum potentials and equivalence of curves	73
13.5	Area function and commutator	75
14	Canonical variables and Möbius transformations	76
14.1	The canonical variables of the state $\mathcal{W}^0$	76
14.2	Time and elliptic curve as moduli of the state $\mathcal{W}^0$	77
14.3	The canonical variables of arbitrary states	78
14.4	The symmetries of the wave-function	83

15 Trajectories and the Equivalence Principle	87
15.1 Differential properties of the reduced action	87
15.2 The Equivalence Principle and the trivializing map	88
15.3 Equivalence Principle and local homeomorphicity of v -maps	89
15.4 The Möbius symmetry of the trivializing map	90
15.5 The free particle	93
15.6 Time parameterization	94
15.7 Quantum mass field: dynamical equation	95
16 Equivalence Principle and fundamental constants	96
16.1 $\hbar \longrightarrow 0, E \longrightarrow 0$ and Planck length	97
16.2 Time, energy and trajectories	102
17 Equivalence Principle, tunnelling and quantized spectra	104
17.1 Tunnelling	104
17.2 Quantized spectra from the Equivalence Principle	105
17.3 The index of the trivializing map	111
17.4 Equivalence Principle and admissible potentials	111
18 The potential well and the harmonic oscillators	112
18.1 The potential well	112
18.2 The infinitely deep potential well and the free particle	116
18.3 The simple and double harmonic oscillators	118
18.4 The general case	122
19 Generalizations	124
19.1 Time-dependent Schrödinger equation	124
19.2 Higher dimension	127
19.3 Schrödinger equation and space compactification	131
19.4 Time-dependent case	132
19.5 Relativistic extension	132
20 Conclusions	133

1 Introduction

Twentieth century physics has revolutionized our view and understanding of the fundamental principles underlying all physical phenomena. At its core, General Relativity (GR) and Quantum Mechanics (QM) reign unchallenged in their respective domains. Yet, the mutual coexistence of these two fundamental theories remains elusive after many decades since their inception. While GR is based on underlying geometrical concepts, general covariance and the Equivalence Principle, the traditional formulation of QM does not follow from such principles. Rather, the conventional formulation of QM is based on concepts concerning the measurement process and the related axiomatic interpretation of the wave-function. This indicates that difficulties in understanding the synthesis of these two fundamental theories trace back to the apparently different nature of such principles. A deeper comprehension of such principles constitutes the basic theoretical challenge of the new millennium.

Recently we showed in [1][2] that QM can in fact be derived from an Equivalence Principle (EP) which is reminiscent of Einstein's EP [3]. In this paper we will expand and develop the approach initiated in Refs.[1][2]. We start in sect.2 with a critical investigation which will lead to formulate the EP. This includes a basic analysis of the role of time in Classical Mechanics (CM) and of the distinguished nature of the rest frame. In this process we will reexamine the basic relations among the canonical variables which are at the heart of Hamilton-Jacobi (HJ) theory. In particular, we will uncover new algebraic symmetries which shed surprising new view on the fundamental role of the canonical variables.

A basic view point of our approach may be regarded as addressing an important question. In order to formulate it, let us first recall that in HJ theory the generalized coordinate and momentum q and p are initially regarded as independent variables. The Classical HJ Equation (CHJE) follows from looking for canonical transformations leading to the free system with vanishing energy. The functional dependence among the canonical variables is only extracted after the HJ equation is solved. Let us now consider a variation of this approach and suppose that we start, for a stationary system, from $p = \partial_q \mathcal{S}_0$, with $\mathcal{S}_0$ the Hamiltonian characteristic function, that we will also call reduced action. Let $\mathcal{S}_0(q)$ and $\mathcal{S}_0^v(q^v)$ be the reduced actions of two arbitrary physical systems. The question we are considering is to find the coordinate transformation $q \longrightarrow q^v$ connecting them. This is reminiscent of HJ theory as it would also imply the existence of the coordinate transformation reducing an arbitrary physical system to the free one with vanishing energy. The basic difference with respect to classical HJ theory is that we are considering coordinate transformations only, with the transformation of p being induced by the relation $p = \partial_q \mathcal{S}_0$. Of course, to find the transformation rule of p we need to know how $\mathcal{S}_0$ transforms. On general grounds we can just choose

$$\mathcal{S}_0^v(q^v) = \mathcal{S}_0(q), \tag{1.1}$$

which defines the “ v -transformation” (VT) $q \longrightarrow q^v = v(q) = \mathcal{S}_0^{v^{-1}} \circ \mathcal{S}_0(q)$. However, although (1.1) seems innocuous, requiring the existence of these transformations for any pair of states, is the same that imposing equivalence of states under coordinate transformations. We will show that this leads to QM. For the time being we note that the requirement that (1.1) be well-defined for any system rules

out CM. This follows because of the peculiar nature of the particle at rest, as can be seen from the fact that for the free particle of vanishing energy we have $\mathcal{S}_0^{cl} = cnst$ and no coordinate transformations can connect a constant to a non-constant function. As we will see, this feature is related to the fact that in our formulation, unlike in Bohm's theory [4], the quantum potential, as much as $\mathcal{S}_0$, is never trivial. This is a property at the heart of basic phenomena such as tunnelling and energy quantization.

The impossibility of connecting different states in CM by a coordinate transformation, a consequence of the peculiar nature of the rest frame, disappears if one considers time-dependent coordinate transformations. Observe that in order to introduce time parameterization in HJ theory, one has to identify the conjugate momentum p with the mechanical one $m\dot{q}$. In CM this is equivalent to use Jacobi's theorem $t - t_0 = \partial_E \mathcal{S}_0^{cl}$. Before doing this, time-dependent coordinate transformations cannot be defined. Describing dynamics in terms of the functional relation between p and q is a deep feature of HJ theory. As we said, once time parameterization is introduced, the peculiar nature of the rest frame disappears. For example, the effect of the time-dependent coordinate transformation

$$q' = q - \frac{1}{2}gt^2, \quad (1.2)$$

is that of reducing the motion of a particle in an external gravitational field, $m\ddot{q} = mg$, to $m\ddot{q}' = 0$, corresponding to a free particle of energy E . This includes the particle at rest for which $E = 0$ and $\mathcal{S}_0^{cl} = cnst$, showing that there are no distinguished frames if one uses time-dependent coordinate transformations. Therefore, while with the dynamical description based on the reduced action it is not always possible to connect two systems by a coordinate transformation, this is not the case if one describes the dynamics using Newton's equation. Thus, formulating the EP can be seen as the natural way to remove this asymmetry between the role of space and time, a basic topic discussed in sect.2.

Note that in (1.1), both the old and new coordinates can be seen as independent variables in their own systems. Thus, while $q \longrightarrow q^v$ defines a functional relation between q^v and q , the physics of the old and new systems is described by the functional dependence of $\mathcal{S}_0$ and $\mathcal{S}_0^v$ on q and q^v respectively. While a canonical transformation maps an independent pair (p, q) to a new one (P, Q) , we will see that requiring that all systems be related by a coordinate transformation, fixes $\mathcal{S}_0$ to be solution of the Quantum Stationary HJ Equation (QSHJE), which in turn implies the Schrödinger Equation (SE).

A feature of CM concerns the symmetry between q and p , a topic considered in sect.3. However, this symmetry, which is reflected in the form of the Hamilton equations of motion, is in general lost by the choice of a potential for a given physical system. A central theme of our approach is to seek a formulation in which duality between the canonical variables is manifest. We will show that this is provided by the Legendre transformation. In particular, we will introduce a new generating function

$$\mathcal{S}_0 = p \frac{\partial \mathcal{T}_0}{\partial p} - \mathcal{T}_0, \quad \mathcal{T}_0 = q \frac{\partial \mathcal{S}_0}{\partial q} - \mathcal{S}_0, \quad (1.3)$$

i.e. $q = \partial_p \mathcal{T}_0$. In the stationary case the Hamilton's principal function can be expressed as $\mathcal{S} = \mathcal{S}_0 - Et$. We also have $\mathcal{T} = \mathcal{T}_0 + Et$, where $\mathcal{T} = q \partial_q \mathcal{S} - \mathcal{S}$. In sect.3 we will show that

$$\frac{\partial \mathcal{S}}{\partial t} = -\frac{\partial \mathcal{T}}{\partial t}, \quad (1.4)$$

which holds also in the non-stationary case. This equation guarantees stability of the $\mathcal{S}$ – $\mathcal{T}$ Legendre duality under time evolution. In this context there appears an imaginary factor which arises by selecting the distinguished, or self-dual, states defined as those states which are invariant under the interchange of q and p . Thus, loosely speaking, since the transformation is of order two, one has to consider the “square root” of the minus sign in (1.4). This results in an imaginary factor appearing in the self-dual states, which in turn reflects in its appearance in the expression of $\mathcal{S}_0$ in terms of solutions of the SE. So, Legendre duality, and, in particular, the minus sign in (1.4), is at the heart of the imaginary factor characterizing quantum time evolution.

A sequential important step in our construction concerns the relationship between second-order linear differential equations and the Legendre transformation. Its involutive nature reflects in the existence of dual differential equations. In particular, taking the second derivative with respect to $\mathbf{s} = \mathcal{S}_0(q)$ and $\mathbf{t} = \mathcal{T}_0(p)$ of the first and second equation in (1.3) respectively, we obtain

$$\left(\frac{\partial^2}{\partial \mathbf{s}^2} + \mathcal{U}(\mathbf{s})\right) \begin{pmatrix} q\sqrt{p} \\ \sqrt{p} \end{pmatrix} = 0, \quad \left(\frac{\partial^2}{\partial \mathbf{t}^2} + \mathcal{V}(\mathbf{t})\right) \begin{pmatrix} p\sqrt{q} \\ \sqrt{q} \end{pmatrix} = 0. \quad (1.5)$$

Apparently, there is no new information in these equations. In particular, since they are equivalent to $\mathcal{U}(\mathbf{s}) = \{q, \mathbf{s}\}/2$ and $\mathcal{V}(\mathbf{t}) = \{p, \mathbf{t}\}/2$, where $\{f, x\} = f'''/f' - 3(f''/f')^2/2$ denotes the Schwarzian derivative, one may consider (1.5) as definition of $\mathcal{U}$ and $\mathcal{V}$. However, a first signal that these equations may be relevant, follows from their manifest p – q duality, which is a direct consequence of the dual structure of the Legendre transformations (1.3). Furthermore, suppose that either $\mathcal{U}$ or $\mathcal{V}$ is given. In this case one can consider (1.5) as equations of motion to be solved with respect to the “potential” $\mathcal{U}$ or $\mathcal{V}$. Thus, we would have a dynamical description with manifest p – q duality. Investigation of these equations will shed light on our formulation of QM. In particular, using the transformation properties of the Schwarzian derivative, we see that the “canonical potential” $\mathcal{U}(\mathbf{s})$ and the quantum potential $Q = \hbar^2 \{\mathcal{S}_0, q\}/4m$, which, as we will see, appears in the QSHJE, satisfy the simple relation

$$2mQ + \hbar^2 p^2 \mathcal{U} = 0. \quad (1.6)$$

The issue of p – q duality has an old origin (see *e.g.* Born’s paper [5]) and is a matter of research also in recent literature. Actually, the above relation between the Legendre transformation and second-order linear differential equations, which can in fact be seen to arise from the basic definition of generating function, was introduced in [6] in the framework of Seiberg–Witten theory [7]. In particular, this relation was introduced as a way of inverting the solution for the moduli parameter in Seiberg–Witten theory [7] from $a(u)$ to $u(a)$, where a is the VEV of the scalar component of the $N = 2$ adjoint superfield and $u = \langle \text{tr } \phi^2 \rangle$ is the gauge invariant parameter. In Ref.[8], inspired by the use of the inversion formula in Seiberg–Witten theory [6], we applied the same procedure to the SE. We introduced in QM a prepotential function $\mathcal{F}$, defined by $\psi^D = \partial_\psi \mathcal{F}$, where ψ^D and ψ are two linearly independent solutions of the SE. Thereby, we showed that in QM the spatial coordinate can be regarded as the Legendre transform of $\mathcal{F}$ with respect to the probability density, a proposal further investigated in [9][10]. This relationship between the Legendre transformation and second-order linear differential equations, which shows a ψ^D – ψ duality just as the p – q duality implied (1.5), can be better understood by considering the following basic question posed in [6]

Given a second-order linear differential equation, with linearly independent solutions ψ^D and ψ , find the function $\mathcal{F}(\psi)$ ($\mathcal{F}_D(\psi^D)$) such that $\psi^D = \partial\mathcal{F}/\partial\psi$ ($\psi = \partial\mathcal{F}_D/\partial\psi^D$).

The answer is that the variable q of the differential equation is proportional to the Legendre transform of $\mathcal{F}$ with respect to ψ^2 . Posing the same question with ψ^D and ψ replaced by $q\sqrt{p}$ and $\sqrt{p}$, one sees that the corresponding $\mathcal{F}$ is proportional to the dual reduced action $\mathcal{T}_0$ and

$$2q\sqrt{p} = \frac{\partial\mathcal{T}_0}{\partial\sqrt{p}}. \quad (1.7)$$

The association between the Legendre transformation and second-order linear differential equations is seen to provide complementary ways to access information on a physical theory, with the symmetry properties of the Legendre transformation and its dual reflecting in symmetry properties of the associated differential equations. It was shown in [11] that there are natural “Legendre brackets” underlying this association. Let us consider a “generating function” $U(z)$ and its Legendre transform $V(w) = z\partial_z U(z) - U(z)$, where $w = \partial_z U$ (note that $z = \partial_w V$). Set $u = U(z)$ and

$$\tau = \frac{\partial z \sqrt{w}}{\partial \sqrt{w}} = z + 2 \frac{w}{\partial_z w} = \frac{1}{2} \frac{\partial^2 V}{\partial \sqrt{w}^2}. \quad (1.8)$$

The Legendre brackets are [11]

$$\{X, Y\}_{(u)} = (\partial_u \tau)^{-1} \left(\frac{\partial X}{\partial \sqrt{w}} \frac{\partial}{\partial u} \frac{\partial Y}{\partial \sqrt{w}} - \frac{\partial Y}{\partial \sqrt{w}} \frac{\partial}{\partial u} \frac{\partial X}{\partial \sqrt{w}} \right). \quad (1.9)$$

If the generating function is $\mathcal{S}_0(q)$, then the dual solutions of the first equation in (1.5) satisfy

$$\{\sqrt{p}, \sqrt{p}\}_{(s)} = 0 = \{q\sqrt{p}, q\sqrt{p}\}_{(s)}, \quad \{\sqrt{p}, q\sqrt{p}\}_{(s)} = 1. \quad (1.10)$$

Similarly, considering the generating function $\mathcal{T}_0(p)$, we have

$$\{\sqrt{q}, \sqrt{q}\}_{(t)} = 0 = \{p\sqrt{q}, p\sqrt{q}\}_{(t)}, \quad \{\sqrt{q}, p\sqrt{q}\}_{(t)} = 1. \quad (1.11)$$

Repeating the construction with $\mathcal{F}(\psi^2)$, we obtain

$$\{\psi, \psi\}_{(q)} = 0 = \{\psi^D, \psi^D\}_{(q)}, \quad \{\psi, \psi^D\}_{(q)} = 1. \quad (1.12)$$

Observe that (1.1), under which p transforms as ∂_q , can be seen as the invariance of the Legendre transform of $\mathcal{T}_0(p)$ under VTs.¹ This is reflected in the covariance of the associated differential equation. In general, these equations have a Möbius symmetry which is exhibited by the representation of the potential as a Schwarzian derivative. However, the potential does not remain invariant if the transformations are extended to general coordinate transformations. This indicates that different physical systems, characterized by different potentials, can be connected by coordinate transformations.

¹In sect.5 we will see that this structure admits a dual formulation in which one considers the u -transformations $p \longrightarrow p^u = u(p)$ defined by $\mathcal{T}_0^u(p^u) = \mathcal{T}_0(p)$ which is the dual version of (1.1).

This fact, based on p - q duality, and the necessity of removing the asymmetry between space and time, which appears in considering the specific role of the rest frame in CM, constitute the basic motivations to formulate the EP. The relationship between duality and time parameterization is investigated in sect.3. These two sections, together with sections 4–6, devoted to the algebraic structures underlying p - q duality, constitute the investigation culminating with the EP formulated in sect.7. In this respect we stress that sections 3–7 considerably extend the structures we introduced in [1][2][12]. The content of the equivalence postulate is [1][2]

For each pair $\mathcal{W}^a \equiv V^a(q^a) - E^a$ and $\mathcal{W}^b \equiv V^b(q^b) - E^b$, there is a VT $q^a \longrightarrow q^b = v(q^a)$ such that

$$\mathcal{W}^a(q^a) \longrightarrow \mathcal{W}^{av}(q^b) = \mathcal{W}^b(q^b). \quad (1.13)$$

Note that according to this postulate, all physical systems can be mapped to the one corresponding to $\mathcal{W}^0 \equiv 0$. In this respect we stress that while we do not define a priori to which spaces the quantities $\mathcal{W}$ and $\mathcal{S}_0$ belong, these will be determined by the request that the QSHJE be defined.

A key point in [1] concerns the transformation properties of the Classical Stationary HJ Equation (CSHJE). These properties, discussed in sect.7, can be summarized as follows

Given two physical systems with reduced actions $\mathcal{S}_0^{clv}$ and $\mathcal{S}_0^{cl}$, consider the *classical* v -map

$$q \longrightarrow q^v = v(q) = \mathcal{S}_0^{clv^{-1}} \circ \mathcal{S}_0^{cl}(q), \quad (1.14)$$

that is $\mathcal{S}_0^{clv}(q^v) = \mathcal{S}_0^{cl}(q)$. Then, from the CSHJEs

$$\frac{1}{2m} \left(\frac{\partial \mathcal{S}_0^{clv}(q^v)}{\partial q^v} \right)^2 + \mathcal{W}^v(q^v) = 0, \quad \frac{1}{2m} \left(\frac{\partial \mathcal{S}_0^{cl}(q)}{\partial q} \right)^2 + \mathcal{W}(q) = 0, \quad (1.15)$$

it follows that under *classical* v -maps $\mathcal{W}(q) \longrightarrow \mathcal{W}^v(q^v) = (\partial_{q^v} q)^2 \mathcal{W}(q)$. In particular

$$\mathcal{W}^0(q^0) \longrightarrow (\partial_{q^v} q^0)^2 \mathcal{W}^0(q^0) = 0. \quad (1.16)$$

Hence, in CM it is not possible to connect by a coordinate transformation the state $\mathcal{W}^0$ to other states with $\mathcal{W} \neq \mathcal{W}^0$. Of course, this is the same fact we observed in considering the degeneration which arises for $\mathcal{S}_0^{cl} = \text{const}$. Nevertheless, the above observation clearly indicates that the EP requires a modification of the CSHJE. We therefore add to the CSHJE a function $Q(q)$ to be determined. Existence of CM imposes that in some limit $Q \longrightarrow 0$. Repeating the above analysis with the additional term Q , we see that it is now the term $\mathcal{W} + Q$ that should transform as a quadratic differential under v -maps. That is, as discussed in sect.7, we have

$$(\mathcal{W}^v(q^v) + Q^v(q^v)) (dq^v)^2 = (\mathcal{W}(q) + Q(q)) (dq)^2. \quad (1.17)$$

This property, together with the request that all possible states be connected by a coordinate transformation to the trivial one corresponding to $\mathcal{W}^0 \equiv 0$, is the starting point of the investigation in sect.8.

In particular, this property indicates that $\mathcal{W}$ and Q pick an inhomogeneous term under coordinate transformations. Therefore, we have

$$\mathcal{W}^v(q^v) = (\partial_{q^v} q^a)^2 \mathcal{W}^a(q^a) + (q^a; q^v), \quad (1.18)$$

and

$$Q^v(q^v) = (\partial_{q^v} q^a)^2 Q^a(q^a) - (q^a; q^v). \quad (1.19)$$

Considering the consistency condition for which the transformation connecting $\mathcal{W}^a$ and $\mathcal{W}^c$ should be equivalent to the composition of the two transformations mapping $\mathcal{W}^a$ into $\mathcal{W}^b$, and then $\mathcal{W}^b$ into $\mathcal{W}^c$, we arrive to the basic cocycle condition [1]

$$(q^a; q^c) = (\partial_{q^c} q^b)^2 [(q^a; q^b) - (q^c; q^b)]. \quad (1.20)$$

In sect.8 we will show that (1.20) implies a basic Möbius invariance of $(q^a; q^b)$. Next, in sect.9 we will first show that this symmetry fixes $(q^a; q^b) \propto \{q^a, q^b\}$. More precisely, we will prove that

The cocycle condition (1.20) uniquely defines the Schwarzian derivative up to a global constant and a coboundary term.

We will see that this univocally implies that $Q = \beta^2 \{\mathcal{S}_0, q\}/4m$ and therefore the QSHJE. In sect.10 we will show that the QSHJE implies the SE once β is identified with Planck's reduced constant $\hbar$. Therefore, we will arrive to the remarkable conclusion that one necessitates the appearance of QM if the EP is to be implemented consistently.

In sect.11 we will show that the formulation has a well-defined classical limit. We will also examine the relation with other versions of the QSHJE, in particular with Bohm's representation. In sect.12 we will derive the representation of the CSHJE in terms of a modified SE with the resulting classical wave-function interpreted in terms of probability amplitude.

In sect.13 the trivializing transformations, mapping the SE with non-trivial potentials to the SE with the trivial one, will be derived and discussed in analogy with the trivializing canonical transformations in the classical HJ formalism. In this context we consider the basic analogy with uniformization theory of Riemann surfaces.

The fact that QM may in fact arise from our EP suggests a reconsideration of the role of the potential. Just as in GR the Equivalence Principle leads to a deformation of the geometry induced by the stress tensor, also in QM it should be possible to relate the EP to a deformation of the geometry induced by the potential. In sect.13 we will follow [12] to show that both the classical and quantum potentials are curvature terms arising in projective geometry. Furthermore, we will see that the QSHJE takes the classical form with the spatial derivative ∂_q replaced by $\partial_{\hat{q}}$, where $(\beta^2 = \hbar^2 \mathcal{U})$

$$d\hat{q} = \frac{dq}{\sqrt{1 - \beta^2(q)}}. \quad (1.21)$$

We will conclude sect.13 by showing that the framework of projective geometry considered in [12], allows us to express the Heisenberg commutation relations by means of the area function.

The outcome of the EP is that the CSHJE is modified by a uniquely determined function

$$\frac{1}{2m} \left(\frac{\partial \mathcal{S}_0}{\partial q} \right)^2 + V(q) - E + \frac{\hbar^2}{4m} \{\mathcal{S}_0, q\} = 0, \quad (1.22)$$

where, as we will see in sect.11, the term $Q = \hbar^2 \{\mathcal{S}_0, q\} / 4m$ can be identified as the genuine quantum potential which plays the role of particle's self-energy. Eq.(1.22) is related to the one considered in Bohm's approach to QM [4]. However, $\mathcal{S}_0$ differs from Bohm's phase in important ways. In particular, as a result of the EP, $\mathcal{S}_0$ is never a constant. This reflects in the fact that a general solution of the SE, and therefore also the wave-function, will have the form²

$$\psi = \frac{1}{\sqrt{\mathcal{S}'_0}} \left(A e^{-\frac{i}{\hbar} \mathcal{S}_0} + B e^{\frac{i}{\hbar} \mathcal{S}_0} \right). \quad (1.23)$$

The basic difference with Bohm's theory [4] arises for bound states. In this case the wave-function is proportional to a real function. This implies that with Bohm's identification $\psi = R \exp(i\mathcal{S}_0/\hbar)$, which is also usually considered in standard textbooks, one would have $\mathcal{S}_0 = \text{const}$.³ This seems an unsatisfactory feature of Bohm's theory as there are difficulties in getting a non-trivial classical limit for $\mathcal{S}_0 = \text{const}$, and therefore for $p = 0$. Whereas this would be consistent in the case of classically forbidden regions as there $\mathcal{S}_0^{cl} = \text{const}$, one has troubles when considers regions which are no classically forbidden. In this case the reduced classical action is not trivial whereas with the standard identification $\psi = R \exp(i\mathcal{S}_0/\hbar)$, one would have $\mathcal{S}_0 = \text{const}$. So, for example, while quantum mechanically the particle would be at rest, after taking the $\hbar \rightarrow 0$ limit, the particle should start moving. This problem disappears if one uses Eq.(1.23) which directly follows from the QSHJE: the fact that $\psi \propto \bar{\psi}$ simply implies that $|A| = |B|$ and $\mathcal{S}_0 \neq \text{const}$. Furthermore, we would like to remark that in Bohm's approach some interpretational aspects are related to the concept of pilot-wave guide. There is no need for this in the present formulation. Nevertheless, there are some similarities between our formulation and Bohm's interpretation of QM. In particular, by solving the QSHJE for $\mathcal{S}_0$, we can evaluate $p = \partial_q \mathcal{S}_0$ as a function of the initial conditions, so that we have a determined orbit in phase space. The solutions of the QSHJE, which is a third-order non-linear differential equation, are obtained by utilizing two real linearly independent solutions of the corresponding SE. In particular, denoting by ψ^D and ψ these solutions, we will see in sect.14 that

$$e^{\frac{2i}{\hbar} \mathcal{S}_0 \{\delta\}} = e^{i\alpha} \frac{w + i\bar{\ell}}{w - i\ell}, \quad (1.24)$$

where $\delta = \{\alpha, \ell\}$, with α and ℓ , $\text{Re } \ell \neq 0$, integration constants, and

$$w = \frac{\psi^D}{\psi} \in \mathbb{R}. \quad (1.25)$$

²In the following, unless otherwise specified, by wave-function we will mean the Hamiltonian eigenfunction.

³To be precise, bound states would correspond to $\mathcal{S}_0 = \text{const}$ outside the nodes of the wave-function.

Denoting by $W = \psi' \psi^D - \psi^{D'} \psi = \text{cnst}$ the Wronskian, we have

$$p = \frac{\hbar W(\ell + \bar{\ell})}{2|\psi^D - i\ell\psi|^2}. \quad (1.26)$$

In sect.14 we will consider the effect of the transformation of the initial conditions α and ℓ on the wave-function. In particular, following the second reference in [1], we will see that there are transformations of α and ℓ that leave the wave-function invariant. This is another important signal of the general fact that the SE contains less information than the QSHJE.

A basic geometrical quantity of the formalism concerns the *trivializing map* reducing the system with a given $\mathcal{W}$ to the one corresponding to $\mathcal{W}^0$. In sect.15 we will see that this map is expressed in terms of a Möbius transformation of the ratio $w = \psi^D/\psi$, whose coefficients depend on the initial conditions associated to the states $\mathcal{W}^0$ and $\mathcal{W}$. The role of the phase α reflects in the possibility of reducing this transformation of w to an affine one. More generally, we will see that the trivializing map has a Möbius symmetry.

A property of the formulation is that $p = \partial_q \mathcal{S}_0 \neq m\dot{q}$. In particular, in sect.15 we will follow Floyd [15] who defined time parameterization by means of Jacobi's theorem, according to which $t - t_0 = \partial_E \mathcal{S}_0$. Using the expression for $\dot{q}$, we set $p = m_Q \dot{q}$, where $m_Q = m(1 - \partial_E Q)$ can be seen as an effective quantum mass. Furthermore, we will rewrite the QSHJE (1.22) in terms of m_Q and its derivatives with respect to q , and in terms of q and its time derivatives. This is the dynamical quantum equation for q . An interesting property of this new equation concerns the appearance of the third-order time derivative of q . This is of course related to the appearance of the integration constant ℓ . For this reason, we can consider $\text{Re } \ell$ and $\text{Im } \ell$ as a sort of hidden variables.

We stress that the consistent implementation of the EP is related to the existence of the Legendre transformation for all physical states. This brings to another basic observation concerning the existence of the self-dual states mentioned above and discussed in sect.6. These are the states that remain invariant under the Möbius transformations of q and p corresponding to the common symmetry of the Legendre transformation and of its dual. Alternatively, the self-dual states are precisely those states which are simultaneous solutions of the two dual differential equations (1.5) associated to the two dual Legendre transformations (1.3). From the view point of the QSHJE, we will see that among the self-dual states there is a reference state which precisely corresponds to the trivial one $\mathcal{W}^0 \equiv 0$. While in the classical case this corresponds to $\mathcal{S}_0^{\text{cl}} = \text{cnst}$, we will see that in the quantum case we always have $\mathcal{S}_0 \neq \text{cnst}$. On the other hand, since the free particle with $E = 0$ can be obtained from the one with $E \neq 0$ in the $E \rightarrow 0$ limit, we have that even the more general case $\mathcal{S}_0 = Aq + B$, for which Legendre duality breaks down, is ruled out. Since the QSHJE contains the Schwarzian derivative of $\mathcal{S}_0$, we have that existence of the QSHJE and the exclusion of the solution $\mathcal{S}_0 = Aq + B$, imply that the Legendre transformation is defined for any state. Following [13], we will see in sect.16 that considering the classical limit and the $E \rightarrow 0$ limit of the free particle, leads to introduce a dependence of the initial conditions, or hidden variables, $\text{Re } \ell$ and $\text{Im } \ell$, of the QSHJE on the Planck length. This suggests a connection between GR and our formulation of QM.

Besides the tunnel effect, another crucial phenomenon of QM concerns energy quantization. In our formulation this arises as a consequence of the existence of the QSHJE and therefore is a direct

consequence of the EP itself! In particular, since the identity

$$(\partial_q \mathcal{S}_0)^2 = \frac{\hbar^2}{2} (\{e^{\frac{2i}{\hbar} \mathcal{S}_0}, q\} - \{\mathcal{S}_0, q\}), \quad (1.27)$$

implies that the QSHJE is equivalent to the Schwarzian equation

$$\{w, q\} = -\frac{4m}{\hbar^2} (V - E), \quad (1.28)$$

one has that the QSHJE is well-defined if and only if w is a local self-homeomorphism of the extended real line $\hat{\mathbb{R}} = \mathbb{R} \cup \{\infty\}$. To show this we first note that existence of Eq.(1.28) implies

$$w \neq \text{const}, \quad w \in C^2(\mathbb{R}), \quad \text{and } \partial_q^2 w \text{ differentiable on } \mathbb{R}. \quad (1.29)$$

Observe that by (1.25) these conditions, which arise from the EP, imply the continuity of both ψ^D , ψ and their first derivative, with $\partial_q \psi^D$ and $\partial_q \psi$ differentiable. Therefore, we have

$$\text{Equivalence Principle} \longrightarrow \text{continuity of } (\psi^D, \psi), \text{ and } (\partial_q \psi^D, \partial_q \psi) \text{ differentiable}. \quad (1.30)$$

Note that the QSHJE, and therefore (1.28), is a consequence of the cocycle condition (1.20) that should be always satisfied. In particular, Eq.(1.28) must be equivalent to $\{w, q^{-1}\} = -4mq^4(V - E)/\hbar^2$. On the other hand, since under the map $q \longrightarrow q^{-1}$, the point 0^- (0^+) maps to $-\infty$ ($+\infty$), we have that gluing 0^- to 0^+ corresponds gluing $-\infty$ to $+\infty$. This means that (1.29) extends to $\hat{\mathbb{R}}$, *i.e.*

$$w \neq \text{const}, \quad w \in C^2(\hat{\mathbb{R}}), \quad \text{and } \partial_q^2 w \text{ differentiable on } \hat{\mathbb{R}}. \quad (1.31)$$

We now show that there is another relevant condition. Namely, equivalence under coordinate transformations implies that these transformations must be locally invertible. A property which can be also seen from the cocycle condition (1.20). This amounts to require that transformations such as $\{w, q\} = -(\partial_q w)^2 \{q, w\}$ be well-defined, so that $w(q)$ should be locally invertible, that is $\partial_q w \neq 0$, $\forall q \in \mathbb{R}$. On the other hand, considering the “ S -transformation” $q \longrightarrow q^{-1}$, one sees that also this condition of local univalence should be extended to $\hat{\mathbb{R}}$. It is easy to see that this reflects in the following joining conditions for w at spatial infinities

$$w(-\infty) = \begin{cases} w(+\infty), & \text{for } w(-\infty) \neq \pm\infty, \\ -w(+\infty), & \text{for } w(-\infty) = \pm\infty. \end{cases} \quad (1.32)$$

As we will see, this condition is at the heart of energy quantization. Observe that, as illustrated by the non-univalent function $w = q^2$, making the choice $w(-\infty) = w(+\infty)$ in the case in which $w(-\infty) = \pm\infty$, would break local univalence. The above analysis is the essence of the proof that, according to the EP, w is a local self-homeomorphism of $\hat{\mathbb{R}}$.

The physical interest underlying the above results is due to the following basic theorem [2] that we will review in sect.17. Let us denote by q_- (q_+) the lowest (highest) q for which $V(q) - E$ changes sign. We have that

If

$$V(q) - E \geq \begin{cases} P_-^2 > 0, & q < q_-, \\ P_+^2 > 0, & q > q_+, \end{cases} \quad (1.33)$$

then w is a local self-homeomorphism of $\hat{\mathbb{R}}$ if and only if the corresponding SE has an $L^2(\mathbb{R})$ solution.

Thus, since the QSHJE is defined if and only if w is a local self-homeomorphism of $\hat{\mathbb{R}}$, this theorem implies that energy quantization *directly* follows from the QSHJE itself. Therefore, while in the usual approach to QM one needs the SE with the *additional* $L^2(\mathbb{R})$ condition, in our formulation the EP itself implies a dynamical equation that does not require any further assumption and reproduces both the tunnel effect and energy quantization. Another feature of our approach is that, as we will see in sect.17, the geometrical properties of w define the space of admissible potentials.

It is useful to provide an explicit example illustrating some features of our approach. Other relevant examples are considered in sect.18. We consider the potential well

$$V(q) = \begin{cases} 0, & |q| \leq L, \\ V_0, & |q| > L. \end{cases} \quad (1.34)$$

According to (1.24), in order to solve Eq.(1.22) we need two real linearly independent solutions of the SE. Let us set $k = \sqrt{2mE}/\hbar$, $\kappa = \sqrt{2m(V_0 - E)}/\hbar$ and $Q_{\pm} = \kappa(q \pm L)$. A solution of the SE is

$$\psi = k^{-1} \cdot \begin{cases} -a \exp Q_+ - b \exp -Q_+, & q < -L, \\ \sin(kq), & |q| \leq L, \\ a \exp -Q_- + b \exp Q_-, & q > L, \end{cases} \quad (1.35)$$

where for any $E \geq 0$ the continuity conditions (1.30), implied by the EP, give

$$a = \frac{1}{2} \sin(kL) - \frac{k}{2\kappa} \cos(kL), \quad b = \frac{1}{2} \sin(kL) + \frac{k}{2\kappa} \cos(kL). \quad (1.36)$$

It is easy to see that any solution of the SE can be expressed as a linear combination of ψ and

$$\psi^D = \begin{cases} c \exp Q_+ + d \exp -Q_+, & q < -L, \\ \cos(kq), & |q| \leq L, \\ c \exp -Q_- + d \exp Q_-, & q > L, \end{cases} \quad (1.37)$$

where

$$c = \frac{1}{2} \cos(kL) + \frac{k}{2\kappa} \sin(kL), \quad d = \frac{1}{2} \cos(kL) - \frac{k}{2\kappa} \sin(kL). \quad (1.38)$$

The ratio of the solutions is given by

$$w = k \cdot \begin{cases} -(c \exp Q_+ + d \exp -Q_+)/(a \exp Q_+ + b \exp -Q_+), & q < -L, \\ \cot(kq), & |q| \leq L, \\ (c \exp -Q_- + d \exp Q_-)/(a \exp -Q_- + b \exp Q_-), & q > L. \end{cases} \quad (1.39)$$

Since the asymptotic behavior of w is

$$\lim_{q \rightarrow \pm\infty} w = \pm \frac{d}{b} k, \quad (1.40)$$

we immediately see that the gluing conditions at $\pm\infty$, at $q = \pm\infty$, Eq.(1.32), imply that either

$$b = 0, \quad (1.41)$$

so that $w(-\infty) = -\text{sgn } d \cdot \infty = -w(+\infty)$, or

$$d = 0, \quad (1.42)$$

so that $w(-\infty) = 0 = w(+\infty)$. Eq.(1.41) and (1.42) precisely correspond to the spectrum derived in the usual approach. In particular, according to the above general theorem, the gluing conditions (1.41) and (1.42) correspond to the unique cases in which the SE has an $L^2(\mathbb{R})$ solution. Furthermore, note that if $b = 0$, then $\psi \in L^2(\mathbb{R})$ (and, of course, $\psi^D \notin L^2(\mathbb{R})$), while if $d = 0$, then $\psi^D \in L^2(\mathbb{R})$ (and $\psi \notin L^2(\mathbb{R})$).⁴ Let us find the explicit form of the trivializing map in the case of the potential well. A pair of linearly independent solutions of the SE with $V - E = 0$, is given by $\psi^{D^0} = q^0$, $\psi^0 = 1$. Therefore, $\exp(2i\mathcal{S}_0^0/\hbar) = \exp(i\alpha_0)(q^0 + i\bar{\ell}_0)/(q^0 - i\ell_0)$, and by (1.1)

$$e^{i\alpha_0} \frac{q^0 + i\bar{\ell}_0}{q^0 - i\ell_0} = e^{i\alpha} \frac{w + i\bar{\ell}}{w - i\ell}. \quad (1.43)$$

Since α_0 and α have no effect on the conjugate momenta, we set $\alpha = \alpha_0 + 2\pi k$ to have (see sect.15)

$$q^0 = \frac{(\ell_0 + \bar{\ell}_0)w + i\ell_0\bar{\ell} - i\bar{\ell}_0\ell}{\ell + \bar{\ell}}. \quad (1.44)$$

Let us consider the potential well with $b = 0$. We have

$$q^0 = \frac{k}{\ell + \bar{\ell}} \cdot \begin{cases} -a^{-1}(\ell_0 + \bar{\ell}_0)(c + de^{-2Q_+}) + i\ell_0\bar{\ell} - i\bar{\ell}_0\ell, & q < -L, \\ (\ell_0 + \bar{\ell}_0) \cot(kq) + i\ell_0\bar{\ell} - i\bar{\ell}_0\ell, & |q| \leq L, \\ a^{-1}(\ell_0 + \bar{\ell}_0)(c + de^{2Q_-}) + i\ell_0\bar{\ell} - i\bar{\ell}_0\ell, & q > L, \end{cases} \quad (1.45)$$

where

$$a = -\frac{k}{\kappa} \cos(kL), \quad c = \frac{\kappa^2 - k^2}{2\kappa^2} \cos(kL), \quad d = \frac{\kappa^2 + k^2}{2\kappa^2} \cos(kL). \quad (1.46)$$

One sees that q^0 satisfies all the properties of the trivializing map. In particular, we have

$$\{q^0, q\} = -4m(V - E)/\hbar^2. \quad (1.47)$$

⁴Manifest duality of our formulation, reflects in the appearance in many formulas of both ψ^D and ψ . As illustrated in sect.10, this is the effect of the underlying Legendre duality which reflects in the ψ^D - ψ duality. In this context we note that the above example explicitly shows a general fact that, even if well-known, should be stressed. Namely, since the SE is a second-order linear differential equation, it always has a pair of linearly independent solutions. In particular, by Wronskian arguments, illustrated in sect.17, it can be seen that if the SE has an $L^2(\mathbb{R})$ solution, then any other linearly independent solution is divergent at $q = \pm\infty$. Therefore, the fact that for some system the wave-function belongs to the $L^2(\mathbb{R})$ space, does not mean that for the same E the SE has no divergent solutions. In other words, one should not confuse uniqueness of the wave-function for bound states, which is the well-known non-degeneration theorem of the spectrum for bound states, with the wrong one: “the SE for bound states has only one solution”. Also note that the fact that the divergent solution cannot be the wave-function does not imply that it cannot appear in relevant expressions which in fact characterize the present formulation.

As follows by (1.18) with $\mathcal{W}^a(q^a) = 0$, this is in accordance with the general fact that any $\mathcal{W}$ can be expressed in terms of the inhomogeneous part in the transformation properties of $\mathcal{W}^0$.

We now consider a further simple example which clearly shows the relevance of the continuity at spatial infinity. In particular, we provide an example of the general fact, proved in sect.18, according to which if $V(q) > E$, $\forall q \in \mathbb{R}$, then there are no admissible solutions of the QSHJE. In other words, if there are no classically accessible regions in $\mathbb{R}$, then this also happens quantum mechanically. In the usual approach the absence of solutions when $V - E > 0$, $\forall q \in \mathbb{R}$ is, once again, a consequence of the axiomatic interpretation of the wave-function. Thus we have to understand how this fact emerges (note that we have no turning points, so the previous theorem does not concern this case). The example we consider is the QSHJE with $V - E = a^2$. The associated SE

$$\frac{\hbar^2}{2m} \partial_q^2 \psi = a^2 \psi, \quad (1.48)$$

has the following linearly independent solutions $\psi^D = Ae^{aq} + Be^{-aq}$, $\psi = Ce^{aq} + De^{-aq}$, $AD - BC \neq 0$. Their ratio $w = (Ae^{2aq} + B)/(Ce^{2aq} + D)$, has the asymptotics (we consider $a > 0$)

$$\lim_{q \rightarrow -\infty} w = \frac{B}{D}, \quad \lim_{q \rightarrow +\infty} w = \frac{A}{C}, \quad (1.49)$$

so that neither the $w(-\infty) = \text{finite} = w(+\infty)$ case, nor $w(-\infty) = -w(+\infty) = \pm\infty$ can occur. Hence, for any $a \in \mathbb{R} \setminus \{0\}$, w is not a local homeomorphism of $\hat{\mathbb{R}}$ into itself and therefore the QSHJE does not admit solutions. This means that the EP cannot be implemented in this case.

A peculiarity of QM is that many of its characterizing properties, such as tunnel effect, energy quantization, Hilbert space structure and Heisenberg uncertainty relations, already appear in one spatial dimension. Thus it is not surprising that our formulation extends to the higher dimensional case. This generalization, including the time-dependent case and the relativistic extension, will be investigated in [16] and is shortly considered in sect.19.

In sect.20 we will make some concluding comments concerning the results obtained and possible developments of our formulation.

We conclude this Introduction by making some bibliographic remarks. First, we note that geometrical concepts have been used in relevant approaches to QM such as geometric quantization, developed by Kirillov, Konstant, Guillemin, Soriau, Sternberg et al. (see *e.g.* [17]), and coherent states quantization (see *e.g.* [18]). Furthermore, besides Bohm's theory, another interesting formulation based on the quantum HJ equation, seen as describing the Madelung fluid, concerns the stochastic quantization initiated by Nelson and further developed by Guerra et al. (see *e.g.* [19]). As shown by Gozzi [20], HJ theory appears in QM also in deriving the SE from the anomalous conservation law associated to the overall rescaling of the Lagrangian. We also mention Periwal's paper [21], where a version of the quantum HJ equation has been obtained in the path-integral framework in a way related to our approach. Furthermore, we note that an EP in the framework of QM has been recently considered by Anandan [22], while an interesting investigation of Einstein's EP in QM has been done in [23], where a free fall in a constant gravitational field is considered in the case of quantum states with and without classical analogue. Another interesting approach to QM has been recently considered by Cini

[24] who formulated QM as a generalization of Classical Statistical Mechanics, and in which it is used Feynman's proposal of dropping the assumption that the probability for an event must always be a positive number. Last but not least, we mention some of the relevant approaches to QM proposed in the last decade, such as the decoherent histories [25], quantum state diffusion [26], quaternionic QM [27] and the GRW spontaneous localization model of wave-function's collapse [28].

2 Canonical and coordinate transformations

In this section we first consider basic properties of canonical transformations in CM. This analysis will bring us to consider the problem of finding the coordinate transformations connecting different systems in the case in which p is considered as *dependent* on q . In this context we will show how in CM the rest frame is a privileged one. Existence of this frame can be seen as a possible motivation for introducing time parameterization of trajectories: this makes it possible to define non singular, although time-dependent, coordinate transformations connecting different systems. This investigation, based on the peculiar nature of the rest frame of CM, is the first step towards the modification of classical HJ theory that will lead to a formalism with manifest p - q duality which, as we will see, underlies the EP. We conclude this section by considering a geometrical picture in which transformations connecting different states are seen as patch transformations on a suitable manifold.

2.1 The classical HJ equation

Let us consider some properties of classical canonical transformations. Given Hamilton's equations

$$\dot{q} = \frac{\partial H}{\partial p}, \quad \dot{p} = -\frac{\partial H}{\partial q}, \quad (2.1)$$

the canonical transformations are the transformations

$$q \longrightarrow Q = Q(q, p, t), \quad p \longrightarrow P = P(q, p, t), \quad (2.2)$$

which leave Hamilton's equations form invariant, that is

$$\dot{Q} = \frac{\partial \tilde{H}}{\partial P}, \quad \dot{P} = -\frac{\partial \tilde{H}}{\partial Q}, \quad (2.3)$$

where $\tilde{H}(Q, P, t)$ is the new Hamiltonian. Furthermore, there is the condition that the dynamics must be equivalent to (2.1), that is

$$p\dot{q} - H = P\dot{Q} - \tilde{H} + \frac{dF}{dt}, \quad (2.4)$$

where the generating function F depends on t and on any of the pairs (q, Q) , (q, P) , (p, Q) and (p, P) considered as independent variables. Choosing (q, Q) yields

$$p = \frac{\partial F}{\partial q}, \quad P = -\frac{\partial F}{\partial Q}, \quad (2.5)$$

so that

$$H(q, p, t) + \frac{\partial F}{\partial t} = \tilde{H}(Q, P, t). \quad (2.6)$$

The HJ equation arises by considering the transformation that leads to a vanishing Hamiltonian

$$\tilde{H} = 0, \quad (2.7)$$

which is equivalent to $\dot{Q} = 0$, $\dot{P} = 0$. Once the canonical transformation $Q = cnst = Q(q, p, t)$, $P = cnst = P(q, p, t)$ has been found, the dynamical problem reduces to the inversion problem

$$\begin{cases} Q = cnst = Q(q, p, t), \\ P = cnst = P(q, p, t), \end{cases} \implies \begin{cases} p = p(Q, P, t), \\ q = q(Q, P, t), \end{cases} \quad (2.8)$$

so that Q and P play the role of initial conditions. By (2.5) the associated generating function $\mathcal{S}^{cl}(q, Q, t)$, called Hamilton's principal function, satisfies

$$p = \frac{\partial \mathcal{S}^{cl}}{\partial q}, \quad P = cnst = -\frac{\partial \mathcal{S}^{cl}}{\partial Q}|_{Q=cnst}. \quad (2.9)$$

Eqs.(2.6)(2.7) and (2.9) imply the classical HJ equation

$$H\left(q, p = \frac{\partial \mathcal{S}^{cl}}{\partial q}, t\right) + \frac{\partial \mathcal{S}^{cl}}{\partial t} = 0, \quad (2.10)$$

that for Hamiltonians of the form

$$H = \frac{p^2}{2m} + V(q, t), \quad (2.11)$$

becomes

$$\frac{1}{2m} \left(\frac{\partial \mathcal{S}^{cl}}{\partial q} \right)^2 + V(q, t) + \frac{\partial \mathcal{S}^{cl}}{\partial t} = 0. \quad (2.12)$$

For a time-independent potential there is the decomposition

$$\mathcal{S}^{cl}(q, Q, t) = \mathcal{S}_0^{cl}(q, Q) - Et, \quad (2.13)$$

with E the energy of the stationary state. The function $\mathcal{S}_0^{cl}$ is called Hamilton's characteristic function or reduced action that by (2.10) satisfies the Classical Stationary HJ Equation (CSHJE)

$$H\left(q, p = \frac{\partial \mathcal{S}_0^{cl}}{\partial q}\right) - E = 0, \quad (2.14)$$

that is ($\mathcal{W}(q) \equiv V(q) - E$)

$$\frac{1}{2m} \left(\frac{\partial \mathcal{S}_0^{cl}}{\partial q} \right)^2 + \mathcal{W} = 0. \quad (2.15)$$

2.2 Coordinate transformations and the distinguished frame

At this stage it is worth making some remarks concerning the derivation of the classical HJ equation. First of all, observe that while for an arbitrary canonical transformation one passes from a couple of independent variables (q, p) to another one (Q, P) , in the case in which one uses the Hamilton's principal function as generating function, one has that the right hand side of (2.6) vanishes with the effect that the independent variables become dependent, that is

$$\tilde{H}(Q, P, t) = 0 \quad \longrightarrow \quad p = \partial_q \mathcal{S}^{cl}(q, Q, t). \quad (2.16)$$

Hence, the effect of having $\tilde{H}(Q, P, t) = 0$ is just that of collapsing p and q to be dependent variables. In the case of stationary systems we have

$$H(q, p) \xrightarrow{\text{Canon. Transf.}} \tilde{H}(Q, P) \xrightarrow{\tilde{H}=0} \text{CSHJE}. \quad (2.17)$$

Let us now consider a similar question to that leading to the CSHJE but starting with

$$p = \frac{\partial \mathcal{S}_0^{cl}}{\partial q}, \quad (2.18)$$

rather than with p and q considered as independent variables. In other words, we are looking for the coordinate transformation defined by the following analogue of the transformation (2.17)

$$\mathcal{S}_0^{cl}(q) \xrightarrow{\text{Coord. Transf.}} \tilde{\mathcal{S}}_0^{cl}(\tilde{q}), \quad (2.19)$$

with $\tilde{\mathcal{S}}_0^{cl}(\tilde{q})$ denoting the reduced action of the system with vanishing Hamiltonian. Note that since we consider p and q dependent, the transformation we are considering is not a canonical one. Even if we still have to define the specific structure of the coordinate transformation, since $\tilde{\mathcal{S}}_0^{cl}(\tilde{q}) = \text{cnst}$ it is clear that (2.19) is a degenerate transformation. The existence of this degenerate case is a rather peculiar one. Let us consider two free particles of mass m_A and m_B moving with relative velocity v . For an observer at rest with respect to the particle A the two reduced actions are

$$\mathcal{S}_0^{cl A}(q_A) = \text{cnst}, \quad \mathcal{S}_0^{cl B}(q_B) = m_B v q_B. \quad (2.20)$$

It is clear that there is no way to have an equivalence under coordinate transformations by setting $\mathcal{S}_0^{cl B}(q_B) = \mathcal{S}_0^{cl A}(q_A)$. This means that at the level of the reduced action there is no coordinate transformation making the two systems equivalent. However, note that this coordinate transformation exists if we consider the same problem described by an observer in a frame in which both particles have a non-vanishing velocity so that the two particles are described by non-constant reduced actions. Therefore, in CM, it is possible to connect different systems by a coordinate transformation except in the case in which one of the systems is described by a constant reduced action. This means that in CM equivalence under coordinate transformations is frame dependent.

Let us consider the case in which the two particles are described by an observer in a frame which is accelerated with respect to them. The reduced actions of the A and B particles as seen by the observer with constant acceleration a are

$$\tilde{\mathcal{S}}_0^{cl A}(Q_A) = \frac{m_A}{3a} (2aQ_A)^{\frac{3}{2}}, \quad \tilde{\mathcal{S}}_0^{cl B}(Q_B) = \frac{m_B}{3a} (v^2 + 2aQ_B)^{\frac{3}{2}}, \quad (2.21)$$

where Q_A (Q_B) is the coordinate of the particle A (B) in the accelerated frame. If in describing the particle B in the accelerated frame one uses the coordinate Q_A defined by $\tilde{\mathcal{S}}_0^{cl A}(Q_A) = \tilde{\mathcal{S}}_0^{cl B}(Q_B)$, one has that the resulting dynamics coincides with the one of the particle A , that is

$$\tilde{\mathcal{S}}_0^{cl B}(Q_B(Q_A)) = \tilde{\mathcal{S}}_0^{cl A}(Q_A). \quad (2.22)$$

This simply means that the system B , described in terms of the coordinate Q_A , coincides with the system A . Hence, in CM the equivalence under coordinate transformations requires choosing a frame in which no particle is at rest. In other words, while in any frame two systems are equivalent under coordinate transformations, this is not anymore true once the frame coincides with the one in which the particle is at rest, so that in the CSHJE description there is a distinguished frame. This seems peculiar as on general grounds what is equivalent under coordinate transformations in all frames should remain so even in the one at rest.

2.3 Time parameterization and space in Classical Mechanics

A quite remarkable property of the CSHJE is that it provides a functional relation between p and q . In particular, note that in HJ theory, time-dependent coordinate transformations cannot be defined. Thus, loosely speaking, dynamics is described without using time parameterization. The relevance of this is that we can consider possible relations among different systems without introducing time parameterization, a concept that, as we will see, is related to the privileged nature of the rest frame. Experience in Special and General Relativity indicates that privileged situations may in fact be a consequence of underlying unjustified and somehow hidden assumptions. In Special Relativity the concept of absolute time was shown to be inconsistent. Here we are in similar situation, but our approach is more drastic as we need to start in the framework of HJ theory in which the concept of time does not appear directly. As a matter of fact, this property of HJ theory is in fact at the heart of our formulation of Quantum Mechanics. HJ theory provides the equation for the reduced action which in turns fixes the relationship between p and q . While in the Hamilton and Lagrange equations time derivatives appear also in the stationary case, in HJ theory the time parameterization is introduced only after one uses Jacobi's theorem $t - t_0 = \partial_E \mathcal{S}_0^{cl}$. In CM this is equivalent to identify the conjugate momentum with the mechanical one. Namely, setting

$$p = \partial_q \mathcal{S}_0^{cl} = m\dot{q}, \quad (2.23)$$

yields

$$t - t_0 = m \int_{q_0}^q \frac{dx}{\partial_x \mathcal{S}_0^{cl}(x)}, \quad (2.24)$$

which also provides the solution of the equation of motion $q = f(t)$. It is just the implicit role of time in the CSHJE at the basis of the peculiar nature of the frame at rest we considered before. Let us consider the equation of motion of a particle in an external gravitational field

$$m\ddot{q} = mg. \quad (2.25)$$

Performing the time-dependent coordinate transformation

$$q' = q - \frac{1}{2}gt^2, \quad (2.26)$$

we have

$$m\ddot{q}' = 0, \quad (2.27)$$

for any value of the energy E of the particle, including the free particle at rest for which $E = 0$. So that, depending on the initial conditions of (2.27), we may have q' to be constant, say $q' = 0$. Therefore, there are no selected frames if one uses time-dependent coordinate transformations. Hence, while with the CSHJE description it is not always possible to connect two systems by a coordinate transformation, this is not the case if one describes the dynamics using Newton's equation. In particular, in finding the coordinate transformation reducing the CSHJE description of Eq.(2.25) one has

$$\frac{1}{2m} \left(\frac{\partial \mathcal{S}_0^{cl}(q)}{\partial q} \right)^2 - mgq + E = 0, \quad (2.28)$$

for which there is no coordinate transformation $q \longrightarrow \tilde{q}(q)$ such that $\mathcal{S}_0^{cl}(q) = \tilde{\mathcal{S}}_0^{cl}(\tilde{q})$ with $\tilde{\mathcal{S}}_0^{cl}(\tilde{q})$ the reduced action of the free particle with $E = 0$. In this context we observe that if for some reason $V - E$ were never vanishing, then the above peculiarity of the CSHJE would not exist.

Time parameterization can be seen as a way to express a constant, say 0, by means of the solution of the equation of motions, $q = f(t)$. For example, for a particle with constant velocity, we have

$$0 = q - vt, \quad (2.29)$$

so that particle's position can be denoted by either q itself or vt . In this way one can always reduce to the particle at rest by simply setting $q' = q - f(t)$. While in the case of the CSHJE description there is the degenerate case $cnst = mvq$, corresponding to $\tilde{\mathcal{S}}_0^{cl}(\tilde{q}) = \mathcal{S}_0^{cl}(q)$, time parameterization provides a well-defined and invertible transformation *i.e.* $q' = q - f(t) \longrightarrow q = q' + f(t)$. The reason underlying the differences in considering the role of space and time is that fixed values of q and t correspond to quite different situations. Even if the particle is at rest, say at $q = 0$, *time continues to flow*. It is just the use of time that allows to connect different systems by a coordinate transformation.

The basic difference in the nature of space and time is rather peculiar of CM. On the other hand, in special relativity space and time are intrinsically related. In particular, this property is not unrelated to the fact that in the framework of special relativity the energy of the particle at rest is non-zero. Recalling that in the relativistic case the CSHJE has the form

$$\frac{1}{2m} \left(\frac{\partial \mathcal{S}_0^{cl}}{\partial q} \right)^2 + \mathcal{W}_{rel} = 0, \quad (2.30)$$

where

$$\mathcal{W}_{rel} \equiv \frac{1}{2mc^2} [m^2 c^4 - (V - E)^2], \quad (2.31)$$

we see that the critical $\mathcal{W}_{rel} = 0$ case corresponds to

$$V - E = \pm mc^2. \quad (2.32)$$

Our aim is to find a dynamical description in which there always exists coordinate transformations connecting arbitrary systems. We can see from the above discussion that such an investigation should imply the existence of a sort of “energy function” which is never vanishing, whatever structure the potential V may have. This would avoid the degenerate situation in which the reduced action is a constant and will remove the consequence of the asymmetric role of space and time discussed above. We will see that imposing the EP will univocally lead to the deformation of the CSHJE by a term which is identified with the quantum potential.

We anticipated above some features underlying the EP, which we will formulate later. As we will see, looking for coordinate transformations connecting different physical systems, and in particular for those reducing any system to that corresponding to $\mathcal{W}^0 \equiv 0$, will bring us to the conclusion that the CSHJE must be modified. This will define a new function $\mathcal{S}_0$ that by abuse of language we will continue to call reduced action. Thus, we will have

$$\mathcal{S}_0 \neq \mathcal{S}_0^{cl}. \quad (2.33)$$

Information about a physical system is encoded in the functional dependence of $\mathcal{S}_0$ on its argument. Therefore, a given transformation would reflect in a change of the functional structure of $\mathcal{S}_0$. In this context the specific choice of the coordinate reduces to a matter of notation with the new coordinate playing the role of independent variable for the new system. The coordinate transformation defined in (2.22) is a natural way to define an induced transformation of $\mathcal{S}_0$. Let us consider the following simple example. Given two functions, say

$$f_1(x_1) = x_1^m, \quad f_2(x_2) = x_2^n, \quad (2.34)$$

there is the associated coordinate transformation $x_1 \longrightarrow x_2 = v(x_1) = x_1^{m/n}$, defined by

$$f_2(x_2) = f_1(x_1). \quad (2.35)$$

This is equivalent to saying that starting from the function $f_1(x_1) = x_1^m$, the transformation $x_1 \longrightarrow x_2 = v(x_1) = x_1^{m/n}$, induces the functional transformation $f_1 \longrightarrow f_2 = f_1 \circ v^{-1}$.

2.4 v -transformations

Let us now consider the case of the reduced action $\mathcal{S}_0$ and let

$$q \longrightarrow q^v = v(q), \quad (2.36)$$

be a locally invertible coordinate transformation. Similarly to Eq.(2.35), setting

$$\mathcal{S}_0^v(q^v) = \mathcal{S}_0(q(q^v)), \quad (2.37)$$

naturally defines a new reduced action $\mathcal{S}_0^v$ associated to such “ v -transformations” (VTs). Note that we may also consider alternative transformations of $\mathcal{S}_0$. In general, we can also choose

$$\mathcal{S}_0 \longrightarrow \tilde{\mathcal{S}}_0 = \mathcal{S}_0 \circ v^{-1} + f, \quad (2.38)$$

for some function f . The construction in this case would be equivalent. In particular, since for any transformation (2.38) we can always find another map $q \longrightarrow q^w = w(q)$, such that $\mathcal{S}_0 \longrightarrow \tilde{\mathcal{S}}_0 = \mathcal{S}_0^w = \mathcal{S}_0 \circ w^{-1}$, it is clear that we can directly consider (2.37). This is a convenient choice as our formulation will be particularly simplified and will exhibit duality properties which are generally more difficult to recognize in working with the form (2.38). We note that this is not a restriction as we are considering the problem of connecting different systems, with q^v seen as the new independent coordinate. Then, what is of interest in this context is the functional change $\mathcal{S}_0 \longrightarrow \mathcal{S}_0^v$. Therefore, considering

$$\mathcal{S}_0 \longrightarrow \mathcal{S}_0^v = \mathcal{S}_0 \circ v^{-1}, \quad (2.39)$$

which is equivalent to (2.37), does not imply loss of generality. Note that this is equivalent to saying that associated to $v = \mathcal{S}_0^{v^{-1}} \circ \mathcal{S}_0$ there is the induced map $v^{-1*} : \mathcal{S}_0 \mapsto v^{-1*}(\mathcal{S}_0)$. In other words, $\mathcal{S}_0^v$ is the pullback of $\mathcal{S}_0$ by v^{-1*} .

Let us anticipate a motivation which illustrates why the formalism will simplify in considering the induced transformation in the form (2.39). The point is that as $p_v = \partial_{q^v} \mathcal{S}_0^v(q^v)$, we have

$$p \longrightarrow p_v = (\partial_q q^v)^{-1} p, \quad (2.40)$$

that is, from the point of view of the conjugate momentum, one has that the choice (2.37) implies that p transforms as ∂_q . As a consequence, the resulting formalism will be manifestly covariant.

2.5 A geometrical picture

Let us conclude this section by observing that Eq.(2.37) admits the following geometrical interpretation. Suppose that the system has an underlying geometry induced by $\mathcal{W} \equiv V - E$ and that the use of q corresponds to a choice of a local coordinate representing a point x of an underlying manifold. Then the map $q \longrightarrow q^v = v(q)$ can be seen as a patch transformation with $\mathcal{S}_0$ seen as a scalar function, that is a zero-differential. Since $p = \partial_q \mathcal{S}_0$, we have that the conjugate momentum transforms as a one-differential. Let us recall that a λ -differential $f^{(\lambda)}$ is a set of functions $f^{(\lambda)} = \{f_\alpha^{(\lambda)}(q_\alpha) | \alpha \in I\}$, where each $f_\alpha^{(\lambda)}$ is defined on the patch U_α of the atlas $\{(U_\alpha, q_\alpha) | \alpha \in I\}$, and satisfying

$$f_\alpha^{(\lambda)}(q_\alpha)(dq_\alpha)^\lambda = f_\beta^{(\lambda)}(q_\beta)(dq_\beta)^\lambda, \quad (2.41)$$

in $U_\alpha \cap U_\beta$. In other words, $f^{(\lambda)}(q)$ transforms as $(dq)^{-\lambda}$, that is $f_\alpha^{(\lambda)}(q_\alpha) = (\partial_{q_\alpha} q_\beta)^\lambda f_\beta^{(\lambda)}(q_\beta)$. In this context the structure of the transition functions between the patches U_α and U_β is determined by the α - and β -physical systems, so that there is the correspondence

$$(\mathcal{W}^\alpha, \mathcal{W}^\beta) \longleftrightarrow (U_\alpha, U_\beta). \quad (2.42)$$

3 p - q duality and the Legendre transformation

A feature of CM is that while in considering canonical transformations and the phase space there is a formal p - q duality, this is broken in the explicit solution of the equations of motion. For example, for the Hamiltonian one usually considers the structure $H = p^2/2m + V(q)$, so that, despite the symmetric structure of the equations of motion $\dot{q} = \partial_p H$, $\dot{p} = -\partial_q H$, the effective dynamical solution breaks p - q duality due to the difference between the kinetic term and $V(q)$ structures. There is however a particular case in which H has explicit p - q duality. This happens for the harmonic oscillator for which $V(q) = m\omega^2 q^2/2$. This is an interesting property if one thinks that quantum field theories are described by infinitely many interacting oscillators. However, rather than explicit p - q duality, what is lacking in CM is a formulation in which the p and q descriptions have the same structure.

To understand p - q duality of Hamilton's equations, it is useful to first investigate some properties of CM's formalism. This will be useful in setting the basis for a formulation with manifest duality between the p and q descriptions. In doing this we will see that there is a remarkable way to characterize the Hamilton equations of motion. This shows, in a way illustrated in the next subsection, how the issue of dependence – independence of the canonical variables is connected to dynamical features.

3.1 Energy conservation and classical p - q duality

We have seen that there are interesting questions which arise in considering p and q as dependent quantities through the still unknown $\mathcal{S}_0$. We now show that, considering the Hamiltonian equations of motion as a way to reduce independent variables to dependent ones, will give a connection between energy conservation and classical p - q duality. A step in this subsection concerns the analysis of the relation among the structures of relevant equations which arise by considering the canonical variables as dependent on one side, and those with the variables considered as independent on the other. We will start with an arbitrary function and then we will derive the Hamilton equations of motion from a perspective which is somewhat different from the standard one. This analysis will connect energy conservation and p - q duality.

Let $F(x_1, x_2, x_3)$ be an arbitrary function of the independent variables x_1, x_2, x_3 , so that

$$\frac{dF}{dx_k} = \frac{\partial F}{\partial x_k}, \quad (3.1)$$

$k = 1, 2, 3$, and suppose that F satisfies a differential equation

$$\mathcal{P}(F) = 0, \quad (3.2)$$

with the property that its solution defines

$$x_1 = x_1(x_3), \quad x_2 = x_2(x_3). \quad (3.3)$$

The pair $(F, \mathcal{P})$ naturally defines the function in one variable

$$G(x_3) = F(x_1(x_3), x_2(x_3), x_3), \quad (3.4)$$

whose derivative is

$$\frac{\partial G}{\partial x_3} = \frac{dG}{dx_3} = \frac{dF}{dx_3} = \frac{\partial F}{\partial x_1} \frac{\partial x_1}{\partial x_3} + \frac{\partial F}{\partial x_2} \frac{\partial x_2}{\partial x_3} + \frac{\partial F}{\partial x_3}. \quad (3.5)$$

Note that, while the total and partial derivatives of F with respect to x_3 coincide for F seen as function of the independent variables x_1, x_2, x_3 , in the case in which F is considered as solution of (3.2) one has

$$\frac{dF}{dx_3} \neq \frac{\partial F}{\partial x_3}. \quad (3.6)$$

There is however an exception to this. This happens if (3.2) implies

$$\frac{\partial F}{\partial x_1} \frac{\partial x_1}{\partial x_3} = -\frac{\partial F}{\partial x_2} \frac{\partial x_2}{\partial x_3}, \quad (3.7)$$

so that

$$\frac{dF}{dx_3} = \frac{\partial F}{\partial x_3}. \quad (3.8)$$

Therefore, Eq.(3.1) is preserved for $k = 3$ even after the “equations of motion” (3.7) are imposed. It follows that from the equation

$$\left(\frac{d}{dx_3} - \frac{\partial}{\partial x_3} \right) F = 0, \quad (3.9)$$

it is not possible to distinguish whether F is to be considered as a function of three independent variables x_1, x_2, x_3 or rather if these satisfy the equations of motion (3.7) so that $x_1 = x_1(x_3)$ and $x_2 = x_2(x_3)$. A property of Eq.(3.7) is that it is invariant under the interchange

$$x_1 \longleftrightarrow x_2. \quad (3.10)$$

However, if one considers

$$\frac{\partial x_1}{\partial x_3} = \frac{\partial F}{\partial x_2}, \quad \frac{\partial x_2}{\partial x_3} = -\frac{\partial F}{\partial x_1}, \quad (3.11)$$

as a particular solution of (3.7), then there is a slightly different version of the duality (3.10), since the symmetry of Eq.(3.11) is now

$$x_1 \longrightarrow -x_2, \quad x_2 \longrightarrow x_1. \quad (3.12)$$

Making the identification

$$F = H, \quad (3.13)$$

where H is the Hamiltonian and $x_1 = q, x_2 = p, x_3 = t$, the equation (3.11) corresponds to

$$\dot{q} = \frac{\partial H}{\partial p}, \quad \dot{p} = -\frac{\partial H}{\partial q}. \quad (3.14)$$

Hence, the Hamiltonian equations can be seen as a particular solution to the problem of finding the structure of $\mathcal{P}(F) = 0$ such that (3.9) is satisfied irrespectively of considering x_1 and x_2 as dependent of x_3 or not. In other words, both H as function of the independent variables q, p, t , and H evaluated on the classical trajectory, satisfy

$$\left(\frac{d}{dt} - \frac{\partial}{\partial t} \right) H(q, p, t) = 0 = \left(\frac{d}{dt} - \frac{\partial}{\partial t} \right) H(q^{cl}(t), p^{cl}(t), t). \quad (3.15)$$

This aspect is related to duality as interchanging $x_1 \longleftrightarrow x_2$ leaves (3.7) invariant. As we have seen, since Hamilton's equations (3.14) correspond to a particular solution of Eq.(3.7), it follows that (3.14) is invariant under

$$q \longrightarrow -p, \quad p \longrightarrow q. \quad (3.16)$$

The above discussion shows that classical p - q duality is related to the way in which Eq.(3.7), which makes (3.9) insufficient to distinguish between dependent and independent variables, is specifically satisfied. In particular, we have seen that the particular solution (3.11) of (3.7) breaks the $p \longleftrightarrow q$ duality to the duality (3.16). Therefore, classical p - q duality is related to the dependence that the structure of relevant equations may have on the canonical variables in the dependent and independent cases. In particular, since (3.15) implies energy conservation if the Hamiltonian does not explicitly depend on time, we see that classical p - q duality is a manifestation of the role of time as parameter for classical trajectories and to energy conservation. The fact that in QM position and momentum cannot be simultaneously measured, indicates that the issue of dependence and independence of the canonical variables is a basic feature.⁵ In this context we will see that the p - q duality of Hamilton mechanics is a restrictive one. This lack of the Hamiltonian formalism indicates that a manifest full p - q duality deserves formulating an alternative description of the dynamical problem. In fact, there is a natural structure, with explicit p - q duality, which is a consequence of the involutive nature of the Legendre transformation, and that cannot be a feature of CM. This duality, which arises by considering the canonical variables on equal footing, will make it quite natural to formulate the EP. However, we stress that the QSHJE will be derived from this principle, formulated in sect.7, without assumptions about the existence of dualities.

3.2 The dual reduced action

As a first step towards a formulation with manifest p - q duality, we note that among the possible transformations of $\mathcal{S}_0$ we can consider that induced by

$$q \longrightarrow q^v = v(q) = p, \quad (3.17)$$

with p the momentum. According to (2.37) we have

$$\mathcal{S}_0(q) \longrightarrow \mathcal{S}_0^v(p) = \mathcal{S}_0(q(p)). \quad (3.18)$$

In order to understand the structure of the function $\mathcal{S}_0^v(p)$, we first observe that there is a natural way to introduce the tools we need in going towards a formulation with manifest p - q duality. Let us introduce the dual reduced action $\mathcal{T}_0(p)$ defined as the Legendre transform of $\mathcal{S}_0(q)$

$$\mathcal{T}_0 = q \frac{\partial \mathcal{S}_0}{\partial q} - \mathcal{S}_0, \quad \mathcal{S}_0 = p \frac{\partial \mathcal{T}_0}{\partial p} - \mathcal{T}_0, \quad (3.19)$$

⁵In this respect we note that according to the Kochen–Specker theorem [29] it is not possible to preserve functional relations between physical quantities with assigned values. We note that this topic, related to hidden variables [30][31], has been recently considered in [32].

$$p = \frac{\partial \mathcal{S}_0}{\partial q}, \quad q = \frac{\partial \mathcal{T}_0}{\partial p}. \quad (3.20)$$

It follows that the transformation (3.18) is nothing but (3.19) seen as function of p

$$\mathcal{S}_0^v(p) = p \frac{\partial \mathcal{T}_0(p)}{\partial p} - \mathcal{T}_0(p). \quad (3.21)$$

Above we discussed the dependence of the structure of some relevant equations on the canonical variables considered either as dependent or independent and noticed that this issue may be connected to the fact that in QM position and momentum cannot be simultaneously measured. In this respect it is interesting to note that for the pair $\mathcal{S}_0$ – $\mathcal{T}_0$, in the case in which p and q are considered as independent variables, we have $F(q, p) = qp - \mathcal{T}_0(p) \leq F(q, p(q)) = \mathcal{S}_0(q)$, $\forall q, p$, with $p = p(q)$ given by $\partial_p \mathcal{T}_0(p) = q$. Therefore, considering p and q as independent variables we have the Young inequality (see *e.g.* [33])

$$\mathcal{T}_0(p) + \mathcal{S}_0(q) \geq qp, \quad \forall q, p. \quad (3.22)$$

3.3 The dual HJ equation

Despite $\mathcal{T}_0$ is not usually considered in the literature, the involutive character of the Legendre transformation makes it clear that $\mathcal{T}_0$ is a natural function to consider in looking for a formulation with manifest p – q duality. Let us introduce the Legendre transform of the Hamilton principal function $\mathcal{S}$

$$\mathcal{T} = q \frac{\partial \mathcal{S}}{\partial q} - \mathcal{S}, \quad \mathcal{S} = p \frac{\partial \mathcal{T}}{\partial p} - \mathcal{T}, \quad (3.23)$$

$$p = \frac{\partial \mathcal{S}}{\partial q}, \quad q = \frac{\partial \mathcal{T}}{\partial p}. \quad (3.24)$$

Observe that in the stationary case

$$\mathcal{S}(q, t) = \mathcal{S}_0(q) - Et, \quad \mathcal{T}(p, t) = \mathcal{T}_0(p) + Et. \quad (3.25)$$

A first question to investigate concerns of the classical HJ equation in the $\mathcal{T}^{cl}$ –representation. Let us consider the differentials

$$d\mathcal{S} = \frac{\partial \mathcal{S}}{\partial q} dq + \frac{\partial \mathcal{S}}{\partial t} dt = pdq + \frac{\partial \mathcal{S}}{\partial t} dt, \quad (3.26)$$

$$d\mathcal{T} = \frac{\partial \mathcal{T}}{\partial p} dp + \frac{\partial \mathcal{T}}{\partial t} dt = qdp + \frac{\partial \mathcal{T}}{\partial t} dt, \quad (3.27)$$

which imply

$$d\mathcal{S} = d(pq - \mathcal{T}) = pdq + qdp - qdp - \frac{\partial \mathcal{T}}{\partial t} dt, \quad (3.28)$$

that is

$$\frac{\partial \mathcal{S}}{\partial t} = -\frac{\partial \mathcal{T}}{\partial t}. \quad (3.29)$$

Relevance of this equation resides in the fact it connects the $\mathcal{S}$ and $\mathcal{T}$ pictures through time evolution. For example, we will see that Eq.(3.29) will fix a sign ambiguity in determining the “self–dual states”.

Furthermore, observe that in the classical case it is precisely the partial derivative of $\mathcal{S}^{cl}$ with respect to time that appears in the HJ equation. In particular, it follows from (2.10) and (3.29) that $\mathcal{T}^{cl}$ satisfies the following dual version of the classical HJ equation

$$H\left(q = \frac{\partial \mathcal{T}^{cl}}{\partial p}, p, t\right) - \frac{\partial \mathcal{T}^{cl}}{\partial t} = 0, \quad (3.30)$$

which for general Hamiltonians

$$H = \frac{p^2}{2m} + V(q, p, t), \quad (3.31)$$

becomes

$$\frac{1}{2m}p^2 + V\left(q = \frac{\partial \mathcal{T}^{cl}}{\partial p}, p, t\right) - \frac{\partial \mathcal{T}^{cl}}{\partial t} = 0. \quad (3.32)$$

We can derive an equivalent version of (3.32) with manifest dual structure. In particular, by analogy with the HJ equation, we may impose that its dual formulation contains the “*dual kinetic term*”

$$\frac{m}{2}\alpha^2 q^2. \quad (3.33)$$

The request of equivalence between the dual equations defines a potential $U^{(\alpha)}$ dual to V , namely

$$\frac{m\alpha^2}{2}\left(\frac{\partial \mathcal{T}^{cl}}{\partial p}\right)^2 + U^{(\alpha)}\left(p, q = \frac{\partial \mathcal{T}^{cl}}{\partial p}, t\right) + \frac{\partial \mathcal{T}^{cl}}{\partial t} = 0, \quad (3.34)$$

where by (3.32)

$$U^{(\alpha)}(p, q, t) = -\frac{1}{2m}(p^2 + m^2\alpha^2 q^2) - V(q, p, t), \quad (3.35)$$

and α is a constant with the dimension of a frequency. Associated with $\mathcal{T}^{cl}$ there is the “Hamiltonian” $K(p, q, t)$, dual to $H(q, p, t)$

$$K(p, q, t) = \frac{m\alpha^2}{2}q^2 + U^{(\alpha)}(p, q, t), \quad (3.36)$$

which has the simple relation with H

$$K(p, q, t) = -H(q, p, t). \quad (3.37)$$

It follows that the classical equations of motion are equivalent to

$$\dot{q} = -\frac{\partial K}{\partial p}, \quad \dot{p} = \frac{\partial K}{\partial q}, \quad (3.38)$$

and the dual classical HJ equation can be written as

$$K\left(p, q = \frac{\partial \mathcal{T}^{cl}}{\partial p}, t\right) + \frac{\partial \mathcal{T}^{cl}}{\partial t} = 0. \quad (3.39)$$

There is a particular system in which the $\mathcal{S}^{cl}$ and $\mathcal{T}^{cl}$ pictures have basically the same HJ equation. This happens for the harmonic oscillator

$$H_\omega(q, p) = \frac{1}{2m}p^2 + V_\omega(q), \quad (3.40)$$

where $V_\omega(q) = m\omega^2 q^2/2$. By (3.35) we have

$$U_\omega^{(\alpha)} = -\frac{1}{2m}p^2 - \frac{m(\omega^2 + \alpha^2)}{2}q^2, \quad (3.41)$$

so that $U_\omega^{(\alpha)} = -H_{\omega_D}$, where the frequency of the new oscillator is $\omega_D^2 = \omega^2 + \alpha^2$. It follows that for $\alpha = \pm i\omega$, the dual potential corresponds to minus the free particle Hamiltonian

$$U_\omega^{(\pm i\omega)} = -H_0 = -\frac{1}{2m}p^2. \quad (3.42)$$

We also observe that as $H_\omega(q, p) = H_\omega(p/m\omega, m\omega q)$, we have

$$K_\omega(p, q) = -H_\omega(p/m\omega, m\omega q). \quad (3.43)$$

4 Möbius symmetry and the canonical equation

In this section we will first consider the effect of the VT on $\mathcal{T}_0$. Next, we will observe that by (2.37), it follows by construction that the Legendre transform of $\mathcal{T}_0$ is invariant under VTs. We will see that a $GL(2, \mathbb{C})$ -transformation corresponds to a rotation of the two-dimensional kernel of a second-order linear operator which we will determine by considering the second derivative of the Legendre transform of $\mathcal{T}_0$ with respect to $\mathbf{s} = \mathcal{S}_0(q)$. As a consequence, the potential $\mathcal{U}(\mathbf{s})$ appearing in this operator turns out to be invariant under the VT corresponding to the Möbius group. In this context covariance of the relevant second-order linear differential equation follows by consistency.

4.1 $\mathcal{T}_0$ -transformation induced by v

Since $\mathcal{S}_0$ and $\mathcal{T}_0$ are a Legendre pair, specifying the transformation properties of $\mathcal{S}_0$ will fix the transformation of $\mathcal{T}_0$ and vice versa. In particular, consistency implies that if $\mathcal{S}_0(q) \longrightarrow \tilde{\mathcal{S}}_0(\tilde{q})$ is an arbitrary transformation, not necessarily of v -type, that is in general $\tilde{\mathcal{S}}_0(\tilde{q}) \neq \mathcal{S}_0(q)$, then the new dual reduced action will be the Legendre transform of $\tilde{\mathcal{S}}_0(\tilde{q})$. In other words, the transformation $\mathcal{S}_0(q) \longrightarrow \tilde{\mathcal{S}}_0(\tilde{q})$ induces the following transformation of $\mathcal{T}_0(p)$

$$\mathcal{T}_0(p) \longrightarrow \tilde{\mathcal{T}}_0(\tilde{p}) = \tilde{q}\tilde{p} - \tilde{\mathcal{S}}_0(\tilde{q}). \quad (4.1)$$

In the case in which the transformation of the functional structure of $\mathcal{S}_0$ is performed by a v -map we have $p_v = (\partial_q q^v)^{-1}p$, and replacing $\mathcal{S}_0^v(q^v)$ with $pq - \mathcal{T}_0$ in (4.1) yields

$$\delta_v \mathcal{T}_0 = \delta_v(qp), \quad (4.2)$$

where $\delta_v \mathcal{T}_0 \equiv \mathcal{T}_{0v}(p_v) - \mathcal{T}_0(p)$ and $\delta_v(qp) \equiv q^v p_v - qp$. Observe that by Eq.(2.37), the VTs can be equivalently seen as a symmetry of the Legendre transform of $\mathcal{T}_0$.

4.2 The canonical equation

Let us now consider the VT corresponding to the Möbius transformation

$$q \longrightarrow q^v = \frac{Aq + B}{Cq + D}, \quad (4.3)$$

that due to the dimensionality of p and q , we consider Eq.(4.3) as a $GL(2, \mathbb{C})$ -transformation rather than a $PGL(2, \mathbb{C}) \cong PSL(2, \mathbb{C})$ -transformation. Eqs.(2.37) and (4.3) imply

$$p \longrightarrow p_v = \rho^{-1}(Cq + D)^2 p, \quad (4.4)$$

where $\rho \equiv AD - BC$. Observe that ρ has the dimension of a length times the dimension of A^2 so that ρ would be dimensionless if $Dim[A] = length^{-1/2}$. By (4.2)(4.3) and (4.4) it follows that the action on $\mathcal{T}_0$ induced by the v -maps is

$$\delta_v \mathcal{T}_0 = \rho^{-1}(ACq^2 + 2BCq + BD)p. \quad (4.5)$$

Taking the square root of (4.4), we see that ($\epsilon = \pm 1$)

$$q^v \sqrt{p_v} = \epsilon \sqrt{\rho^{-1}}(Aq\sqrt{p} + B\sqrt{p}), \quad \sqrt{p_v} = \epsilon \sqrt{\rho^{-1}}(Cq\sqrt{p} + D\sqrt{p}). \quad (4.6)$$

This version of Eqs.(4.3) and (4.4) indicates that $q^v \sqrt{p_v}$ and $\sqrt{p_v}$ are linear combinations of the solutions of a second-order linear differential equation. In particular, Eq.(4.6) can be seen as a rotation of the elements in the kernel of a second-order linear operator which can be simply identified by observing that the second derivative of the second equation in (3.19) with respect to $\mathbf{s} = \mathcal{S}_0(q)$

$$\frac{\partial q}{\partial \mathbf{s}} \frac{\partial p}{\partial \mathbf{s}} + p \frac{\partial^2 q}{\partial \mathbf{s}^2} = 0, \quad (4.7)$$

is equivalent to

$$\frac{1}{q\sqrt{p}} \frac{\partial^2 (q\sqrt{p})}{\partial \mathbf{s}^2} = \frac{1}{\sqrt{p}} \frac{\partial^2 \sqrt{p}}{\partial \mathbf{s}^2}. \quad (4.8)$$

On the other hand, this is equivalent to the “canonical equation”

$$\left(\frac{\partial^2}{\partial \mathbf{s}^2} + \mathcal{U}(\mathbf{s}) \right) q\sqrt{p} = 0 = \left(\frac{\partial^2}{\partial \mathbf{s}^2} + \mathcal{U}(\mathbf{s}) \right) \sqrt{p}, \quad (4.9)$$

where $\mathcal{U}$ is the “canonical potential”

$$\mathcal{U}(\mathbf{s}) = \frac{1}{2} \{q\sqrt{p}/\sqrt{p}, \mathbf{s}\} = \frac{1}{2} \{q, \mathbf{s}\}, \quad (4.10)$$

with

$$\{h(x), x\} = \frac{h'''}{h'} - \frac{3}{2} \left(\frac{h''}{h'} \right)^2 = (\ln h')'' - \frac{1}{2} (\ln h')'^2, \quad (4.11)$$

denoting the Schwarzian derivative whose chain rule is

$$\{h(x), x(y)\} = \left(\frac{\partial y}{\partial x} \right)^2 \{h(x), y\} - \left(\frac{\partial y}{\partial x} \right)^2 \{x, y\}. \quad (4.12)$$

4.3 Legendre transformation as Möbius transformation

Another interesting property due to (2.37) is that $\mathcal{S}_0$ can be expressed in terms of a Möbius transformation of $\mathcal{T}_0$. Let us first consider the effect of an S -transformation

$$S = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}. \quad (4.13)$$

We have by (4.5)

$$S \circ \mathcal{T}_0 = \mathcal{T}_0 - 2pq, \quad (4.14)$$

so that the Legendre transform of $\mathcal{T}_0$ can be written in the form

$$\mathcal{S}_0(q) = -\frac{1}{2}(\mathcal{T}_0(p) + S \circ \mathcal{T}_0(p)). \quad (4.15)$$

Observe that from (4.5) it follows that for any B and C , $BC \neq 0$, the action of $R = \begin{pmatrix} 0 & B \\ C & 0 \end{pmatrix}$ on $\mathcal{T}_0$ does not depend on B and C . However, by (4.3) the action of R on q depends on $\delta = B/C$

$$R : q \longrightarrow q^v = v(q) = \frac{\delta}{q}, \quad (4.16)$$

whereas by (4.4) the action on p is

$$R : p \longrightarrow p_v = -\frac{1}{\delta}q^2p. \quad (4.17)$$

In particular, we have

$$R : pq \longrightarrow p_v q^v = -pq. \quad (4.18)$$

4.4 Möbius invariance of $\mathcal{U}$

Eqs.(4.9) and (4.10) follow from (4.8) or, equivalently, from the observation that the identities

$$\frac{\partial}{\partial x} h'^{1/2} h'^{-1/2} = 0 = \frac{\partial}{\partial x} \frac{1}{h'} \frac{\partial}{\partial x} h'^{1/2} h'^{-1/2} h, \quad (4.19)$$

imply that the kernel of the second-order linear operator

$$h'^{1/2} \frac{\partial}{\partial x} \frac{1}{h'} \frac{\partial}{\partial x} h'^{1/2} = \frac{\partial^2}{\partial x^2} + \frac{1}{2}\{h, x\}, \quad (4.20)$$

is given by the linear span of $h'^{-1/2}$ and $h'^{-1/2}h$, that is

$$\left(\frac{\partial^2}{\partial x^2} + \frac{1}{2}\{h, x\} \right) h'^{-1/2}(Ah + B) = 0. \quad (4.21)$$

Eqs.(4.19)–(4.21) show that the Schwarzian derivative of the ratio of two linearly independent elements in the kernel of a second-order linear operator $(\partial_x^2 + V(x))$ is twice $V(x)$. Noticing that for any A

and B , not simultaneously vanishing, $(\partial_x^2 + V(x))f_k(x) = 0$, $k = 1, 2$, is equivalent to $V = -(Af_1'' + Bf_2'')/(Af_1 + Bf_2)$, we have the well-known fact

$$\{\gamma(h), x\} = \{h, x\}, \quad (4.22)$$

where $\gamma(h)$ is the Möbius transformation of h

$$\gamma(h) = \frac{Ah + B}{Ch + D}. \quad (4.23)$$

Eq.(4.22) implies $\{\gamma(x), x\} = \{x, x\} = 0$. Conversely, if $\{h, x\} = 0$, then, solving

$$(\ln h')'' - \frac{1}{2}(\ln h')'^2 = 0, \quad (4.24)$$

yields $h = \gamma(x)$. By (4.12) these properties of the Schwarzian derivative are equivalent to the fact that

$$\{f, x\} = \{h, x\}, \quad (4.25)$$

if and only if $f = \gamma(h)$. This implies that $\mathcal{U}(\mathbf{s})$, associated to the Legendre transform of $\mathcal{T}_0$, is invariant under v -maps corresponding to the Möbius transformations $q^v = \gamma(q)$, that is

$$\mathcal{U}^v(\mathbf{s}^v) = \frac{1}{2}\{q^v, \mathbf{s}^v\} = \frac{1}{2}\{\gamma(q), \mathbf{s}\} = \mathcal{U}(\mathbf{s}), \quad (4.26)$$

where $\mathbf{s}^v = \mathcal{S}_0^v(q^v) = \mathcal{S}_0(q) = \mathbf{s}$. However, if q^v is not a Möbius transformation of q , then

$$\mathcal{U}^v(\mathbf{s}^v) = \frac{1}{2}\{q^v, \mathbf{s}^v\} \neq \mathcal{U}(\mathbf{s}). \quad (4.27)$$

Therefore, although by definition any v -map leaves the Legendre transform of $\mathcal{T}_0$ invariant, we have that $\mathcal{U}$ is invariant only under $GL(2, \mathbb{C})$ Möbius transformations of q .

4.5 Canonical equation, dynamics and initial conditions

We have seen that the canonical equation associated with the Legendre transform of $\mathcal{T}_0$ is $GL(2, \mathbb{C})$ -invariant. Another property of the canonical equation (4.9) is that it can be seen as an equation of motion. In fact, suppose that the canonical potential $\mathcal{U}(\mathbf{s})$ is given. Then, if $y_1(\mathbf{s})$ and $y_2(\mathbf{s})$ are two linearly independent solutions of the canonical equation, we have

$$q\sqrt{p} = Ay_1(\mathbf{s}) + By_2(\mathbf{s}), \quad \sqrt{p} = Cy_1(\mathbf{s}) + Dy_2(\mathbf{s}). \quad (4.28)$$

In order to derive $\mathcal{S}_0(q)$, that is to solve the dynamical problem, we first use (4.28) to obtain q as function of $\mathbf{s}$ and then consider its inverse. Taking the ratio of (4.28) yields

$$q = \frac{Ah(\mathbf{s}) + B}{Ch(\mathbf{s}) + D}, \quad (4.29)$$

where $h(\mathbf{s}) = y_1(\mathbf{s})/y_2(\mathbf{s})$. Inverting (4.29) we have $\mathcal{S}_0 = h^{-1} \circ \gamma^{-1}$ with γ given in (4.23) so that the dynamical problem is solved by

$$\mathcal{S}_0(q) = h^{-1}(\gamma^{-1}(q)). \quad (4.30)$$

Although the canonical equation is a second-order linear differential equation, so that a particular solution is obtained by giving two conditions, we have that as $q(\mathbf{s})$ is the ratio of two linearly independent solutions, one needs three initial conditions to fix it. This follows from the observation that (4.29) can be written in terms of three constants; *e.g.* if $A \neq 0$ and $C \neq 0$, then

$$q = \frac{A}{C} \left(\frac{h(\mathbf{s}) + B/A}{h(\mathbf{s}) + D/C} \right). \quad (4.31)$$

Therefore, the dynamics is obtained by first considering the ratio of two linearly independent solutions of the canonical equation (4.9) and then evaluating its inverse. In particular, all the reduced actions $\mathcal{S}_0^v$ in the $GL(2, \mathbb{C})$ -orbit of $\mathcal{S}_0$ are the inverse of the ratio of two linearly independent solutions of the same canonical equation.

4.6 Covariance

A property of the canonical equation is that it is manifestly covariant under arbitrary transformations. In particular, we have seen that the transformation of $\mathcal{T}_0$, induced by an arbitrary transformation of $\mathcal{S}_0$, not necessarily of v -type, is determined by the fact that $\tilde{\mathcal{T}}_0$ is the Legendre transform of $\tilde{\mathcal{S}}_0$. Then, by Eq.(4.1) the canonical equation (4.9) is covariant under arbitrary transformations. In particular, the second derivative of (4.1) with respect to $\tilde{\mathbf{s}} = \tilde{\mathcal{S}}_0(\tilde{q})$ yields

$$\left(\frac{\partial^2}{\partial \tilde{\mathbf{s}}^2} + \tilde{\mathcal{U}}(\tilde{\mathbf{s}}) \right) \tilde{q} \sqrt{\tilde{p}} = 0 = \left(\frac{\partial^2}{\partial \tilde{\mathbf{s}}^2} + \tilde{\mathcal{U}}(\tilde{\mathbf{s}}) \right) \sqrt{\tilde{p}}. \quad (4.32)$$

Thus, the procedure to associate a second-order linear differential equation with a Legendre transformation is manifestly covariant. There is another property of $\mathcal{U}(\mathbf{s})$. Namely, by the chain rule of the Schwarzian derivative (4.12), we have

$$\{q, (A\mathbf{s} + B)/(C\mathbf{s} + D)\} = \tau^{-2}(C\mathbf{s} + D)^4 \{q, \mathbf{s}\} = 2\tau^{-2}(C\mathbf{s} + D)^4 \mathcal{U}(\mathbf{s}), \quad (4.33)$$

where $\tau \equiv AD - BC \neq 0$. Hence, a Möbius transformation of $\mathcal{S}_0$ corresponds to a rescaling of $\mathcal{U}(\mathbf{s})$. Under such a transformation we have

$$\tilde{q} = q, \quad \tilde{p} = \frac{\partial \tilde{\mathbf{s}}}{\partial \tilde{q}} = \frac{\partial \tilde{\mathbf{s}}}{\partial \mathbf{s}} p, \quad (4.34)$$

so that, setting $\phi(\mathbf{s}) = aq\sqrt{p} + b\sqrt{p}$, with a and b two non simultaneously vanishing constants, we have $\tilde{\phi}(\tilde{\mathbf{s}}) = (\partial_{\tilde{\mathbf{s}}} \tilde{\mathbf{s}})^{1/2} \phi(\mathbf{s})$. Therefore, ϕ transforms as a $-1/2$ -differential. Observe that

$$\frac{\partial^2}{\partial \tilde{\mathbf{s}}^2} = \left(\frac{\partial \tilde{\mathbf{s}}}{\partial \mathbf{s}} \right)^{-2} \frac{\partial^2}{\partial \mathbf{s}^2} - \left(\frac{\partial \tilde{\mathbf{s}}}{\partial \mathbf{s}} \right)^{-3} \left(\frac{\partial^2 \tilde{\mathbf{s}}}{\partial \mathbf{s}^2} \right) \frac{\partial}{\partial \mathbf{s}}, \quad (4.35)$$

so that under the transformation $\mathbf{s} \longrightarrow \tilde{\mathbf{s}} = (A\mathbf{s} + B)/(C\mathbf{s} + D)$, we have

$$\frac{\partial^2}{\partial \tilde{\mathbf{s}}^2} = \tau^{-2}(C\mathbf{s} + D)^4 \left(\frac{\partial^2}{\partial \mathbf{s}^2} + \frac{2C}{C\mathbf{s} + D} \frac{\partial}{\partial \mathbf{s}} \right). \quad (4.36)$$

Hence, by (4.33)

$$\begin{aligned} \left(\frac{\partial^2}{\partial \tilde{\mathbf{s}}^2} + \tilde{\mathcal{U}}(\tilde{\mathbf{s}}) \right) \tilde{\phi}(\tilde{\mathbf{s}}) &= \tau^{-2}(C\mathbf{s} + D)^4 \left(\frac{\partial^2}{\partial \mathbf{s}^2} + \frac{2C}{C\mathbf{s} + D} \frac{\partial}{\partial \mathbf{s}} + \mathcal{U}(\mathbf{s}) \right) \tilde{\phi}(\tilde{\mathbf{s}}) = \\ &= \tau^{-3/2}(C\mathbf{s} + D)^3 \left(\frac{\partial^2}{\partial \mathbf{s}^2} + \mathcal{U}(\mathbf{s}) \right) \phi(\mathbf{s}) = 0. \end{aligned} \quad (4.37)$$

Comparing the VTs corresponding to Möbius transformations

$$q \longrightarrow \frac{Aq + B}{Cq + D}, \quad \mathcal{S}_0 \longrightarrow \mathcal{S}_0, \quad p \longrightarrow \rho^{-1}(Cq + D)^2 p, \quad (4.38)$$

with the above transformations of $\mathbf{s}$

$$\mathcal{S}_0 \longrightarrow \frac{A\mathcal{S}_0 + B}{C\mathcal{S}_0 + D}, \quad q \longrightarrow q, \quad p \longrightarrow \tau(C\mathcal{S}_0 + D)^{-2} p. \quad (4.39)$$

we see that there is a sort of duality between q and $\mathbf{s}$. In particular, Eq.(4.39) is nothing but the analogue of the VTs corresponding to the Möbius transformations, with the role of $\mathbf{s}$ and q interchanged. This duality between $\mathbf{s}$ and q suggests considering the analogue of the canonical equation but with the role of $\mathbf{s}$ and q interchanged. In this case one has

$$\left(\frac{\partial^2}{\partial q^2} + \frac{1}{2}\{\mathcal{S}_0, q\} \right) \mathcal{S}_0 p^{-1/2} = 0 = \left(\frac{\partial^2}{\partial q^2} + \frac{1}{2}\{\mathcal{S}_0, q\} \right) p^{-1/2}. \quad (4.40)$$

We will see that $\{\mathcal{S}_0, q\}$ is proportional to the quantum potential. Generalizing the above Möbius transformations to arbitrary ones

$$\mathbf{s} \longrightarrow \tilde{\mathbf{s}}, \quad (4.41)$$

we obtain

$$\frac{\partial^2}{\partial \tilde{\mathbf{s}}^2} \left(\frac{\partial \tilde{\mathbf{s}}}{\partial \mathbf{s}} \right)^{1/2} = \left(\frac{\partial \tilde{\mathbf{s}}}{\partial \mathbf{s}} \right)^{-3/2} \left(\frac{\partial^2}{\partial \mathbf{s}^2} + \frac{1}{2}\{\tilde{\mathbf{s}}, \mathbf{s}\} \right), \quad (4.42)$$

where now the $\{\tilde{\mathbf{s}}, \mathbf{s}\}$ does not vanish. In the case in which the functional change of q is given by

$$q \longrightarrow \tilde{q}(\tilde{\mathbf{s}}) = q(\mathbf{s}), \quad (4.43)$$

we have by $\tilde{\mathcal{U}}(\tilde{\mathbf{s}}) = \{\tilde{q}, \tilde{\mathbf{s}}\}/2 = \{q, \tilde{\mathbf{s}}\}/2$, that is

$$0 = \left(\frac{\partial \tilde{\mathbf{s}}}{\partial \mathbf{s}} \right)^{3/2} \left(\frac{\partial^2}{\partial \tilde{\mathbf{s}}^2} + \frac{1}{2}\{q, \tilde{\mathbf{s}}\} \right) \tilde{\phi}(\tilde{\mathbf{s}}) = \left(\frac{\partial^2}{\partial \mathbf{s}^2} + \frac{1}{2}\{\tilde{\mathbf{s}}, \mathbf{s}\} + \frac{1}{2}(\{q, \mathbf{s}\} - \{\tilde{\mathbf{s}}, \mathbf{s}\}) \right) \phi(\mathbf{s}), \quad (4.44)$$

that is the equation $(\partial_{\tilde{\mathbf{s}}}^2 + \frac{1}{2}\{\tilde{q}, \tilde{\mathbf{s}}\}) \tilde{\phi}(\tilde{\mathbf{s}}) = 0$ is equivalent to

$$\left(\frac{\partial^2}{\partial \mathbf{s}^2} + \frac{1}{2}\{q, \mathbf{s}\} \right) \phi(\mathbf{s}) = 0. \quad (4.45)$$

The above analysis shows that though $\mathcal{U}$ transforms as the Schwarzian derivative, its additive term is canceled by that coming from $\partial_{\tilde{\mathbf{s}}}^2$ acting on $(\partial \tilde{\mathbf{s}}/\partial \mathbf{s})^{1/2}$.

5 Legendre duality

The involutive nature of the Legendre transformation implies that the construction considered in the previous section admits a dual formulation obtained by the exchange

$$\mathcal{S}_0 \longleftrightarrow \mathcal{T}_0, \quad q \longleftrightarrow p. \quad (5.1)$$

In particular, we will see how both the VTs and the canonical equation have a dual version. The correspondence manifests the equivalence of the q and p descriptions and shows how this duality relies on the properties of the Legendre transformation.

5.1 u -transformations

Let us start by considering a locally invertible momentum transformation

$$p \longrightarrow p^u = u(p). \quad (5.2)$$

Similarly to the transformation properties of the reduced action $\mathcal{S}_0$, we can associate in a natural way a new $\mathcal{T}_0$ with the u -map $\mathcal{T}_0 \longrightarrow \mathcal{T}_0^u = \mathcal{T}_0 \circ u^{-1}$, which is equivalent to

$$\mathcal{T}_0^u(p^u) = \mathcal{T}_0(p(p^u)). \quad (5.3)$$

Since any pair $\mathcal{T}_0$ and $\mathcal{T}_0^u$ induces the p -transformation defined by (5.3), we can consider the previous construction in the reversed order, *i.e.*

$$p \longrightarrow p^u = \mathcal{T}_0^{u^{-1}} \circ \mathcal{T}_0(p). \quad (5.4)$$

Since $q \longrightarrow q_u = \partial_{p^u} \mathcal{T}_0^u(p^u)$, under (5.2) q transforms as ∂_p , that is $q_u = (\partial_p p^u)^{-1} q$. In summary, under VTs we have

$$q^v = v(q), \quad \mathcal{S}_0^v(q^v) = \mathcal{S}_0(q), \quad \delta_v \mathcal{T}_0 = \delta_v(qp), \quad (5.5)$$

$$p \sim \frac{\partial}{\partial q}, \quad (5.6)$$

where $\delta_v \mathcal{T}_0 \equiv \mathcal{T}_{0v}(p_v) - \mathcal{T}_0(p)$ and $\delta_v(qp) \equiv q^v p_v - qp$. Similarly, for the u -transformations

$$p^u = u(p), \quad \mathcal{T}_0^u(p^u) = \mathcal{T}_0(p), \quad \delta_u \mathcal{S}_0 = \delta_u(pq), \quad (5.7)$$

$$q \sim \frac{\partial}{\partial p}, \quad (5.8)$$

where $\delta_u \mathcal{S}_0 \equiv \mathcal{S}_{0u}(q_u) - \mathcal{S}_0(q)$ and $\delta_u(pq) \equiv p^u q_u - pq$.

5.2 The dual canonical equation

Let us now consider the $GL(2, \mathbb{C})$ -transformations

$$p \longrightarrow p^u = \frac{Ap + B}{Cp + D}, \quad (5.9)$$

$\sigma \equiv AD - BC$. Since $q_u = \partial_{p^u} \mathcal{T}_0^u(p^u) = \partial_{p^u} \mathcal{T}_0(p)$, we have that the effect on q of (5.9) is

$$q \longrightarrow q_u = \sigma^{-1}(Cp + D)^2 q. \quad (5.10)$$

Eqs.(5.9)(5.10) are equivalent to

$$p^u \sqrt{q_u} = \epsilon \sqrt{\sigma^{-1}}(Ap\sqrt{q} + B\sqrt{q}), \quad \sqrt{q_u} = \epsilon \sqrt{\sigma^{-1}}(Cp\sqrt{q} + D\sqrt{q}), \quad (5.11)$$

where $\epsilon = \pm 1$. As $\delta_u \mathcal{S}_0 = \delta_u(pq)$ we have

$$\delta_u \mathcal{S}_0 = \sigma^{-1}(ACp^2 + 2BCp + BD)q. \quad (5.12)$$

Let us recall that the canonical equation (4.9) arises from the identity (4.8). This identity can be also obtained by simply taking the second derivative of $p = (\partial_{\mathbf{s}} q)^{-1}$ with respect to $\mathbf{s}$. Thus we see that the canonical equation stems from the very basic definition of dual variable $p = \partial_q \mathcal{S}_0$, thereby capturing the germ of p - q duality. In this context the fact that $\mathcal{S}_0$ and $\mathcal{T}_0$ are the Legendre transform of each other, leads to the dual version of the canonical equation (4.9)

$$\left(\frac{\partial^2}{\partial \mathbf{t}^2} + \mathcal{V}(\mathbf{t}) \right) p\sqrt{q} = 0 = \left(\frac{\partial^2}{\partial \mathbf{t}^2} + \mathcal{V}(\mathbf{t}) \right) \sqrt{q}, \quad (5.13)$$

where $\mathbf{t} = \mathcal{T}_0(p)$, and

$$\mathcal{V}(\mathbf{t}) = \frac{1}{2} \{p\sqrt{q}/\sqrt{q}, \mathbf{t}\} = \frac{1}{2} \{p, \mathbf{t}\}. \quad (5.14)$$

5.3 Legendre brackets

We now show, following [11][1], that there are natural brackets associated to the relationship between the Legendre transformation and second-order linear differential equations observed in [6]. Let us start by considering a function $U(z)$ and its Legendre transform

$$V(w) = z\partial_z U(z) - U(z), \quad U(z) = w\partial_w V(w) - V(w) \quad (5.15)$$

where $w = \partial_z U$ (note that $z = \partial_w V$). Taking the second derivative of the second equation in (5.15) with respect to $u = U(z)$, we see that $\sqrt{\partial_z U}$ and $z\sqrt{\partial_z U}$ correspond to linearly independent solutions of a second-order linear differential equation. Let us now set

$$\tau = \frac{\partial z \sqrt{w}}{\partial \sqrt{w}} = z + 2 \frac{w}{\partial_z w} = \frac{1}{2} \frac{\partial^2 V}{\partial \sqrt{w}^2}. \quad (5.16)$$

The “Legendre brackets” introduced in [11] are

$$\{X, Y\}_{(u)} = (\partial_u \tau)^{-1} \left(\frac{\partial X}{\partial \sqrt{w}} \frac{\partial}{\partial u} \frac{\partial Y}{\partial \sqrt{w}} - \frac{\partial Y}{\partial \sqrt{w}} \frac{\partial}{\partial u} \frac{\partial X}{\partial \sqrt{w}} \right). \quad (5.17)$$

In the case in which the generating function is $\mathcal{S}_0(q)$, we have that the linearly independent solutions of the canonical equation satisfy

$$\{\sqrt{p}, \sqrt{p}\}_{(\mathbf{s})} = 0 = \{q\sqrt{p}, q\sqrt{p}\}_{(\mathbf{s})}, \quad \{\sqrt{p}, q\sqrt{p}\}_{(\mathbf{s})} = 1, \quad (5.18)$$

where $\mathbf{s} = \mathcal{S}_0(q)$. Similarly, starting from the generating function $\mathcal{T}_0(p)$ we have

$$\{\sqrt{q}, \sqrt{q}\}_{(\mathbf{t})} = 0 = \{p\sqrt{q}, p\sqrt{q}\}_{(\mathbf{t})}, \quad \{\sqrt{q}, p\sqrt{q}\}_{(\mathbf{t})} = 1. \quad (5.19)$$

5.4 p and q as Legendre pair

We have seen that the Legendre transformation is a natural framework to investigate the structure of dualities. In particular, with the “Legendre pair” $\mathcal{S}_0$ – $\mathcal{T}_0$ there is a naturally associated pair of second-order linear differential equations. We now consider the Legendre pair $\mathcal{S}_0$ – $\mathcal{T}_0$ from a slightly different point of view. First of all observe that $\mathcal{S}_0$ and $\mathcal{T}_0$ define the function $\mathbf{t} = f(\mathbf{s})$, where $\mathbf{s} = \mathcal{S}_0(q)$ and $\mathbf{t} = \mathcal{T}_0(p)$. Then suppose that instead of deriving $\mathcal{T}_0$ from $\mathcal{S}_0$ we started from a given $\mathbf{t} = f(\mathbf{s})$. In this case, we can define the dual pair p – q by requiring that

$$\mathcal{T}_0 = q \frac{\partial \mathcal{S}_0}{\partial q} - \mathcal{S}_0, \quad \mathcal{S}_0 = p \frac{\partial \mathcal{T}_0}{\partial p} - \mathcal{T}_0. \quad (5.20)$$

Similarly, we can introduce a new pair x^D – x by considering p as the Legendre dual of q , that is

$$\nu p = x \frac{\partial q}{\partial x} - q, \quad \nu^{-1} q = x^D \frac{\partial p}{\partial x^D} - p, \quad (5.21)$$

where ν is a constant which has been introduced for dimensional reasons. Therefore, considering p and q as Legendre duals induces the definition of the new pair x^D – x . Integrating (5.21) we obtain

$$x = x^0 \exp\left[\int_{q^0}^q dz (z + \nu \partial_z \mathcal{S}_0(z))^{-1}\right], \quad (5.22)$$

and

$$x^D = x^{D0} (q + \nu \partial_q \mathcal{S}_0(q)) \exp\left[-\int_{q^0}^q dz (z + \nu \partial_z \mathcal{S}_0(z))^{-1}\right], \quad (5.23)$$

where $x^0 \equiv x(q^0)$, $x^{D0} \equiv x^D(q^0)$. Regarding p – q duality as Legendre duality suggests considering a sequence of dual variables. In particular, we can consider also the x^D – x pair as Legendre duals. This defines the pair y^D – y by

$$\mu x^D = y \frac{\partial x}{\partial y} - x, \quad \mu^{-1} x = y^D \frac{\partial x^D}{\partial y^D} - x^D, \quad (5.24)$$

where μ is a dimensional constant. Integrating (5.24) yields

$$y = y^0 \exp\left[\int_{x^0}^x dz(z + \mu\partial_z q(z))^{-1}\right], \quad (5.25)$$

and

$$y^D = y^{D^0}(x + \mu\partial_x q(x)) \exp\left[-\int_{x^0}^x dz(z + \mu\partial_z q(z))^{-1}\right], \quad (5.26)$$

where $y^0 \equiv y(x^0)$, $y^{D^0} \equiv y^D(x^0)$. Iteration of this construction leads to the “ $\mathcal{W}$ -sequence”

$$\dots \longrightarrow (x_{i-1}^D, x_{i-1}) \longrightarrow (x_i^D, x_i) \longrightarrow (x_{i+1}^D, x_{i+1}) \longrightarrow \dots, \quad (5.27)$$

and if we convene to set $(\mathcal{T}_0, \mathcal{S}_0) = (x_i^D, x_i)$ for some i , then $(p, q) = (x_{i+1}^D, x_{i+1})$.

6 Self-dual states

We have seen that, as a consequence of $\mathcal{S}_0$ – $\mathcal{T}_0$ duality, there is a full correspondence among the u – and v –transformations. This duality indicates that any physical system with given $\mathcal{W}$ has two equivalent descriptions, corresponding to the $\mathcal{S}_0$ and $\mathcal{T}_0$ pictures. This suggests to investigate whether this dual structure by itself may select some distinguished states. We will see that this question will lead to the determination of states with rather peculiar properties. In particular, based on Legendre duality, we will see that the structure of time evolution will imply the appearance of imaginary numbers in considering the $p \longleftrightarrow q$ interchange for the highest symmetric states.

6.1 Where do the $\mathcal{S}_0$ – $\mathcal{T}_0$ pictures overlap?

Let us consider the naturally selected $\mathcal{W}$ states corresponding to the special case in which the $\mathcal{S}_0$ and $\mathcal{T}_0$ pictures overlap. On general grounds we should expect that the distinguished states, for which $\mathcal{S}_0$ and $\mathcal{T}_0$ have the same functional structure, have some peculiar properties. In order to find this common subspace we consider the interchange of the $\mathcal{S}_0$ and $\mathcal{T}_0$ pictures given by

$$q \longrightarrow \tilde{q} = \alpha p, \quad p \longrightarrow \tilde{p} = \beta q. \quad (6.1)$$

This implies that

$$\frac{\partial \tilde{\mathcal{T}}_0}{\partial \tilde{p}} = \alpha \frac{\partial \mathcal{S}_0}{\partial q}, \quad \frac{\partial \tilde{\mathcal{S}}_0}{\partial \tilde{q}} = \beta \frac{\partial \mathcal{T}_0}{\partial p}, \quad (6.2)$$

which is equivalent to

$$\frac{\partial \tilde{\mathcal{T}}_0}{\partial q} = \alpha \beta \frac{\partial \mathcal{S}_0}{\partial q}, \quad \frac{\partial \tilde{\mathcal{S}}_0}{\partial p} = \alpha \beta \frac{\partial \mathcal{T}_0}{\partial p}, \quad (6.3)$$

that is

$$\tilde{\mathcal{S}}_0(\tilde{q}) = \alpha \beta \mathcal{T}_0(p) + cnst, \quad \tilde{\mathcal{T}}_0(\tilde{p}) = \alpha \beta \mathcal{S}_0(q) + cnst, \quad (6.4)$$

so that $\tilde{\mathcal{S}}_0(\tilde{q})$ and $\tilde{\mathcal{T}}_0(\tilde{p})$ are essentially the Legendre transform of $\mathcal{S}_0(q)$ and $\mathcal{T}_0(p)$ respectively. Since interchanging the role of q and p twice has no effect on the functional dependence relating q and p , we have that up to an additive constant

$$\tilde{\tilde{\mathcal{S}}}_0 = \mathcal{S}_0, \quad \tilde{\tilde{\mathcal{T}}}_0 = \mathcal{T}_0, \quad (6.5)$$

so that

$$(\alpha\beta)^2 = 1. \quad (6.6)$$

The distinguished $\mathcal{W}$ states are precisely those states which are left invariant by (6.1) and (6.4). Hence, up to an additive constant

$$\tilde{\mathcal{S}}_0(\tilde{q}) = \mathcal{S}_0(q), \quad \tilde{\mathcal{T}}_0(\tilde{p}) = \mathcal{T}_0(p). \quad (6.7)$$

Since $\mathcal{S} = \mathcal{S}_0 - Et$ and $\mathcal{T} = \mathcal{T}_0 + Et$, it follows by (6.4)(6.6)(6.7) that these states correspond to

$$\mathcal{S} = \pm\mathcal{T} + \text{cnst}. \quad (6.8)$$

As (6.8) should be stable under time evolution, the relation (3.29) fixes the sign ambiguity and sets

$$\alpha\beta = -1. \quad (6.9)$$

Therefore, the distinguished $\mathcal{W}$ states correspond to

$$\mathcal{S} = -\mathcal{T} + \text{cnst}. \quad (6.10)$$

On the other hand, since $\mathcal{S} = pq - \mathcal{T}$, we have

$$pq = \gamma, \quad (6.11)$$

where γ is a constant. By (6.11) it follows that in the case of distinguished states the transformation (6.1) is equivalent to

$$q \longrightarrow \tilde{q} = \frac{\alpha\gamma}{q}, \quad p \longrightarrow \tilde{p} = \frac{\beta\gamma}{p}. \quad (6.12)$$

Let us consider the R -transformation (4.16)(4.17) in the case of the distinguished states (6.11)

$$R : q \longrightarrow q^v = \frac{\delta}{q}, \quad R : p \longrightarrow p_v = -\frac{\gamma^2}{\delta p}. \quad (6.13)$$

By (6.12) and (6.13) it follows that for the distinguished states the interchanging transformation (6.1) corresponds to an R -transformation with

$$\delta = \alpha\gamma = -\frac{\gamma}{\beta}. \quad (6.14)$$

Among the possible transformations (6.1), labeled by α and β , with $\alpha\beta = -1$, the one with the highest symmetric structure corresponds to the case in which (6.1) remains invariant under the interchange

$$p \longleftrightarrow q, \quad (6.15)$$

that is when

$$\alpha = \beta. \quad (6.16)$$

Observe that, without loss of generality, we are setting to 1 a dimensional constant one should consider in (6.16). It follows by (6.9) and (6.16) that the highest symmetric states are those with

$$\alpha = \beta = \pm i. \quad (6.17)$$

This shows a relation between stability of the Legendre transformation under time evolution, expressed by (3.29), which in turn implies the minus sign in (6.9), and the appearance of the imaginary factor in (6.17). The interesting outcome is that, based on Legendre duality, we have seen that the structure of time evolution implies the appearance of imaginary numbers for the highest symmetric states.

Let us now consider another way to select the distinguished states which will shed new light on their basic nature. First of all note that if the $\mathcal{S}_0$ and $\mathcal{T}_0$ pictures coincide, then, in particular, the v -action on $\mathcal{T}_0$ should be equivalent to the u -action on $\mathcal{S}_0$, that is

$$\delta_v \mathcal{T}_0 = \delta_u \mathcal{S}_0, \quad (6.18)$$

which is satisfied by (6.7). Then, to find the distinguished states we may first look for the Möbius transformations

$$q^v = \frac{A_v q + B_v}{C_v q + D_v}, \quad p^u = \frac{A_u p + B_u}{C_u p + D_u}, \quad (6.19)$$

satisfying (6.18), that by (4.5) and (5.12) is equivalent to

$$\rho^{-1}(A_v C_v q^2 + 2B_v C_v q + B_v D_v)p = \sigma^{-1}(A_u C_u p^2 + 2B_u C_u p + B_u D_u)q. \quad (6.20)$$

6.2 Dilatations

Eq.(6.20) shows that the answer to the above question has two solutions which do not fix any functional relation between p and q . One solution is given by $B_v = 0$, $C_v = 0$ and $B_u = 0$, $C_u = 0$. This corresponds to the dilatations

$$q^v = \frac{A_v}{D_v} q, \quad p^u = \frac{D_u}{A_u} p, \quad (6.21)$$

and

$$p^u = \frac{A_u}{D_u} p, \quad q_u = \frac{D_u}{A_u} q, \quad (6.22)$$

and

$$\delta_v \mathcal{T}_0 = 0 = \delta_u \mathcal{S}_0. \quad (6.23)$$

6.3 Changing sign: $pq \longrightarrow p^u q_u = -pq = p_v q^v$

The other solution of (6.20), which does not fix the functional dependence between p and q , is given by $A_v = 0$, $D_v = 0$ and $A_u = 0$, $D_u = 0$, corresponding to

$$q^v = \frac{B_v}{C_v} \frac{1}{q}, \quad p_v = -\frac{C_v}{B_v} q^2 p, \quad (6.24)$$

and

$$p^u = \frac{B_u}{C_u} \frac{1}{p}, \quad q_u = -\frac{C_u}{B_u} p^2 q. \quad (6.25)$$

Observe that the effect of this transformation is that of changing the sign of pq

$$pq \longrightarrow p^u q_u = -pq = p_v q^v. \quad (6.26)$$

We also have

$$\delta_v \mathcal{T}_0 = -2pq = \delta_u \mathcal{S}_0, \quad (6.27)$$

and as $\rho = -B_v C_v$ and $\sigma = -B_u C_u$, there are no constraints on the coefficients B_u, C_u, B_v, C_v .

6.4 Self-dual states: $q^v = q_u, p^u = p_v$

The solutions of (6.20) we found did not imply any constraint on the state. In other words, these solutions do not fix the dynamics. However, this investigation naturally leads to extend (6.18) by imposing the conditions with the highest symmetry, that is

$$q^v = q_u, \quad (6.28)$$

and

$$p^u = p_v. \quad (6.29)$$

In the case of dilatations, both (6.28) and (6.29) give

$$\frac{A_v}{D_v} = \frac{D_u}{A_u}. \quad (6.30)$$

In the case of Eqs.(6.24) and (6.25), both (6.28) and (6.29) yield

$$\frac{B_v B_u}{C_v C_u} = -p^2 q^2, \quad (6.31)$$

implying that

$$pq = \gamma, \quad (6.32)$$

with γ a constant. By (6.27) and (6.32) we have

$$\delta_v \mathcal{T}_0 = -2\gamma = \delta_u \mathcal{S}_0. \quad (6.33)$$

Therefore, the transformations (6.19) meeting the requirements (6.18)(6.28) and (6.29) should satisfy

$$\frac{B_v B_u}{C_v C_u} = -\gamma^2. \quad (6.34)$$

Observe that this transformation corresponds to interchanging p and q , namely

$$q \longrightarrow q^v = \frac{1}{\gamma} \frac{B_v}{C_v} p = -\gamma \frac{C_u}{B_u} p = q_u, \quad (6.35)$$

which is equivalent to

$$p \longrightarrow p^u = \frac{1}{\gamma} \frac{B_u}{C_u} q = -\gamma \frac{C_v}{B_v} q = p_v. \quad (6.36)$$

For these states the transformations (6.35) and (6.36) correspond to (6.1) with

$$\alpha = \frac{1}{\gamma} \frac{B_v}{C_v}, \quad \beta = -\gamma \frac{C_v}{B_v}, \quad (6.37)$$

and $\alpha\beta = -1$. It follows by (6.9) that these states correspond to the distinguished states. Observe that by (6.34) it follows that if $B_v B_u / C_v C_u > 0$, then γ is purely imaginary. Since

$$\delta_v(pq) = \delta_u(qp) = -pq = -\gamma, \quad (6.38)$$

it follows that if $B_v B_u / C_v C_u > 0$, then $\delta_v(pq) = \delta_u(qp)$ corresponds to the complex conjugation

$$\delta_v(\gamma) = \delta_u(\gamma) = \bar{\gamma}. \quad (6.39)$$

Based on purely symmetry arguments, we arrived to (6.17) in which the imaginary factor appears. This equation is essentially a consequence of (6.9) whose minus sign originates from (3.29). Hence, the origin of the imaginary factor in (6.17) traces back to (3.29) which relates the time evolution of the $\mathcal{S}$ and $\mathcal{T}$ pictures. As we will see, this relation between time evolution and imaginary number is at the heart of the imaginary factor appearing in the relation between $\mathcal{S}_0$ and the solutions of the SE. On the other hand, this factor is the signal that time evolution in QM is related to imaginary numbers. Hence, time evolution in QM is related to time evolution of the $\mathcal{S}$ and $\mathcal{T}$ dual pictures

Due to the highest symmetry (6.28)(6.29), we will call self-dual states the distinguished states. As for these states we have $p = \gamma/q$, the self-dual states correspond to

$$\mathcal{S}_0(q) = \gamma \ln \gamma_q q, \quad \mathcal{T}_0(p) = \gamma \ln \gamma_p p, \quad (6.40)$$

where, due to

$$\mathcal{S}_0 + \mathcal{T}_0 = pq = \gamma, \quad (6.41)$$

the dimensional constants γ_p , γ_q and γ satisfy

$$\gamma_p \gamma_q \gamma = e. \quad (6.42)$$

We also note that when $p = \gamma/q$, the solutions of (4.9) and (5.13) coincide and

$$\mathcal{U}(\mathbf{s}) = -\frac{1}{4\gamma^2} = \mathcal{V}(\mathbf{t}). \quad (6.43)$$

It follows that for the self-dual states the canonical equation

$$\left(\frac{\partial^2}{\partial \mathbf{s}^2} - \frac{1}{4\gamma^2} \right) q \sqrt{p} = 0 = \left(\frac{\partial^2}{\partial \mathbf{s}^2} - \frac{1}{4\gamma^2} \right) \sqrt{p}, \quad (6.44)$$

coincides with its dual version

$$\left(\frac{\partial^2}{\partial \mathbf{t}^2} - \frac{1}{4\gamma^2}\right) p\sqrt{q} = 0 = \left(\frac{\partial^2}{\partial \mathbf{t}^2} - \frac{1}{4\gamma^2}\right) \sqrt{q}. \quad (6.45)$$

Considering the inverse of $\mathcal{S}_0$ and $\mathcal{T}_0$ in (6.40), we have $q = \gamma_q^{-1} \exp(\mathbf{s}/\gamma)$, $p = \gamma_p^{-1} \exp(\mathbf{t}/\gamma)$, so that the solutions of Eqs.(4.9) and (5.13), with $\mathcal{U}(\mathbf{s}) = -1/4\gamma^2 = \mathcal{V}(\mathbf{t})$, are

$$q\sqrt{p} = \sqrt{\gamma\gamma_q^{-1}} e^{\frac{1}{2\gamma}\mathbf{s}} = \sqrt{\gamma}\sqrt{q} = \gamma\sqrt{\gamma_p} e^{-\frac{1}{2\gamma}\mathbf{t}}, \quad (6.46)$$

and

$$\sqrt{p} = \sqrt{\gamma\gamma_q} e^{-\frac{1}{2\gamma}\mathbf{s}} = \sqrt{\gamma^{-1}} p\sqrt{q} = \sqrt{\gamma_p^{-1}} e^{\frac{1}{2\gamma}\mathbf{t}}. \quad (6.47)$$

7 The Equivalence Principle

In sect.2 we proposed that as in the search for canonical transformations leading to a system with vanishing Hamiltonian one obtains the HJ equation, we may similarly look for transformations q and p , seen as dependent variables, reducing to the free system with vanishing energy. This is at the heart of the EP we will formulate in this section.

7.1 Equivalence Principle and v -transformations

The discussion concerning the way of inducing the transformations $\mathcal{S}_0 \longrightarrow \mathcal{S}_0^v$ made in sect.2, allows us to formulate the above question more precisely

Given an arbitrary system with reduced action $\mathcal{S}_0(q)$, find the coordinate transformation $q \longrightarrow q^{v_0} = v_0(q)$, such that the new reduced action $\mathcal{S}_0^{v_0}$, defined by

$$\mathcal{S}_0^{v_0}(q^{v_0}) = \mathcal{S}_0(q), \quad (7.1)$$

corresponds to the system with $V - E = 0$.

In the following we will use the notation $q^0 \equiv q^{v_0}$, $\mathcal{S}_0^0 \equiv \mathcal{S}_0^{v_0}$, and denote by $\mathcal{W}^0(q^0)$ the state corresponding to $\mathcal{W} = 0$. Observe that the structure of the states described by $\mathcal{S}_0^0$ and $\mathcal{S}_0$ determines the “trivializing coordinate” q^0 to be

$$q \longrightarrow q^0 = \mathcal{S}_0^{0^{-1}} \circ \mathcal{S}_0(q), \quad (7.2)$$

The problem (7.1) suggested the following Equivalence Principle (EP) formulated in [1]

For each pair $\mathcal{W}^a, \mathcal{W}^b$, there is a v -transformation $q^a \longrightarrow q^b = v(q^a)$ such that

$$\mathcal{W}^a(q^a) \longrightarrow \mathcal{W}^b(q^b). \quad (7.3)$$

7.2 Classical Mechanics and the Equivalence Principle

We now show the basic fact that the EP cannot be consistently implemented in CM. In order to implement the EP in CM we should first understand the nature of the transformation

$$\mathcal{W}(q) \longrightarrow \mathcal{W}^v(q^v), \quad (7.4)$$

induced by

$$\mathcal{S}_0^{cl}(q) \longrightarrow \mathcal{S}_0^{cl v}(q^v) = \mathcal{S}_0^{cl}(q(q^v)). \quad (7.5)$$

To answer this question we use the fact that the CSHJE (2.15) provides a correspondence between $\mathcal{W}$ and $\mathcal{S}_0^{cl}$. In particular, $\mathcal{S}_0^{cl v}(q^v)$ should satisfy the CSHJE

$$\frac{1}{2m} \left(\frac{\partial \mathcal{S}_0^{cl v}(q^v)}{\partial q^v} \right)^2 + \mathcal{W}^v(q^v) = 0. \quad (7.6)$$

Using $\mathcal{S}_0^{cl v}(q^v) = \mathcal{S}_0^{cl}(q)$, which defines the classical v -transformations $q \longrightarrow q^v = v(q)$, and comparing (2.15) with (7.6) we obtain

$$\mathcal{W}(q) \longrightarrow \mathcal{W}^v(q^v) = (\partial_{q^v} q)^2 \mathcal{W}(q), \quad (7.7)$$

that is

$$\mathcal{W}^v(q^v)(dq^v)^2 = \mathcal{W}(q)(dq)^2. \quad (7.8)$$

Therefore, in CM consistency requires that $\mathcal{W}(q)$ belongs to $\mathcal{Q}^{cl}$, the space of functions transforming as quadratic differentials under the classical v -transformations, that is

$$\mathcal{W} \in \mathcal{Q}^{cl} = \left\{ g \mid g^v(q^v) = (\partial_{q^v} q)^2 g(q) \right\}. \quad (7.9)$$

Let us now consider the case of the state $\mathcal{W}^0$. By (7.7) it follows that

$$\mathcal{W}^0(q^0) \longrightarrow \mathcal{W}^v(q^v) = (\partial_{q^v} q^0)^2 \mathcal{W}^0(q^0) = 0. \quad (7.10)$$

Then we have [1]

$\mathcal{W}$ states transform as quadratic differentials under classical v -maps. It follows that $\mathcal{W}^0$ is a fixed point in $\mathcal{H}$. Equivalently, in CM the space $\mathcal{H}$ cannot be reduced to a point upon factorization by the classical v -transformations. Hence, the EP (7.3) cannot be consistently implemented in CM.

7.3 Modifying the classical HJ equation

It is clear that in order to implement the EP we have to modify the CSHJE. Let us consider the properties that the equation satisfied by $\mathcal{S}_0$ should have. First of all, observe that by Eqs.(3.19)–(3.20) it follows that adding a constant to either $\mathcal{S}_0$ or $\mathcal{T}_0$ does not change the dynamics. Therefore, the most general equation that $\mathcal{S}_0$ should satisfy has the form

$$F(\mathcal{S}'_0, \mathcal{S}''_0, \dots) = 0, \quad (7.11)$$

where $' \equiv \partial_q$. Since in the classical limit we have

$$F(\mathcal{S}'_0, \mathcal{S}''_0, \dots) = 0 \longrightarrow \frac{1}{2m} \left(\frac{\partial \mathcal{S}_0^{cl}}{\partial q} \right)^2 + \mathcal{W}(q) = 0, \quad (7.12)$$

it is useful to write Eq.(7.11) in the form

$$\frac{1}{2m} \left(\frac{\partial \mathcal{S}_0(q)}{\partial q} \right)^2 + \mathcal{W}(q) + Q(q) = 0, \quad (7.13)$$

so that the classical limit corresponds to

$$Q \longrightarrow 0. \quad (7.14)$$

7.4 Covariance as consistency condition

Let us derive the transformation properties of $\mathcal{W} + Q$ by consistency. Namely, we first consider the transformed version of Eq.(7.13)

$$\frac{1}{2m} \left(\frac{\partial \mathcal{S}_0^v(q^v)}{\partial q^v} \right)^2 + \mathcal{W}^v(q^v) + Q^v(q^v) = 0, \quad (7.15)$$

then note that by (2.37) $(\partial_{q^v} \mathcal{S}_0^v)^2 = (\partial_{q^v} q)^2 (\partial_q \mathcal{S}_0)^2$, so that (7.13) and (7.15) yield

$$\mathcal{W}^v(q^v) + Q^v(q^v) = (\partial_{q^v} q)^2 (\mathcal{W}(q) + Q(q)). \quad (7.16)$$

Hence $\mathcal{W} + Q$ belongs to the space $\mathcal{Q}$ of functions transforming as quadratic differentials under VTs

$$(\mathcal{W} + Q) \in \mathcal{Q}. \quad (7.17)$$

Therefore, choosing (2.37) to represent the functional change $\mathcal{S}_0 \longrightarrow \mathcal{S}_0^v$ provides simple transformation properties for $\mathcal{W} + Q$. In particular, Eq.(7.13) transforms in a manifestly covariant way with the choice (2.37). Had we used another representation, *e.g.* (2.38), then the resulting transformation rule for $\mathcal{W} + Q$ would involve also f and $\mathcal{S}_0$. We stress that there is nothing wrong in considering this alternative representation, nevertheless it is clear that the formalism would be far more cumbersome.

8 Implementing the Equivalence Principle

The relevance of the EP is manifest already in considering the transformation properties of $\mathcal{W}$ and Q . Actually, according to the EP, all the $\mathcal{W}$'s are connected with each other by a coordinate transformation. On the other hand, we have seen that if $\mathcal{W}$ transforms as a quadratic differential, then the state $\mathcal{W}^0$ would be a fixed point in the space $\mathcal{H}$. It follows that

$$\mathcal{W} \notin \mathcal{Q}. \quad (8.1)$$

On the other hand, we have seen that by consistency $(\mathcal{W} + Q) \in \mathcal{Q}$, so that by (8.1)

$$Q \notin \mathcal{Q}. \quad (8.2)$$

In this section, following [1], we will see that the transformation properties of $\mathcal{W}$ and Q lead to a cocycle condition which will be crucial to fix Q . In particular, we will first find the transformation properties of an inhomogeneous term one is forced to introduce in the transformation properties of $\mathcal{W}$. In the next section we will see that these properties fix this term and therefore Q . In the present section we will also consider the problem of selecting a self-dual state as “reference state”.

8.1 $\mathcal{W}$ states as inhomogeneous terms

Note that the only possible way to reach an arbitrary state $\mathcal{W}^v \neq 0$ starting from the state $\mathcal{W}^0$, is that it transforms with an inhomogeneous term. This is the physical content of Eq.(8.1). This implies that under a VT, in general we have

$$\mathcal{W}^0 \longrightarrow \mathcal{W}^v \neq 0. \quad (8.3)$$

In particular, any two arbitrary states $\mathcal{W}$ and $\mathcal{W}^v$ are related by

$$\mathcal{W}^v(q^v) = (\partial_{q^v} q)^2 \mathcal{W}(q) + (q; q^v), \quad (8.4)$$

where $(q; q^v)$ is the inhomogeneous term to be determined. By (7.16) we have

$$\mathcal{W}^v(q^v) + Q^v(q^v) = (\partial_{q^v} q)^2 (\mathcal{W}(q) + Q(q)), \quad (8.5)$$

so that, replacing $\mathcal{W}^v(q^v)$ in (8.5) with the right hand side of (8.4), we obtain

$$(\partial_{q^v} q)^2 \mathcal{W}(q) + (q; q^v) + Q^v(q^v) = (\partial_{q^v} q)^2 (\mathcal{W}(q) + Q(q)), \quad (8.6)$$

that is

$$Q^v(q^v) = (\partial_{q^v} q)^2 Q(q) - (q; q^v). \quad (8.7)$$

Replacing $\mathcal{W}(q)$ in (8.4) with $\mathcal{W}^0(q^0)$, we have $\mathcal{W}^v(q^v) = (q^0; q^v)$, so that

$$\mathcal{W}(q) = (q^0; q). \quad (8.8)$$

Therefore, all the states correspond to the inhomogeneous, and unique, part in the transformation of $\mathcal{W}^0$ induced by v -maps. In other words, all the states originate from the map $q^0 \longrightarrow q = v_0^{-1}(q^0)$. We also note that, in view of (8.8), $\mathcal{W}^0$ plays a distinguished role, as any state can be written in the universal form (8.8) in which the distinguished coordinate q^0 , representing the state $\mathcal{W}^0$, appears.

Observe that by construction, Eq(7.13) has the simple transformation property

$$\frac{1}{2m} \left(\frac{\partial \mathcal{S}_0^v(q^v)}{\partial q^v} \right)^2 + \mathcal{W}^v(q^v) + Q^v(q^v) = \left(\frac{\partial q}{\partial q^v} \right)^2 \left[\frac{1}{2m} \left(\frac{\partial \mathcal{S}_0(q)}{\partial q} \right)^2 + \mathcal{W}(q) + Q(q) \right] = 0. \quad (8.9)$$

We saw in sect.6 that there is a set of distinguished, or self-dual, states which are naturally selected by purely symmetry arguments. In doing this we found an explicit expression for $\mathcal{S}_0$ and $\mathcal{T}_0$ corresponding to the self-dual states. Nevertheless, we did not consider the problem of finding the corresponding expression of $\mathcal{W}$. However, even if the form of $\mathcal{W}$ has not been derived,⁶ one expects that its structure should reflect the highest symmetry we used to fix the explicit form (6.40) of $\mathcal{S}_0$ and $\mathcal{T}_0$. We will see that the state $\mathcal{W}^0$ precisely corresponds to a self-dual state. As such the state $\mathcal{W}^0$ can be seen as the state in which the $\mathcal{S}_0$ and $\mathcal{T}_0$ pictures overlap.⁷

8.2 The cocycle condition

We now consider the transformation properties of $(q^a; q^b)$. Comparing

$$\mathcal{W}^b(q^b) = \left(\partial_{q^b} q^a\right)^2 \mathcal{W}^a(q^a) + (q^a; q^b) = (q^0; q^b), \quad (8.10)$$

with the same formula with q^a and q^b interchanged we have

$$(q^b; q^a) = -(\partial_{q^a} q^b)^2 (q^a; q^b), \quad (8.11)$$

in particular

$$(q; q) = 0. \quad (8.12)$$

More generally, comparing

$$\begin{aligned} \mathcal{W}^b(q^b) &= \left(\partial_{q^b} q^c\right)^2 \mathcal{W}^c(q^c) + (q^c; q^b) = \left(\partial_{q^b} q^c\right)^2 \left[(\partial_{q^c} q^a)^2 \mathcal{W}^a(q^a) + (q^a; q^c) \right] + (q^c; q^b) = \\ &= \left(\partial_{q^b} q^a\right)^2 \mathcal{W}^a(q^a) + \left(\partial_{q^b} q^c\right)^2 (q^a; q^c) + (q^c; q^b), \end{aligned} \quad (8.13)$$

with Eq.(8.10), we obtain the basic cocycle condition

$$(q^a; q^c) = \left(\partial_{q^c} q^b\right)^2 \left[(q^a; q^b) - (q^c; q^b) \right], \quad (8.14)$$

which can be seen as the essence of the EP.

Before going further some remarks are in order. We started by noticing that since the coordinate of a physical system can be seen as an independent variable, its choice essentially reduces to a notational one. This suggested considering the transformation among different physical systems to be performed through VTs which are equivalent to consider $\mathcal{S}_0$ as a scalar function. The power of this approach is evident in implementing the EP. In particular, we obtained a manifestly covariant description where

⁶Note that finding the explicit form of $\mathcal{W}$ associated to the self-dual states, would essentially be equivalent to finding the dynamical equation for $\mathcal{S}_0$.

⁷We stress that in deriving the quantum version of the HJ equation, we will only use the EP and the existence of CM. In doing this we frequently discuss the role of the self-dual states. One aim of these discussions is to show that the quantum HJ equation fully reflects the natural dual structure which emerges in considering the Legendre pair $\mathcal{S}_0$ - $\mathcal{T}_0$. Therefore, in deriving the quantum HJ equation, we do not assume any kind of duality to be satisfied, rather we will see that the solution following from the EP satisfies $\mathcal{S}_0$ - $\mathcal{T}_0$ duality.

the scalar nature of $\mathcal{S}_0$ implied that $(\mathcal{W} + Q) \in \mathcal{Q}$. The whole investigation culminated in the basic equation (8.14), which makes clear the basic nature of the EP. As already observed in sect.2, even if considering another kind of representation for the transformation of $\mathcal{S}_0$ would yield the same results, these would not be manifestly covariant and the elegance of the construction would be lost. In this sense representing the state transformation by the pullback of $\mathcal{S}_0$ by v^{-1*} is the natural choice.

By eqs.(8.4)(8.7) and (8.14) we have

$$\mathcal{W} \in \mathcal{P}, \quad Q \in \mathcal{P}, \quad (\cdot; q) \in \mathcal{P}, \quad (8.15)$$

where $\mathcal{P}$ is the space of “inhomogeneous” quadratic differentials

$$\mathcal{P} = \left\{ f \mid f^v(q^v) = (\partial_{q^v} q)^2 f(q) + \alpha_f(q; q^v) \right\}, \quad (8.16)$$

with α_f a constant. Note that the difference between any two elements f_i and f_j in $\mathcal{P}$ with $\alpha_{f_i} = \alpha_{f_j}$, belongs to the space $\mathcal{Q}$, that is $(f_i - f_j) \in \mathcal{Q}$, $f_i, f_j \in \mathcal{P}$, $\alpha_{f_i} = \alpha_{f_j}$.

As we will see, Eq.(8.14), which follows as a consistency condition from the EP, univocally fixes the differential equation for $\mathcal{S}_0$. In particular, in the following we will first show that Eq.(8.14), and (7.11) implies $(q; \gamma(q)) = 0 = (\gamma(q); q)$, with γ a Möbius transformation. In the next section, we will prove that $(q^a; q^b)$ is proportional to the Schwarzian derivative $\{q^a, q^b\}$. It follows that $\mathcal{P}$ is the space of functions transforming as projective connections under v -maps. The fact that $(q^a; q^b) \propto \{q^a, q^b\}$, is related to the so-called pseudogroup property [34] associated to (8.14). This is the basic defining property of complex analytic structures as it implies that the composition of two complex analytic local homeomorphisms is again a local homeomorphism. It turns out that any element g in the family of one-dimensional complex analytic local homeomorphisms, solutions of a system of differential equations involving only the first and higher derivatives, and satisfying the pseudogroup property, satisfies either $g''(x)/g'(x) = 0$ or $\{g(x), x\} = 0$ [34][35][36]. It is worth noticing that this issue is related to the cohomology of Lie algebras [37][38].

8.3 Cocycle condition and Möbius symmetry

Let us start observing that if the cocycle condition (8.14) is satisfied by $(f(q); q)$, then this is still satisfied by adding a coboundary term

$$(f(q); q) \longrightarrow (f(q); q) + (\partial_q f)^2 G(f(q)) - G(q). \quad (8.17)$$

Since $(Aq; q)$ evaluated at $q = 0$ is independent of A , we have

$$0 = (q; q) = (q; q)|_{q=0} = (Aq; q)|_{q=0}. \quad (8.18)$$

Therefore, if both $(f(q); q)$ and (8.17) satisfy (8.14), then $G(0) = 0$, which is the unique condition that G should satisfy. We now use (7.11) to fix the ambiguity (8.17). First of all observe that by (8.8) the differential equation we are looking for is

$$(q^0; q) = \mathcal{W}(q). \quad (8.19)$$

Then, recalling that $q^0 = \mathcal{S}_0^{-1} \circ \mathcal{S}_0(q)$ (see Eq.(7.2)), we have that a necessary condition to satisfy (7.11) is that $(q^0; q)$ depends only on the first and higher derivatives of q^0 . This in turn implies that for any constant B we have $(q^a + B; q^b) = (q^a; q^b)$, that together with (8.11) gives

$$(q^a + B; q^b) = (q^a; q^b) = (q^a; q^b + B). \quad (8.20)$$

Let A be a non-vanishing constant and set $h(A, q) = (Aq; q)$. By (8.20) we have $h(A, q + B) = h(A, q)$, that is $h(A, q)$ is independent of q . On the other hand by (8.18) we have $h(A, 0) = 0$. This fact and (8.11), imply

$$(Aq; q) = 0 = (q; Aq). \quad (8.21)$$

Eq.(8.14) implies $(q^a; Aq^b) = A^{-2}((q^a; q^b) - (Aq^b; q^b))$, so that by (8.21)

$$(q^a; Aq^b) = A^{-2}(q^a; q^b). \quad (8.22)$$

By (8.11) and (8.22) we have $(Aq^a; q^b) = -A^{-2}(\partial_{q^b} q^a)^2(q^b; Aq^a) = -(\partial_{q^b} q^a)^2(q^b; q^a) = (q^a; q^b)$, that is

$$(Aq^a; q^b) = (q^a; q^b). \quad (8.23)$$

Setting $f(q) = q^{-2}(q; q^{-1})$ and noticing that by (8.11) and (8.23) $f(Aq) = -f(q^{-1})$, we obtain

$$(q; q^{-1}) = 0 = (q^{-1}; q). \quad (8.24)$$

Furthermore, since by (8.14) and (8.24) one has $(q^a; q^{b^{-1}}) = q^{b^4}(q^a; q^b)$, it follows that

$$(q^{a^{-1}}; q^b) = -(\partial_{q^b} q^{a^{-1}})^2(q^b; q^{a^{-1}}) = -(\partial_{q^b} q^a)^2(q^b; q^a) = (q^a; q^b), \quad (8.25)$$

so that

$$(q^{a^{-1}}; q^b) = (q^a; q^b) = q^{b^{-4}}(q^a; q^{b^{-1}}). \quad (8.26)$$

Since translations, dilatations and inversion are the generators of the Möbius group, it follows by (8.20)(8.22)(8.23) and (8.26) that

$$(\gamma(q^a); q^b) = (q^a; q^b), \quad (8.27)$$

and

$$(q^a; \gamma(q^b)) = (\partial_{q^b} \gamma(q^b))^{-2}(q^a; q^b), \quad (8.28)$$

where

$$\gamma(q) = \frac{Aq + B}{Cq + D}, \quad (8.29)$$

with $\begin{pmatrix} A & B \\ C & D \end{pmatrix} \in GL(2, \mathbb{C})$. In particular

$$(\gamma(q); q) = 0 = (q; \gamma(q)). \quad (8.30)$$

For sake of completeness we recall that the infinitesimal versions of translations, dilatations and special conformal transformations of q are $q \longrightarrow q - \epsilon$, $q \longrightarrow q - \epsilon q$ and $q \longrightarrow q - \epsilon q^2$ respectively. Recall that a special conformal transformation has the form $q \longrightarrow q/(1 - Bq)$. In the case of infinitesimal Möbius transformations of $q + \epsilon f(q)$, we have

$$q + \epsilon f(q) \longrightarrow q + \epsilon f(q) - \epsilon(q + \epsilon f(q))^n, \quad n = 0, 1, 2, \quad (8.31)$$

so that the first-order contribution to the variation of $q + \epsilon f(q)$ coincides with that of q .

8.4 Selecting a self-dual state

Let us now go back to the question posed at the beginning of sect.7, Eq.(7.1), and to the EP. We saw that, if $\mathcal{S}_0 = \text{const}$, then the EP cannot be consistently implemented. Similarly, the Legendre transformation cannot be defined for $\mathcal{S}_0 = Aq + B$ and $\mathcal{T}_0 = Ap + B$. In CM, the state with $\mathcal{S}_0^{cl} = \text{const}$ corresponds to the free particle with vanishing energy, whereas in the case in which $\mathcal{S}_0^{cl} \propto q$ one has the free particle with $E \neq 0$. Therefore, problems arise just for the free particle. The $\mathcal{S}_0$ – $\mathcal{T}_0$ duality investigated above, and the fact that the problems which arise in applying the EP reflect in the impossibility of defining the Legendre transformation, indicate that the solution to this question is related to the observed dualities. Let us recall that the self-dual (or distinguished) states

$$\mathcal{S}_0(q) = \gamma \ln \gamma_q q, \quad \mathcal{T}_0(p) = \gamma \ln \gamma_p p, \quad (8.32)$$

parameterized by γ , are those with the highest symmetric properties as they are characterized by

$$\delta_v \mathcal{T}_0 = \delta_u \mathcal{S}_0, \quad q^v = q_u, \quad p^u = p_v. \quad (8.33)$$

We have seen that the states (8.32) play a special role. On the other hand, the fact that all the states satisfy the relation $\mathcal{S}_0^v(q^v) = \mathcal{S}_0(q)$, suggests selecting a particular self-dual state (8.32) as reference state. This means choosing a given value γ_{sd} for the constant γ in (8.32). We denote by q^{sd} and p^{sd} the corresponding coordinate and the momentum respectively. Therefore, we have the reference state

$$\mathcal{S}_0^{sd}(q^{sd}) = \gamma_{sd} \ln \gamma_q q^{sd}, \quad \mathcal{T}_0(p^{sd}) = \gamma_{sd} \ln \gamma_p p^{sd}. \quad (8.34)$$

We will denote by $\mathcal{W}^{sd}$ the distinguished self-dual state $\mathcal{W}$ corresponding to $\mathcal{S}_0^{sd}(q^{sd})$. Therefore, all possible states have a reduced action of the form

$$\mathcal{S}_0(q) = \mathcal{S}_0^{sd}(q^{sd}) = \gamma_{sd} \ln \gamma_q q^{sd}(q), \quad (8.35)$$

which implies that the coordinate transformation reducing an arbitrary system with reduced action $\mathcal{S}_0(q)$ to the distinguished self-dual state $\mathcal{S}_0^{sd}(q^{sd})$ is

$$q \longrightarrow q^{sd} = v_{sd}(q) = \gamma_q^{-1} e^{\frac{1}{\gamma_{sd}} \mathcal{S}_0(q)}. \quad (8.36)$$

Note that, according to the EP, the transformation should be locally invertible. Therefore, there is also the map transforming $\mathcal{S}_0^{sd}$ into $\mathcal{S}_0$, that is $q^{sd} \longrightarrow q = v_{sd}^{-1}(q^{sd})$.

9 The quantum stationary HJ equation

We can now determine the explicit expression of Q in Eq.(7.13). We first consider some facts about the Schwarzian derivative. Next, we will prove the basic identification $(q^a; q^b) = -\beta^2 \{q^a, q^b\}/4m$. This result will lead to the explicit determination of Q . After proving that Q is univocally fixed, we will show in the next section that the modified stationary HJ equation implies the SE.

9.1 Complex entries and real values of the Schwarzian derivative

Let us recall a few facts about the Schwarzian derivative. Let us first consider $\{e^h, x\}$. Since by (4.12) $\{y, x\} = y'^2(\{y, y\} - \{x, y\}) = -y'^2\{x, y\}$, we have

$$\{e^h, x\} = -\left(\frac{\partial e^h}{\partial x}\right)^2 \{x, e^h\}. \quad (9.1)$$

On the other hand, (4.12) implies

$$\{x, e^h\} = e^{-2h}\{x, h\} - \{e^h, h\} = e^{-2h}\{x, h\} + \frac{1}{2}, \quad (9.2)$$

and by (9.1)

$$\{e^h, x\} = \{h, x\} - \frac{1}{2}\left(\frac{\partial h}{\partial x}\right)^2. \quad (9.3)$$

A property of $\{h, x\}$ is that it may take real values even if h and x take complex values. The basic reason is already evident from the invariance under the complex Möbius transformations $\{\gamma(h), x\} = \{h, x\}$. Another example is provided by $\{e^{ih}, x\} = \{h, x\} + (\partial_x h)^2/2$. This implies that if h and/or x take real or purely imaginary values, then $\{e^{ih}, x\} \in \mathbb{R}$. Similarly, if α is independent of x , we have

$$\{e^{i\alpha h}, x\} = \{h, x\} + \frac{\alpha^2}{2}\left(\frac{\partial h}{\partial x}\right)^2. \quad (9.4)$$

Observe that if h is independent of α , then

$$(\partial_x h)^2 = \alpha^{-1} \partial_\alpha \{e^{i\alpha h}, x\}. \quad (9.5)$$

The fact that $\{e^{i\alpha h}, x\}$ splits into two parts is a key point in our construction. This is a consequence of the following property of the Schwarzian derivative

$$\{h^\alpha, x\} = \{h, x\} + \frac{1}{2}(1 - \alpha^2)\left(\frac{h'}{h}\right)^2. \quad (9.6)$$

The identity (9.4) implies that $(\partial_q \mathcal{S}_0)^2$, which transforms as a quadratic differential under v -maps, can be expressed as the difference of two Schwarzian derivatives

$$\left(\frac{\partial \mathcal{S}_0}{\partial q}\right)^2 = \frac{\beta^2}{2} \left(\{e^{\frac{2i}{\beta} \mathcal{S}_0}, q\} - \{\mathcal{S}_0, q\} \right). \quad (9.7)$$

Observe that this basic identity forces us to introduce the dimensional constant β . We also note the appearance of the imaginary factor.

$$\mathbf{9.2} \quad (q^a; q^b) = -\frac{\beta^2}{4m} \{q^a, q^b\}$$

We now start showing that $(q^a; q^b) \propto \{q^a, q^b\}$. Since we are interested to what corresponds $(\cdot; q)$, we can consider, without loss of generality, the case in which q^a differs infinitesimally from $q \equiv q^b$, that is $q^a = q + \epsilon f(q)$. Since $(q^a; q^b)$ depends only on the first and higher derivatives of q^a , we have

$$(q + \epsilon f(q); q) = c_1 \epsilon f^{(k)}(q) + \mathcal{O}(\epsilon^2), \quad (9.8)$$

where $f^{(k)} \equiv \partial_q^k f$. We will see that

$$c_1 \neq 0. \quad (9.9)$$

Let us first fix the value of k in (9.8). To this end we note that

$$(Aq + \epsilon A f(q); Aq) = (q + \epsilon f(q); Aq) = A^{-2}(q + \epsilon f(q); q), \quad (9.10)$$

on the other hand, setting $F(Aq) = A f(q)$, by Eq.(9.8) we have

$$(Aq + \epsilon A f(q); Aq) = (Aq + \epsilon F(Aq); Aq) = c_1 \epsilon \partial_{Aq}^k F(Aq) = A^{1-k} c_1 \epsilon f^{(k)}(q), \quad (9.11)$$

so that $k = 3$, that is

$$(q + \epsilon f(q); q) = c_1 \epsilon f^{(3)}(q) + \mathcal{O}(\epsilon^2). \quad (9.12)$$

The above scaling property generalizes to higher-order contributions in ϵ . In particular, the contribution at order ϵ^n to $(Aq + \epsilon A f(q); Aq)$ is given by a sum of terms of the form

$$c_{i_1 \dots i_n} \partial_{Aq}^{i_1} \epsilon F(Aq) \dots \partial_{Aq}^{i_n} \epsilon F(Aq) = c_{i_1 \dots i_n} \epsilon^n A^{n - \sum i_k} f^{(i_1)}(q) \dots f^{(i_n)}(q), \quad (9.13)$$

and by (9.10)

$$\sum_{k=1}^n i_k = n + 2. \quad (9.14)$$

On the other hand, since $(q^a; q^b)$ depends only on the first and higher derivatives of q^a , we have

$$i_k \geq 1, \quad k \in [1, n]. \quad (9.15)$$

Eqs.(9.14)(9.15) imply that either

$$i_k = 3, \quad i_j = 1, \quad j \in [1, n], \quad j \neq k, \quad (9.16)$$

or

$$i_k = i_j = 2, \quad i_l = 1, \quad l \in [1, n], \quad l \neq k, l \neq j, \quad (9.17)$$

so that

$$(q + \epsilon f(q); q) = \sum_{n=1}^{\infty} \epsilon^n \left(c_n f^{(3)} f^{(1)^{n-1}} + d_n f^{(2)^2} f^{(1)^{n-2}} \right), \quad d_1 = 0. \quad (9.18)$$

We now show that either $c_1 \neq 0$ or $(q + \epsilon f(q); q) = 0$. Let us consider the transformations

$$q^b = v^{ba}(q^a), \quad q^c = v^{cb}(q^b) = v^{cb} \circ v^{ba}(q^a), \quad q^c = v^{ca}(q^a). \quad (9.19)$$

Note that $v^{ab} = v^{ba^{-1}}$, and

$$v^{ca} = v^{cb} \circ v^{ba}. \quad (9.20)$$

We can write these transformations in the form

$$q^b = q^a + \epsilon^{ba}(q^a), \quad q^c = q^b + \epsilon^{cb}(q^b) = q^b + \epsilon^{cb}(q^a + \epsilon^{ba}(q^a)), \quad q^c = q^a + \epsilon^{ca}(q^a). \quad (9.21)$$

Since $q^b = q^a - \epsilon^{ab}(q^b)$, we have $q^b = q^a - \epsilon^{ab}(q^a + \epsilon^{ba}(q^a))$ that compared with $q^b = q^a + \epsilon^{ba}(q^a)$ yields

$$\epsilon^{ba} + \epsilon^{ab} \circ (\mathbf{1} + \epsilon^{ba}) = 0, \quad (9.22)$$

where $\mathbf{1}$ denotes the identity map. More generally, Eq.(9.21) gives

$$\epsilon^{ca}(q^a) = \epsilon^{cb}(q^b) + \epsilon^{ba}(q^a) = \epsilon^{cb}(q^b) - \epsilon^{ab}(q^b), \quad (9.23)$$

so that we obtain (9.20) with $v^{yx} = \mathbf{1} + \epsilon^{yx}$

$$\epsilon^{ca} = \epsilon^{cb} \circ (\mathbf{1} + \epsilon^{ba}) + \epsilon^{ba} = (\mathbf{1} + \epsilon^{cb}) \circ (\mathbf{1} + \epsilon^{ba}) - \mathbf{1}. \quad (9.24)$$

Let us consider the case in which $\epsilon^{yx}(q^x) = \epsilon f_{yx}(q^x)$, with ϵ infinitesimal. At first-order in ϵ Eq.(9.24) reads

$$\epsilon^{ca} = \epsilon^{cb} + \epsilon^{ba}, \quad (9.25)$$

in particular, $\epsilon^{ab} = -\epsilon^{ba}$. Since $(q^a; q^b) = c_1 \epsilon^{ab''' } (q^b) + \mathcal{O}^{ab}(\epsilon^2)$, where $'$ denotes the derivative with respect to the argument, we can use the cocycle condition (8.14) to get

$$c_1 \epsilon^{ac''' } (q^c) + \mathcal{O}^{ac}(\epsilon^2) = (1 + \epsilon^{bc'}(q^c))^2 \left(c_1 \epsilon^{ab''' } (q^b) + \mathcal{O}^{ab}(\epsilon^2) - c_1 \epsilon^{cb''' } (q^b) - \mathcal{O}^{cb}(\epsilon^2) \right), \quad (9.26)$$

that at first-order in ϵ corresponds to (9.25). We see that $c_1 \neq 0$. For, if $c_1 = 0$, then by (9.26), at second-order in ϵ one would have

$$\mathcal{O}^{ac}(\epsilon^2) = \mathcal{O}^{ab}(\epsilon^2) - \mathcal{O}^{cb}(\epsilon^2), \quad (9.27)$$

which contradicts (9.25). In fact, by (9.18) we have

$$\mathcal{O}^{ab}(\epsilon^2) = c_2 \epsilon^{ab''' } (q^b) \epsilon^{ab'}(q^b) + d_2 \epsilon^{ab''^2}(q^b) + \mathcal{O}^{ab}(\epsilon^3), \quad (9.28)$$

that together with (9.27) provides a relation which cannot be consistent with $\epsilon^{ac}(q^c) = \epsilon^{ab}(q^b) - \epsilon^{cb}(q^b)$. A possibility is that $(q^a; q^b) = 0$. However, this is ruled out by the EP, so that $c_1 \neq 0$. Higher-order terms due to a non-vanishing c_1 are obtained by using $q^c = q^b + \epsilon^{cb}(q^b)$, $\epsilon^{ac}(q^c) = \epsilon^{ab}(q^b) - \epsilon^{cb}(q^b)$ and $\epsilon^{bc}(q^c) = -\epsilon^{cb}(q^b)$ in $c_1 \partial_{q^c}^3 \epsilon^{ac}(q^c)$ and in $c_1 \left(2 \partial_{q^c} \epsilon^{bc}(q^c) + \partial_{q^c} \epsilon^{bc}(q^c)^2 \right) \partial_{q^b}^3 \left(\epsilon^{ab}(q^b) - \epsilon^{cb}(q^b) \right)$. Note that one can also consider the case in which both the first- and second-order contributions to $(q^a; q^b)$ are vanishing. However, this possibility is ruled out by a similar analysis. In general, one has that if the first non-vanishing contribution to $(q^a; q^b)$ is of order ϵ^n , $n \geq 2$, then, unless $(q^a; q^b) = 0$, the cocycle condition (8.14) cannot be consistent with linearity of (9.25). Observe that we proved that $c_1 \neq 0$ is a necessary condition for the existence of solutions $(q^a; q^b)$ of the cocycle condition (8.14), depending

only on the first and higher derivatives of q^a . Existence of solutions follows from the fact that the Schwarzian derivative $\{q^a, q^b\}$ solves (8.14) and depends only on the first and higher derivatives of q^a .

The fact that $c_1 = 0$ implies $(q^a; q^b) = 0$, can be also seen by explicitly evaluating the coefficients c_n and d_n . These can be obtained using the same procedure considered above to prove that $c_1 \neq 0$. Namely, inserting the expansion (9.18) in (8.14) and using $q^c = q^b + \epsilon^{cb}(q^b)$, $\epsilon^{ac}(q^c) = \epsilon^{ab}(q^b) - \epsilon^{cb}(q^b)$ and $\epsilon^{bc}(q^c) = -\epsilon^{cb}(q^b)$, we obtain

$$c_n = (-1)^{n-1}c_1, \quad d_n = \frac{3}{2}(-1)^{n-1}(n-1)c_1, \quad (9.29)$$

which in fact are the coefficients one obtains expanding $c_1\{q + \epsilon f(q), q\}$. However, we now use only the fact that $c_1 \neq 0$, as the relation $(q + \epsilon f(q); q) = c_1\{q + \epsilon f(q), q\}$ can be proved without making the calculations leading to (9.29). Summarizing, we have

Lemma

The cocycle condition (8.14) admits solutions $(q^a; q^b)$ depending only on the first and higher derivatives of q^a , if and only if in the case $q^a = q^b + \epsilon^{ab}(q^b)$, one has $(q^a; q^b) = c_1\epsilon^{ab''' }(q^b) + \mathcal{O}^{ab}(\epsilon^2)$, with $c_1 \neq 0$.

We are now ready to prove that, up to a multiplicative constant and a coboundary term, the Schwarzian derivative is the unique solution of the cocycle condition (8.14). Let us first note that

$$[q^a; q^b] = (q^a; q^b) - c_1\{q^a; q^b\}, \quad (9.30)$$

satisfies the cocycle condition

$$[q^a; q^c] = \left(\partial_{q^c} q^b\right)^2 \left([q^a; q^b] - [q^c; q^b]\right). \quad (9.31)$$

In particular, since both $(q^a; q^b)$ and $\{q^a; q^b\}$ depend only on the first and higher derivatives of q^a , we have, as in the case of $(q + \epsilon f(q); q)$, that

$$[q + \epsilon f(q); q] = \tilde{c}_1 \epsilon f^{(3)}(q) + \mathcal{O}(\epsilon^2), \quad (9.32)$$

where either $\tilde{c}_1 \neq 0$ or $[q + \epsilon f(q); q] = 0$. However, since $\{q + \epsilon f(q); q\} = \epsilon f^{(3)}(q) + \mathcal{O}(\epsilon^2)$, by Eq.(9.12) we have $\tilde{c}_1 = 0$ and the Lemma yields $[q + \epsilon f(q); q] = 0$. Therefore, we have that the EP univocally implies the following central result

$$(q^a; q^b) = -\frac{\beta^2}{4m}\{q^a, q^b\}, \quad (9.33)$$

where for convenience we replaced c_1 by $-\beta^2/4m$. Observe that from a mathematical point of view we proved the following

Theorem

The cocycle condition (8.14) uniquely defines the Schwarzian derivative up to a global constant and a coboundary term.

We observe that despite some claims [39], we have not been able to find in the literature a complete and close proof of the above theorem.⁸

Let us now consider the issue of the classical limit. Comparing $\mathcal{W}^v(q^v) = (\partial_{q^v} q)^2 \mathcal{W}(q) + (q; q^v)$ with the classical case $\mathcal{W}^v(q^v) = (\partial_{q^v} q)^2 \mathcal{W}(q)$, we have that in the classical limit $(q^a; q^b) \longrightarrow 0$. Hence, by (9.33)

$$\frac{\beta^2}{4m} \{q, q^v\} \longrightarrow 0. \quad (9.34)$$

Thus β is precisely the parameter we are looking for. Since in the classical limit $Q \longrightarrow 0$, we have

$$\lim_{\beta \longrightarrow 0} Q = 0. \quad (9.35)$$

Note that in this limit $\mathcal{W}$ is kept fixed. Since for $Q \longrightarrow 0$ one obtains the CSHJE (2.15), we have

$$\lim_{\beta \longrightarrow 0} \mathcal{S}_0 = \mathcal{S}_0^{cl}. \quad (9.36)$$

From (8.8) and (9.33) it follows that $\mathcal{W}$ itself is a Schwarzian derivative

$$\mathcal{W}(q) = -\frac{\beta^2}{4m} \{q^0, q\}. \quad (9.37)$$

On the other hand, by (7.13) and (9.7) we have

$$\mathcal{W}(q) = \frac{\beta^2}{4m} \left(\{\mathcal{S}_0, q\} - \{e^{\frac{2i}{\beta} \mathcal{S}_0}, q\} \right) - Q(q), \quad (9.38)$$

so that by (9.37)

$$Q(q) = \frac{\beta^2}{4m} \left(\{\mathcal{S}_0, q\} - \{e^{\frac{2i}{\beta} \mathcal{S}_0}, q\} + \{q^0, q\} \right). \quad (9.39)$$

In particular, the correction to the CSHJE in the case of the state $\mathcal{W}^0$ has the form

$$Q^0(q^0) = \frac{\beta^2}{4m} \left(\{\mathcal{S}_0^0, q^0\} - \{e^{\frac{2i}{\beta} \mathcal{S}_0^0}, q^0\} \right), \quad (9.40)$$

where $\mathcal{S}_0 \equiv \mathcal{S}_0(q)$, $\mathcal{S}_0^0 \equiv \mathcal{S}_0^0(q^0)$. Observe that we still have to determine what q^0 is. Namely, we have to find the expression of the reduced action $\mathcal{S}_0^0(q^0)$. Since $\mathcal{S}_0^0(q^0) = \mathcal{S}_0(q)$ this will also give $q^0 = v_0(q)$. We will see that the unique possible solution is

$$q^0 = \frac{Ae^{\frac{2i}{\beta} \mathcal{S}_0(q)} + B}{Ce^{\frac{2i}{\beta} \mathcal{S}_0(q)} + D}, \quad (9.41)$$

so that, Eq.(9.37) and the Möbius symmetry $\{(Ah + B)/(Ch + D), x\} = \{h, x\}$, yield

$$\mathcal{W}(q) = -\frac{\beta^2}{4m} \{e^{\frac{2i}{\beta} \mathcal{S}_0}, q\}, \quad (9.42)$$

and by (9.39)

$$Q(q) = \frac{\beta^2}{4m} \{\mathcal{S}_0, q\}. \quad (9.43)$$

It follows from (7.13) and (9.43) that the equation for $\mathcal{S}_0$ we were looking for is

$$\frac{1}{2m} \left(\frac{\partial \mathcal{S}_0(q)}{\partial q} \right)^2 + V(q) - E + \frac{\beta^2}{4m} \{\mathcal{S}_0, q\} = 0. \quad (9.44)$$

⁸We thank D.B. Fuchs for a bibliographic comment concerning the above theorem.

9.3 Uniqueness of the solution

To show the uniqueness of the solution (9.42), that by (9.37) is equivalent to (9.41), we first set

$$Q = \frac{\beta^2}{4m} \{\mathcal{S}_0, q\} - g(q), \quad (9.45)$$

so that by (9.38)

$$\mathcal{W} = -\frac{\beta^2}{4m} \{e^{\frac{2i}{\beta}\mathcal{S}_0}, q\} + g(q). \quad (9.46)$$

We now consider, for any fixed $\mathcal{W}$, the $\beta \rightarrow 0$ limit of Q in (9.45). Suppose that $\lim_{\beta \rightarrow 0} \{\mathcal{S}_0, q\} = \{\lim_{\beta \rightarrow 0} \mathcal{S}_0, q\}$, so that

$$\lim_{\beta \rightarrow 0} \{\mathcal{S}_0, q\} = \{\mathcal{S}_0^{cl}, q\}. \quad (9.47)$$

The exception arises when $\{\mathcal{S}_0^{cl}, q\}$ is not defined so that one has to consider $\lim_{\beta \rightarrow 0} \{\mathcal{S}_0, q\}$. This happens, for example, in the case of the state $\mathcal{W}^0$ corresponding to $\mathcal{S}_0^{0cl} = c \text{nst}$. Let us then first consider an arbitrary state for which $\{\mathcal{S}_0^{cl}, q\}$ exists. By (9.47) we have

$$\lim_{\beta \rightarrow 0} \left(\frac{\beta^2}{4m} \{\mathcal{S}_0, q\} - g(q) \right) = \lim_{\beta \rightarrow 0} \frac{\beta^2}{4m} \{\mathcal{S}_0^{cl}, q\} - g^{cl}(q) = -g^{cl}(q), \quad (9.48)$$

and by (9.35)

$$g^{cl} = 0. \quad (9.49)$$

Next, note that since $\{\mathcal{S}_0, q\} \in \mathcal{P}$, it follows from (8.15) that $g(q)$ must transform as a quadratic differential under v -maps, that is consistency requires

$$g \in \mathcal{Q}. \quad (9.50)$$

On the other hand, the only quadratic differential that can be built from $\mathcal{S}_0$ has the form

$$g(q) = \frac{1}{4m} (\partial_q \mathcal{S}_0)^2 G(\mathcal{S}_0), \quad (9.51)$$

with G a function of $\mathcal{S}_0$. In other words, there is no way to construct a quadratic differential by means of higher derivatives of $\mathcal{S}_0$, as these terms would break the consistency condition (9.50). Furthermore, (7.11) implies $G(\mathcal{S}_0) = c$, where c is a constant. On the other hand, since by (9.51) c is dimensionless whereas β has the dimension of an action, Eq.(9.49) gives $c = 0$. Hence

$$g = 0. \quad (9.52)$$

The extension of (9.52) to arbitrary states simply follows from the observation that by (9.50) we have $g(q) \rightarrow g^v(q^v) = (\partial_q q^v)^{-2} g(q)$, so that by (9.52) $g^v(q^v) = 0$. In other words, (9.52) holds for all $\mathcal{W} \in \mathcal{H}$.

10 The Schrödinger equation

We have seen that the EP univocally leads to Eq.(9.44). In this section we will first see that this equation implies the Schrödinger equation, so that we will call Eq.(9.44) the Quantum Stationary HJ Equation (QSHJE). Next, we will consider the role of the wave-function in comparison with the appearance of a pair of linearly independent solutions of the SE arising in the present formulation. Furthermore, we will derive a dual SE in which $\mathcal{W}$ is replaced by (minus) the quantum potential Q . Furthermore, we will see that the self-dual state $\mathcal{W}^{sd}$, we selected by symmetry arguments, coincides with $\mathcal{W}^0$. Subsequently, we will consider the explicit form of the trivializing map, an investigation that will be further considered later on.

10.1 QSHJE and Schrödinger equation

It is easy to see that Eq.(9.44) implies the SE. It simply arises by considering Eqs.(4.19)–(4.21) with $h = e^{\frac{2i}{\beta}S_0}$. In particular, we immediately see that (9.44) yields

$$e^{\frac{2i}{\beta}S_0} = \frac{\psi^D}{\psi}, \quad (10.1)$$

where ψ^D and ψ are linearly independent solutions of the stationary SE

$$\left(-\frac{\beta^2}{2m} \frac{\partial^2}{\partial q^2} + V(q) \right) \psi = E\psi. \quad (10.2)$$

Thus, for the “covariantizing parameter” we have

$$\beta = \hbar, \quad (10.3)$$

where $\hbar = h/2\pi$ is Planck’s reduced constant. Therefore, in our formulation the SE arises as the equation which linearizes the QSHJE.

Although the QSHJE (9.44) is not familiar, it can be found in [15][40][41][42][43]. In this respect we note that (9.44) differs from other more familiar versions of the quantum HJ equations. A well-known version arises by identifying the wave-function with $Re^{\frac{i}{\hbar}\hat{S}_0}$, $R, \hat{S}_0 \in \mathbb{R}$, in the SE. As we will see, the basic difference between this stationary quantum HJ equation and (9.44) is that while in the standard approach one considers $Re^{\frac{i}{\hbar}\hat{S}_0}$ to be the wave-function, (9.44) is obtained by considering the wave-function as a linear combination of $Re^{\frac{i}{\hbar}\hat{S}_0}$ and $Re^{-\frac{i}{\hbar}\hat{S}_0}$ and S_0 is never a constant. In particular, in the cases in which the wave-function is proportional to a real function, the standard formulation leads to $\hat{S}_0 = cns$ even for non-trivial systems such as the harmonic oscillator. As a result, while the QSHJE has a well-defined classical limit, in the standard version one has situations in which quantum mechanically the reduced action is trivial even if its classical analogue is not. Another version of the quantum analogue of the HJ equation arises by setting $\psi = e^{\frac{i}{\hbar}\sigma}$, the first step of the WKB approximation. We will further consider these features later on.

10.2 Wave-function and Copenhagen School

Let us now consider the relationship between the solutions of (10.2) and the reduced action. By (10.1) it follows that any solution of (10.2) has the form

$$\psi = \frac{1}{\sqrt{\mathcal{S}'_0}} \left(A e^{-\frac{i}{\hbar} \mathcal{S}_0} + B e^{\frac{i}{\hbar} \mathcal{S}_0} \right). \quad (10.4)$$

In particular, by (10.1) the transformation $\tilde{\psi}^D = A\psi^D + B\psi$, $\tilde{\psi} = C\psi^D + D\psi$, is equivalent to

$$e^{\frac{2i}{\hbar} \mathcal{S}_0} \longrightarrow \frac{A e^{\frac{2i}{\hbar} \mathcal{S}_0} + B}{C e^{\frac{2i}{\hbar} \mathcal{S}_0} + D}, \quad (10.5)$$

which is an invariance of $\mathcal{W} = -\frac{\hbar^2}{4m} \{e^{\frac{2i}{\hbar} \mathcal{S}_0}, q\}$. Note that since ψ^D and ψ are arbitrary linearly independent solutions of (10.2), we have by (10.1) that Eq.(10.5) is equivalent to

$$e^{\frac{2i}{\hbar} \mathcal{S}_0} = \frac{A\psi^D + B\psi}{C\psi^D + D\psi}. \quad (10.6)$$

This expression shows that a distinguished feature of the present formulation concerns the appearance of both ψ^D and ψ . This is the signal of the underlying Legendre duality which reflects in the ψ^D – ψ duality, a consequence of the fact that the SE is a second-order linear differential equation. The appearance of both solutions in relevant formulas is unusual in the standard approach. So, since in the standard approach the solution of the SE one considers is the wave-function, it follows that for bound states, only the $L^2(\mathbb{R})$ solution is considered. On the other hand, as we will see in sect.17, Wronskian arguments show that if the SE has an $L^2(\mathbb{R})$ solution, then any other linearly independent solution of the SE cannot belong to $L^2(\mathbb{R})$.⁹ Therefore, the fact that for some system the wave-function belongs to the $L^2(\mathbb{R})$ space, does not mean that for the same energy level the SE has no divergent solutions. In other words, one should not confuse the fact that the wave-function for bound states is unique, corresponding to the non-degeneration theorem of the spectrum for bound states, with the wrong sentence: “the SE for bound states has only one solution”. Also note that the fact that the divergent solution cannot be the wave-function does not imply that it cannot appear in relevant expressions which in fact characterize the present formulation. In this context it is important to observe that our approach derives from the EP and, as such, we are deriving not only QM from a first principle, which looks different from the axioms of the Copenhagen School, but, as we will see, also its interpretation is quite different. On the other hand, to be consistent, we have to reproduce the basic experimental predictions of QM. As we will see, two basic phenomena, such as the tunnel effect and energy quantization, are in fact a direct consequence of the EP. It is well-known that in the usual approach these phenomena are a consequence of the axiomatic interpretation of the wave-function in terms of probability amplitude. In particular, since in the classically forbidden regions the wave-function does not vanish identically, this interpretation implies a non-vanishing probability of finding

⁹Uniqueness of the wave-function for bound states, may induce to believe that divergent solutions of the SE appear only in the case of a non-physical energy value. Thus, one may erroneously understand that in this context, by linearly independent solution it is meant a solution of the same SE but with a different value of the energy.

a particle in such regions. The probabilistic interpretation also implies that for suitable potentials the wave-function must belong to $L^2(\mathbb{R})$.

At this stage one may wonder how it is possible that tunnel effect and energy quantization also arise in our approach in which divergent solutions appear. Let us denote by Σ_E the space of solutions of the SE (10.2) corresponding to the energy level E and consider the case in which

$$V(q) - E \geq \begin{cases} P_-^2 > 0, & q < q_-, \\ P_+^2 > 0, & q > q_+, \end{cases} \quad (10.7)$$

where q_- (q_+) is the lowest (highest) value of q for which $V(q) - E$ changes sign. We denote by σ^{BD} the associated physical discrete spectrum. Let us then explain in which sense, in the usual approach, the divergent solutions associated to (10.7) cannot be considered physical ones. First note that the precise sentence is that in the case (10.7) the wave-function, and therefore a particular solution of the SE, must belong to $L^2(\mathbb{R})$. From a physical point of view, what is relevant is that this request selects the energy spectrum. In particular, the content of the ‘‘Copenhagen axiom’’ that the physical spectrum associated to (10.7) is determined by the condition

$$\textit{The wave - function belongs to } L^2(\mathbb{R}), \quad (10.8)$$

can be separated in two parts. The physically relevant part, in which the concept of wave-function does not appear, is

$$\textit{The SE admits an } L^2(\mathbb{R}) \textit{ solution}, \quad (10.9)$$

which by itself implies the quantized spectrum, that is

$$E \in \sigma^{BD} \iff \psi \in L^2(\mathbb{R}), \quad (10.10)$$

for some $\psi \in \Sigma_E$. Therefore, in the context of the energy spectrum, the meaning of the wave-function may be seen as a possible way to impose the physical request (10.9), with the remnant part of (10.8) reducing to the meaning of the wave-function and therefore to its probabilistic interpretation. In other words, in order to select the physical spectrum we only need (10.9) which contains less assumptions than those in (10.8). This is a first signal that the Copenhagen interpretation is sufficiently sophisticated to well hide the underlying nature of QM. In this respect we will see that the physical request (10.9) directly follows from the EP without using the axiomatic interpretation of the wave-function! Therefore, as far as the problem of selecting the physical spectrum is concerned, the role of (10.8) reduces to selecting the values of E for which the SE admits an $L^2(\mathbb{R})$ solution. Since if for a fixed E the SE has an $L^2(\mathbb{R})$ solution, then any other linearly independent solution of the SE cannot belong to $L^2(\mathbb{R})$, we have that the effect of (10.8) on the selection of the spectrum is equivalent to

$$E \notin \sigma^{BD} \iff \psi \notin L^2(\mathbb{R}), \quad (10.11)$$

$\forall \psi \in \Sigma_E$. As we said, the fact that the wave-function must belong to $L^2(\mathbb{R})$ does not mean that for the same E all solutions of the SE belong to $L^2(\mathbb{R})$. It is true the opposite as, if for a fixed E the SE

has an $L^2(\mathbb{R})$ solution, then any other linearly independent solution cannot belong to $L^2(\mathbb{R})$. So, for example, if for a potential well there is a value of E for which the SE has a solution ψ which decreases exponentially at both spatial infinities, then any other linearly independent solution ψ^D , therefore solution of the same SE with the same E but with $\psi^D \not\propto \psi$, grows exponentially at $q = \pm\infty$.

The fact that in our formulation both ψ^D and ψ appear, is a consequence of the underlying Legendre duality implied by the EP. In fact, in the same way we obtained p - q duality in considering the canonical equations, here we have a manifest ψ^D - ψ duality. This can be nicely expressed by means of the Legendre brackets introduced in [11]. In particular, repeating the construction in (5.15)–(5.17), we see that considering $\mathcal{F}(\psi^2)$ as generating function, defined by $\psi^D = \partial\mathcal{F}/\partial\psi$, we have

$$\{\psi, \psi\}_{(q)} = 0 = \{\psi^D, \psi^D\}_{(q)}, \quad \{\psi, \psi^D\}_{(q)} = 1. \quad (10.12)$$

We stress that this duality structure, which, due to the preferred nature played by the wave-function, is missing in the usual formulation of QM, is a basic feature which distinguishes the present formulation from the one of the Copenhagen School.

10.3 $\mathcal{W} \longleftrightarrow Q$ and the dual Schrödinger equation

An interesting question is to seek a direct relation between $\mathcal{W}$ and Q . To be more specific, note that while the QSHJE (9.44) connects $\mathcal{W}$ and Q and $\mathcal{S}'_0$, it would be desirable to find a differential equation involving $\mathcal{W}$ and Q only. To this end we observe that (9.7) implies

$$\mathcal{S}_0 = \alpha \int_{q_0}^q dx (\mathcal{W} + Q)^{1/2}, \quad (10.13)$$

where $\alpha^2 = -2m$. As $Q = \hbar^2 \{\mathcal{S}_0, q\}/4m$, we see that the equation relating $\mathcal{W}$ and the quantum potential is $Q = \hbar^2 \{\int_{q_0}^q dx (\mathcal{W} + Q)^{1/2}, q\}/4m$, that is

$$Q = \frac{\hbar^2}{8m} \left(\frac{\mathcal{W}'' + Q''}{\mathcal{W} + Q} - \frac{5}{4} \frac{(\mathcal{W}' + Q')^2}{(\mathcal{W} + Q)^2} \right). \quad (10.14)$$

Note that adding $\mathcal{W}$ to both sides of (10.14) we obtain a similar expression with now $\mathcal{W}$ expressed in terms of $\mathcal{W} + Q$ and its derivatives. This complementary role between $\mathcal{W}$ and Q is not an accident. To show this, we write down again Eq.(4.40) we derived on the basis of duality arguments

$$\left(\frac{\hbar^2}{2m} \frac{\partial^2}{\partial q^2} + Q \right) \phi = 0. \quad (10.15)$$

This shows that Q plays a similar role to that played by $\mathcal{W}$ in the SE. It is easy to see that if ϕ^D and ϕ are solutions of (10.15), then

$$\mathcal{S}_0 = \frac{A\phi^D + B\phi}{C\phi^D + D\phi}. \quad (10.16)$$

It follows by (10.6) that the solutions of (10.16) are related to the solutions of the SE by

$$\gamma_\psi(\psi^D/\psi) = e^{\frac{2i}{\hbar}\gamma_\phi(\phi^D/\phi)}, \quad (10.17)$$

where γ_ψ and γ_ϕ denote two Möbius transformations.

10.4 $\mathcal{W}^{sd} = \mathcal{W}^0$ and existence of the Legendre transformation

We now explicitly show how the solution, that is the QSHJE (9.44), solves the problem of defining the Legendre transformation for any state, so that the $\mathcal{S}_0$ and $\mathcal{T}_0$ descriptions exist for *any* state. Let us first consider the state $\mathcal{W}^0$. In this case we have

$$\frac{1}{2m} \left(\frac{\partial \mathcal{S}_0^0}{\partial q^0} \right)^2 + \frac{\hbar^2}{4m} \{ \mathcal{S}_0^0, q^0 \} = 0, \quad (10.18)$$

that by (9.7) is equivalent to

$$\{ e^{\frac{2i}{\hbar} \mathcal{S}_0^0}, q^0 \} = 0. \quad (10.19)$$

A possible solution is

$$e^{\frac{2i}{\hbar} \mathcal{S}_0^0} = \gamma_q q^0. \quad (10.20)$$

Note that the Legendre transform of $\mathcal{S}_0^0$ in (10.20) is well-defined. Therefore, the QSHJE solves the problem of defining the Legendre transform of the reduced action one meets in CM. Now observe that the self-dual states (6.40), which arose as the highest symmetric points of the Legendre pair $\mathcal{S}_0$ – $\mathcal{T}_0$, have the structure of (10.20). Hence, we can identify the reduced action $\mathcal{S}_0^0$ associated to the state $\mathcal{W}^0$ with the distinguished self-dual state (8.35)! In other words, we can set

$$\gamma_{sd} = \pm \frac{\hbar}{2i}, \quad (10.21)$$

so that

$$\mathcal{W}^{sd} = \mathcal{W}^0, \quad (10.22)$$

and

$$\mathcal{S}_0^0(q^0) = \mathcal{S}_0^{sd}(q^{sd}) = \pm \frac{\hbar}{2i} \ln \gamma_q q^0, \quad q^{sd} = q^0. \quad (10.23)$$

We note that the general solution of (10.19) is

$$\mathcal{S}_0^0 = \frac{\hbar}{2i} \ln \left(\frac{Aq^0 + B}{Cq^0 + D} \right), \quad (10.24)$$

which will be useful in considering both the properties of the physical solutions with real p and the structure of the trivializing map.

The relevant case in which the Legendre transformation is not defined is for $\mathcal{S}_0 = Aq + B$. According to Eq.(9.42), this corresponds to $\mathcal{W} = -A^2/2m$, that for $V = 0$ corresponds to a free system with energy $E = A^2/2m$. On the other hand, by the Möbius invariance of the Schwarzian derivative

$$\mathcal{W}(q) = -\frac{\hbar^2}{4m} \{ e^{\frac{2i}{\hbar} \mathcal{S}_0}, q \} = -\frac{\hbar^2}{4m} \left\{ \frac{Ae^{\frac{2i}{\hbar} \mathcal{S}_0} + B}{Ce^{\frac{2i}{\hbar} \mathcal{S}_0} + D}, q \right\}, \quad (10.25)$$

$AD - BC \neq 0$. It follows that if $\mathcal{S}_0$ is a solution of Eq.(9.44), we have that also

$$\tilde{\mathcal{S}}_0 = \frac{\hbar}{2i} \ln \left(\frac{Ae^{\frac{2i}{\hbar} \mathcal{S}_0} + B}{Ce^{\frac{2i}{\hbar} \mathcal{S}_0} + D} \right), \quad (10.26)$$

is a solution in agreement with (10.6). Therefore, we see that in QM the kinetic term and the quantum term $Q(q)$ are intimately related: they mix under the symmetry (10.25) of $\mathcal{W}$. This allows us to solve the problem of defining the Legendre transformation in the case in which $\mathcal{W}$ is a constant. Indeed, besides $\mathcal{S}_0 = \pm\sqrt{2mE}q$, the equation

$$\frac{1}{2m} \left(\frac{\partial \mathcal{S}_0}{\partial q} \right)^2 - E + \frac{\hbar^2}{4m} \{\mathcal{S}_0, q\} = 0, \quad (10.27)$$

has solutions

$$\mathcal{S}_0 = \frac{\hbar}{2i} \ln \left(\frac{Ae^{\frac{2i}{\hbar}\sqrt{2mE}q} + B}{Ce^{\frac{2i}{\hbar}\sqrt{2mE}q} + D} \right), \quad (10.28)$$

where the constants are chosen in such a way that $\mathcal{S}_0 \not\propto q$. Since $\mathcal{S}_0^{cl} = \pm\sqrt{2mE}q$, we have

$$\lim_{\hbar \rightarrow 0} \ln \left(\frac{Ae^{\frac{2i}{\hbar}\sqrt{2mE}q} + B}{Ce^{\frac{2i}{\hbar}\sqrt{2mE}q} + D} \right)^{\frac{\hbar}{2i}} = \pm\sqrt{2mE}q, \quad (10.29)$$

implying that A, B, C, D should depend on $\hbar$. This important point will be discussed in detail later on (see subsection 16.1). For the time being we just observe that the standard solution $\mathcal{S}_0 = \pm\sqrt{2mE}q$ considered in literature has the property that in the $E \rightarrow 0$ limit one obtains $\mathcal{S}_0 = 0$ for which both the Schwarzian derivative and the Legendre transformation are not defined.

10.5 The trivializing map

We now derive the VT $q \rightarrow q^0$ such that $\mathcal{W} \rightarrow \mathcal{W}^0$. Although, as we will see, there is a direct way to find it, it is instructive to first consider the relevant transformation properties with the derived explicit expression for $\mathcal{W}$. First note that under the v -map $q \rightarrow q^b = v^b(q)$, we have

$$\{e^{\frac{2i}{\hbar}\mathcal{S}_0(q^b)}, q^b\} = \{e^{\frac{2i}{\hbar}\mathcal{S}_0(q)}, q^b\} = \left(\partial_{q^b} q \right)^2 \left[\{e^{\frac{2i}{\hbar}\mathcal{S}_0(q)}, q\} - \{q^b, q\} \right], \quad (10.30)$$

that is

$$\mathcal{W}^b(q^b) = \left(\partial_{q^b} q \right)^2 \left[\mathcal{W}(q) + \frac{\hbar^2}{4m} \{q^b, q\} \right]. \quad (10.31)$$

It follows that if

$$q^b = \frac{Ae^{\frac{2i}{\hbar}\mathcal{S}_0(q)} + B}{Ce^{\frac{2i}{\hbar}\mathcal{S}_0(q)} + D}, \quad (10.32)$$

then, according to (10.30), we have $\{e^{\frac{2i}{\hbar}\mathcal{S}_0(q^b)}, q^b\} = 0$, that is q^b is a Möbius transformation of q^0 and

$$\mathcal{W}^b(q^b) = \mathcal{W}^0(q^0). \quad (10.33)$$

It follows that

$$q \rightarrow q^0 = \frac{Ae^{\frac{2i}{\hbar}\mathcal{S}_0(q)} + B}{Ce^{\frac{2i}{\hbar}\mathcal{S}_0(q)} + D}, \quad (10.34)$$

is the solution of the inversion problem Eq.(7.2). Note that this result can be also directly derived by comparing (7.2) and (10.24).

11 Quantum HJ equation and the reality condition

We saw that the formulation exhibits $\mathcal{S}_0$ – $\mathcal{T}_0$ duality as $\mathcal{S}_0 = Aq + B$ does not belong to the space $\mathcal{K}$ of all possible $\mathcal{S}_0$'s. Due to the Möbius invariance of the Schwarzian derivative, instead of $\mathcal{S}_0 = \sqrt{2mE}q$, which corresponds to $\mathcal{W} = -E$, we can choose for $\mathcal{S}_0$ the expression given in (10.28). Similarly, we have seen that the state $\mathcal{W}^0$ corresponds to $\mathcal{S}_0^0 = \frac{\hbar}{2i} \ln(Aq^0 + B)/(Cq^0 + D)$. Therefore, while setting $\psi = Re^{\frac{i}{\hbar}\hat{\mathcal{S}}_0}$ the solutions corresponding to the states with $\mathcal{W} = \text{const}$ coincide with the classical ones, here we have a basic difference related to the existence of the Legendre transform of $\mathcal{S}_0$ for any system. Here, we first discuss these features in connection with the standard version of the quantum HJ equation, and then we will consider the classical limit. We will see that, while there are some subtleties involved in considering such a limit for the standard versions of the quantum HJ equation, in the case of Eq.(9.44) there is a natural criterion for defining it. We will conclude this section by considering the version of the quantum HJ equation used in the WKB approximation. We will show that also in this case the phase cannot be considered as the quantum reduced action.

11.1 Quantum HJ equation and reality of R and $\hat{\mathcal{S}}_0$

We now compare the QSHJE (9.44) with a more familiar version. Let us denote by ψ the Schrödinger wave-function. While Eq.(9.44) is written in terms of $\mathcal{S}_0$ only, setting

$$\psi(q) = Re^{\frac{i}{\hbar}\hat{\mathcal{S}}_0}, \quad (11.1)$$

with the condition

$$(R, \hat{\mathcal{S}}_0) \in \mathbb{R}^2, \quad (11.2)$$

in Eq.(10.2), leads to the system of real equations

$$\frac{1}{2m} \left(\frac{\partial \hat{\mathcal{S}}_0}{\partial q} \right)^2 + V - E - \frac{\hbar^2}{2mR} \frac{\partial^2 R}{\partial q^2} = 0, \quad (11.3)$$

$$\frac{\partial}{\partial q} \left(R^2 \frac{\partial \hat{\mathcal{S}}_0}{\partial q} \right) = 0. \quad (11.4)$$

Note that to identify the wave-function with $Re^{\frac{i}{\hbar}\hat{\mathcal{S}}_0}$ is an assumption. We will see that in general it should be identified with the linear combination $R(Ae^{-\frac{i}{\hbar}\hat{\mathcal{S}}_0} + Be^{\frac{i}{\hbar}\hat{\mathcal{S}}_0})$.

We now show that (9.44) is the natural quantum analogue of the CSHJE. First of all, when $\hat{\mathcal{S}}_0$ is constant, (11.4) degenerates and (11.3) reduces to $\mathcal{W} = \hbar^2 \partial_q^2 R / 2mR$. Since $\hat{\mathcal{S}}_0$ does not appear in this equation, its classical limit does not arise in a natural way as for (9.44). In this respect it is worth noticing that there are many cases with $\hat{\mathcal{S}}_0 = \text{const}$ despite the fact that $\mathcal{S}_0^{cl}$, as much as $\mathcal{S}_0$, is non-trivial, most notably the harmonic oscillator. More generally, all the states for which the wave-function ψ is proportional to a real function have a trivial $\hat{\mathcal{S}}_0$ (except for possible nodes of the wave-function where $R = 0$). It follows that for bound states the conjugate momentum would vanish.

In going from the QSHJE to the SE we lose some information about $\mathcal{S}_0$. This is a consequence of the invariance of $\mathcal{W}$ under a Möbius transformation of $e^{\frac{2i}{\hbar}\mathcal{S}_0}$. We also note that with the condition

(11.2), which implies the existence of states with $\hat{\mathcal{S}}_0 = \text{const}$, both the EP and the $\mathcal{S}_0$ - $\mathcal{T}_0$ Legendre duality are not naturally implemented. Later on we will discuss the reality condition for $\mathcal{S}_0$ and the identification of the wave-function in terms of $\mathcal{S}_0$. For the time being we consider (10.1) and do not fix any reality restriction on $\mathcal{S}_0$. The derivation of the connection between (11.3)(11.4) and (9.44) implies a distinction between two cases. Let us denote by ψ^D a solution of the SE linearly independent from ψ . We may have either $\bar{\psi} \not\propto \psi$ or $\bar{\psi} \propto \psi$. In the first case, since $\mathcal{W}$ is real, we can set

$$\psi^D = \bar{\psi}, \quad (11.5)$$

that is $\psi^D = R e^{\frac{i}{\hbar} \hat{\mathcal{S}}_0}$, $\psi = R e^{-\frac{i}{\hbar} \hat{\mathcal{S}}_0}$ and by (10.1)

$$\mathcal{S}_0 = \hat{\mathcal{S}}_0 + \pi k \hbar, \quad (11.6)$$

$k \in \mathbb{Z}$. The continuity equation (11.4) gives $R \propto 1/\sqrt{\mathcal{S}'_0}$. Therefore, by (10.1), that together with the condition (11.5) fixes some of the arbitrariness of $\mathcal{S}_0$, we have

$$Q = \frac{\hbar^2}{4m} \{\mathcal{S}_0, q\} = -\frac{\hbar^2}{2mR} \frac{\partial^2 R}{\partial q^2}, \quad (11.7)$$

and (11.3) corresponds to (9.44). The difference between (9.44) and (11.3)(11.4) becomes relevant when $\bar{\psi} \propto \psi$. In particular, (11.1) and the condition (11.2) imply that in this case $\hat{\mathcal{S}}_0$ must be a constant. Since this constant can be absorbed by a normalization of ψ , we can choose $\hat{\mathcal{S}}_0 = 0$. Therefore, up to a normalization, we have

$$\psi = R, \quad (11.8)$$

which is real. Since also

$$\psi^D = R \int_{q_0}^q dx R^{-2}, \quad (11.9)$$

is real, we have that if we identify ψ and ψ^D in (11.8)(11.9) with those in (10.1), then $\mathcal{S}_0$ would be purely imaginary, that is

$$\mathcal{S}_0 = \frac{\hbar}{2i} \ln \int_{q_0}^q dx R^{-2}, \quad (11.10)$$

and

$$\left(\frac{\partial \mathcal{S}_0}{\partial q} \right)^2 + \frac{\hbar^2}{2} \{\mathcal{S}_0, q\} = -\frac{\hbar^2}{R} \frac{\partial^2 R}{\partial q^2}. \quad (11.11)$$

It follows that (9.44) is equivalent to

$$\frac{\hbar^2}{2mR} \frac{\partial^2 R}{\partial q^2} = V - E. \quad (11.12)$$

On the other hand, for $\hat{\mathcal{S}}_0 = \text{const}$, (11.4) degenerates and (11.3) corresponds to (11.12) (and by (11.11) to (9.44)). Later on we will derive the real physical solutions of $\mathcal{S}_0$.

11.2 The classical limit

We saw that if $\bar{\psi} \propto \psi$, then $\hat{\mathcal{S}}_0 = \text{const}$. It follows that $\mathcal{S}_0^{cl}$, which is never a constant unless $\mathcal{W} = 0$, should arise in a rather involved way from (11.3). In particular, the classical limit of (11.3) manifestly does not correspond to the CSHJE. Usually, by quantum potential it is meant the term

$$\hat{Q} = -\frac{\hbar^2}{2mR} \frac{\partial^2 R}{\partial q^2}. \quad (11.13)$$

However, when $\bar{\psi} \propto \psi$, Q and $\hat{Q}$ have different structures. In particular, by (11.11) we have

$$\hat{Q} = Q + \frac{1}{2m} \left(\frac{\partial \mathcal{S}_0}{\partial q} \right)^2. \quad (11.14)$$

We will see that by itself the reality condition (11.2) is not an obstacle to getting the classical limit. The point is that this condition, together with the *assumption* that the wave-function takes the form (11.1), causes difficulties in getting the classical limit of (11.3)(11.4). Thus, since if (11.1) is a solution of the SE, then also

$$\psi = R \left(A e^{-\frac{i}{\hbar} \hat{\mathcal{S}}_0} + B e^{\frac{i}{\hbar} \hat{\mathcal{S}}_0} \right), \quad (11.15)$$

is a solution, we should investigate whether there are suitable A and B leading to a consistent classical limit for (11.3)(11.4). Suppose that the wave-function is proportional to a real function, *i.e.* $\bar{\psi} \propto \psi$. In the case of (11.1) this would imply $\hat{\mathcal{S}}_0 = \text{const}$, so that, unless $\mathcal{W} = 0$, one has $\lim_{\hbar \rightarrow 0} \hat{\mathcal{S}}_0 \neq \mathcal{S}_0^{cl}$. With the representation (11.15) the reality condition $\bar{\psi} \propto \psi$ would translate to a condition on the constants A and B rather than on $\hat{\mathcal{S}}_0$. In this way one can obtain a non-trivial $\hat{\mathcal{S}}_0$ and then, as we will see, a consistent classical limit for $\hat{\mathcal{S}}_0$. We note that the possibility of taking the linear combination (11.15), from which the classical limit may be defined, is a consequence of the linearity of the SE. On the other hand, this linearity is nothing but the manifestation of the invariance of $\mathcal{W}$ under Möbius transformations of $\exp(2i\mathcal{S}_0/\hbar)$. Therefore, we see once again that the Möbius symmetry, which we already met in considering Legendre duality and the related canonical equations, plays a basic role.

Let us further discuss the issue of the classical limit for the version (11.3)(11.4) of the quantum HJ equation. As we observed, the case $\hat{\mathcal{S}}_0 = 0$ is quite different from the classical situation. For example, for the classical harmonic oscillator of energy E one has

$$\frac{\partial \mathcal{S}_0^{cl}}{\partial q} = \pm m\omega(a^2 - q^2)^{1/2}, \quad (11.16)$$

where $a = (2E/m\omega^2)^{1/2}$ is the amplitude of motion. In the quantum case we have

$$\hat{\mathcal{S}}_0 = 0, \quad (11.17)$$

and by (11.12)

$$\hat{Q} = E_n - \frac{1}{2}m\omega^2 q^2, \quad (11.18)$$

where $E_n = (n + 1/2)\hbar\omega$ denotes the energy level of the quantum oscillator.¹⁰ This example shows that the condition (11.2) leads to a description which has nothing in common with the classical one.

We have seen that these features, related to the condition (11.2), do not arise in considering (9.44). In particular, while Q is the universal quantum correction to the CSHJE, this is not the case for $\hat{Q}$, as there are physical systems for which $\hat{Q}$ cannot be seen as quantum correction to the CSHJE. In fact, we saw that for all the states for which $\bar{\psi} \propto \psi$, one has $\hat{S}_0 = \text{const}$. Therefore, troubles in considering the classical limit of (11.3)(11.4) should arise for all bound states. See [41][44][45][46] for related topics.

As observed in [41], in some textbooks the classical limit of (11.3)(11.4) is understood in a naive sense. The standard argument is that due to the $\hbar^2$ factor in $\hat{Q}$ one has that in the $\hbar \rightarrow 0$ limit $\hat{Q}$ vanishes. We have explicitly seen that this is not generally the case. This can be also directly seen by observing that the fact that R and $\hat{S}_0$ in (11.3)(11.4) depend on $\hbar$ implies that taking the classical limit requires a rather different procedure. Furthermore, even admitting that $\lim_{\hbar \rightarrow 0} \hat{Q} = 0$, it remains to understand what the meaning of Eq.(11.4) is in the $\hbar \rightarrow 0$ limit. We saw that all these questions are naturally solved in the version (9.44) of the quantum HJ equation. In particular, while in general $\lim_{\hbar \rightarrow 0} \hat{Q} \neq 0$, in our formulation we have by construction that

$$\lim_{\hbar \rightarrow 0} Q = 0, \quad (11.19)$$

so that

$$\lim_{\hbar \rightarrow 0} \mathcal{S}_0 = \mathcal{S}_0^{cl}. \quad (11.20)$$

In particular, this means that CM is seen as an approximate description of QM. In other words, there are no two worlds, a classical and a quantum one, but only the quantum one with the classical phase $\hbar \rightarrow 0$ corresponding to an approximation. In this framework, it is relevant that there exists a QSHJE different from that considered in Bohm's theory [4]. In particular, while the quantum and classical harmonic oscillators, as any other system with real wave-function, seem to belong to two different worlds, this is not the case for the QSHJE. We also observe that considering CM as an approximation of QM would suggest a reconsideration of the collapse of the wave-function.

Concerning $\hat{Q}$, we note that by (11.12) it follows that in the case in which $\bar{\psi} \propto \psi$, one has

$$\lim_{\hbar \rightarrow 0} \hat{Q} = -V + \lim_{\hbar \rightarrow 0} E. \quad (11.21)$$

In this context we observe that in our approach $\mathcal{W}$ is kept fixed in the classical limit. In this respect the difference between CM and QM resides in the fact that in the quantum case there are possible constraints on the structure of the spectrum σ^{qu} , so that in general

$$\sigma^{qu} \subset \sigma^{cl}. \quad (11.22)$$

This means that any quantum energy level is an admissible one in CM. Of course, for a given E one in general has $\mathcal{S}_0^{cl} \neq \mathcal{S}_0$, so that for a given E one may have regions of space which are classically forbidden, but this does not exclude that the associated CSHJE makes sense. It is sufficient that

¹⁰See [41] for a discussion of the quantum harmonic oscillator in the framework of Eqs.(11.3)(11.4).

$\mathcal{W} \leq 0$ in some region, in order for the CSHJE to admit solutions. On the other hand, if $\mathcal{W} > 0$, $\forall q$, then E belongs neither to σ^{qu} nor σ^{cl} . Therefore, Eq.(11.22) is satisfied for any system. We also note that, as CM is an approximation, the actual observable physical spectrum is σ^{qu} . Therefore, from a fixed $\mathcal{W}$ in the QSHJE, it makes sense to consider the classical limit with $\mathcal{W}$, and therefore E , fixed. In particular, the CSHJE is naturally obtained from Eq.(9.44) with potential V and energy $E \in \sigma^{qu} \subset \sigma^{cl}$. This makes the classical limit (11.19) consistent and well-defined. Therefore, the QSHJE (9.44) admits a universal criterion for defining the classical limit. Note that while keeping E fixed is the natural procedure to get the classical limit, this does not eliminate the difficulties in considering the classical limit of the standard version of the quantum HJ equation (11.3)(11.4). In particular, in the case in which $\bar{\psi} \propto \psi$ we have that the classical limit of $\hat{Q}$, keeping E fixed, is

$$\lim_{\hbar \rightarrow 0} \hat{Q} = E - V = \hat{Q}, \quad (11.23)$$

that is $\hat{Q}$ and its classical limit coincide. Since $\hat{Q}$ does not vanish in the classical limit, unless $V = E$, it seems more appropriate to call Q “quantum potential” rather than $\hat{Q}$.

11.3 The WKB approximation and the quantum HJ equation

It is well-known that another version of the quantum HJ equation arises in considering the WKB approximation. This is obtained by setting $\psi = e^{\frac{i}{\hbar}\sigma}$, in the SE, so that ($' \equiv \partial_q$)

$$\frac{1}{2m}\sigma'^2 - \frac{i\hbar}{2m}\sigma'' = E - V(q), \quad (11.24)$$

and $\sigma = \sigma_0 + \frac{\hbar}{i}\sigma_1 + \left(\frac{\hbar}{i}\right)^2\sigma_2 + \dots$. The first three terms are $\sigma_0 = \pm \int_{q_0}^q dx p^{cl}$, $\sigma_1 = -\frac{1}{2} \ln p^{cl}$, $\sigma_2 = \frac{m}{4} \frac{F}{p^{cl3}} + \frac{m^2}{8} \int^q dx \frac{F^2}{p^{cl5}}$, where $p^{cl} = \sqrt{2m(E - V)}$, denotes the modulus of the classical momentum and $F = -V' = p^{cl}p^{cl'}/m$. We have seen that the Schwarzian derivative, which determines the quantum correction in (9.44), arises quite naturally in our formulation. We note that it also appears in considering the quantum correction σ_2 , that is $\sigma'_2 = \{\sigma_0, q\}/4p^{cl}$.

Also in the case of (11.24) one has to consider non-trivial questions concerning the classical limit. Similarly, if one considers the wave-function in the WKB approximation ψ_{WKB} , and then defines the quantum potential in a similar way to $\hat{Q}$, that is with $R = |\psi_{WKB}|$, one still meets similar problems to those considered with the version Eqs.(11.3)(11.4). We refer to [41] for a discussion concerning similar features. Here we just observe that a property of (11.24) is that it implies $\text{Im } \sigma \neq 0$ unless $\sigma = Aq + B$. On the contrary, we will see that the QSHJE (9.44) admits real solutions.

12 The wave equation of Classical Mechanics

In this section we will see that the relationship between the QSHJE and the SE indicates the way to associate to the CSHJE a wave equation. The latter differs from the SE for the addition to V of a non-linear term whose structure is the same of the quantum potential but with $\mathcal{S}_0$ replaced by

$\mathcal{S}_0^{cl}$. Remarkably, this equation permits an interpretation of the “classical wave-function” in terms of probability amplitude. We will conclude this section by discussing the different properties of the $\mathcal{W}$ states in CM and QM.

12.1 The wave-function of Classical Mechanics

A feature of our formulation is that the existence of both the classical and quantum versions of the HJ equation suggests considering a wave-function for CM. A similar problem was considered by Schiller and Rosen [47]. However, they derived the SE for Classical Statistical Mechanics, using a suitable modification of Eqs.(11.3)(11.4) (see also sect.2.6 in [41]). Here, we consider a different approach. The idea is to add the vanishing term

$$0 = \frac{\delta^2}{4m} \{\mathcal{S}_0^{cl}, q\} - \frac{\delta^2}{4m} \{\mathcal{S}_0^{cl}, q\}, \quad (12.1)$$

in the CSHJE (2.15), that is

$$\frac{1}{2m} \left(\frac{\partial \mathcal{S}_0^{cl}}{\partial q} \right)^2 + \mathcal{W}(q) - \frac{\delta^2}{4m} \{\mathcal{S}_0^{cl}, q\} + \frac{\delta^2}{4m} \{\mathcal{S}_0^{cl}, q\} = 0. \quad (12.2)$$

Similarly to the way one derives the SE from QSHJE (9.44), one can derive the SE for CM by considering $\mathcal{W}(q) - \delta^2 \{\mathcal{S}_0^{cl}, q\}/4m$ as an effective potential. We have

$$e^{\frac{2i}{\delta} \mathcal{S}_0^{cl}} = \frac{\psi_{cl}^D}{\psi_{cl}}, \quad (12.3)$$

where ψ_{cl}^D and ψ_{cl} are linearly independent solutions of

$$\left(-\frac{\delta^2}{2m} \frac{\partial^2}{\partial q^2} + V(q) - \frac{\delta^2}{4m} \{\mathcal{S}_0^{cl}, q\} \right) \psi_{cl} = E \psi_{cl}. \quad (12.4)$$

This equation can be interpreted as the wave-equation of CM.

12.2 Probabilistic interpretation of the classical wave-function

Observe that since for the free particle at rest one has $\mathcal{S}_0^{0cl} = \text{const}$, it follows that the Schwarzian derivative $\{\mathcal{S}_0^{0cl}, q\}$ is not defined in this case. This is solved in QM since the constant function does not belong to $\mathcal{K}$. Another property of Eq.(12.4) is that since an arbitrary solution has the form

$$\psi_{cl} = \frac{1}{\sqrt{\mathcal{S}_0^{cl'}}} \left(A e^{-\frac{i}{\delta} \mathcal{S}_0^{cl}} + B e^{\frac{i}{\delta} \mathcal{S}_0^{cl}} \right), \quad (12.5)$$

one has that the solutions (12.5) with either $A = 0$ or $B = 0$, that is

$$\psi_{cl} = \frac{A}{\sqrt{\mathcal{S}_0^{cl'}}} e^{\pm \frac{i}{\delta} \mathcal{S}_0^{cl}}, \quad (12.6)$$

satisfy the differential equation

$$\left(-\frac{\delta^2}{2m}\frac{\partial^2}{\partial q^2} + V(q) + \frac{\delta^2}{2m}\frac{\partial_q^2 |\psi_{cl}|}{|\psi_{cl}|}\right)\psi_{cl} = E\psi_{cl}, \quad (12.7)$$

which makes clear the non-linearity of the wave equation of CM. The classical wave-function solution of (12.7) has an obvious interpretation. First observe that

$$|\psi_{cl}|^2 = \frac{|A|^2}{p}, \quad (12.8)$$

and note that the probability of finding a particle in an interval is proportional to the time the particle stays in it. On the other hand, this time is inversely proportional to the velocity of the particle. It follows from (12.8) that $|\psi_{cl}(q)|^2 dq$ is proportional to the probability of finding the particle in the interval $[q, q + dq]$. Therefore, in this respect, we have an analogy between the quantum wave-function and the classical one. A feature of (12.2) is that it is independent of δ . As such, we can fix for δ an arbitrary value. Let us set

$$\delta = \hbar, \quad (12.9)$$

so that

$$\frac{1}{2m}\left(\frac{\partial \mathcal{S}_0^{cl}}{\partial q}\right)^2 + \mathcal{W}(q) + \frac{\hbar^2}{4m}\{\mathcal{S}_0^{cl}, q\} - \frac{\hbar^2}{4m}\{\mathcal{S}_0^{cl}, q\} = 0, \quad (12.10)$$

and

$$\left(-\frac{\hbar^2}{2m}\frac{\partial^2}{\partial q^2} + V(q) - \frac{\hbar^2}{4m}\{\mathcal{S}_0^{cl}, q\}\right)\psi_{cl} = E\psi_{cl}. \quad (12.11)$$

These classical equations are the dual versions of (9.44) and (10.2). In particular, in the same way in which (9.44) was seen as the deformation of the CSHJE by the term $Q = \hbar^2\{\mathcal{S}_0, q\}/4m$, one may consider the CSHJE (12.10) to be the deformation of the QSHJE by the term $Q^{cl} = -\hbar^2\{\mathcal{S}_0^{cl}, q\}/4m$. In other words, the (9.44)(10.2) and (12.10)(12.11) show a complementarity between CM and QM.

12.3 $\mathcal{W}$ states in Classical and Quantum Mechanics

There is another aspect related to the Möbius transformations and to the properties of the $\mathcal{W}$ states. Namely, we have seen that the EP implies that $\mathcal{W}$ transforms with an inhomogeneous term under VTs. Now, it is natural to investigate whether, despite the difference between classical and quantum $\mathcal{W}$ states, there are v -maps under which $\mathcal{W}$ transforms in the same way both in CM and QM. To answer this question one simply note that comparing (7.7) and (10.31) it follows that the only VTs under which the classical and quantum transformation properties of $\mathcal{W}$ coincide are the transformations $q^a \longrightarrow q^b$ such that the inhomogeneous term in the right hand side of (10.31) vanishes, that is

$$\{q^b, q^a\} = 0, \quad (12.12)$$

whose solution is

$$q^b = \frac{Aq^a + B}{Cq^a + D}. \quad (12.13)$$

Hence, under VTs corresponding to Möbius transformations, the $\mathcal{W}$ states transform in the same way in CM and QM. In particular, under v -maps of the kind (12.13) we have ($AD - BC = 1$)

$$\mathcal{W}^b(q^b) = (Cq^a + D)^4 \mathcal{W}^a(q^a), \quad Q^b(q^b) = (Cq^a + D)^4 Q^a(q^a). \quad (12.14)$$

It follows that

$$\mathcal{M}(q) = \frac{\mathcal{W}(q)}{Q(q)}, \quad (12.15)$$

is a Möbius invariant function. That is, under (12.12) we have

$$\mathcal{M}^b(q^b) = \mathcal{M}^a(q^a). \quad (12.16)$$

13 The trivializing map and quantum transformations

We have seen that the trivializing coordinate plays a crucial role in our formulation. In this section we will further investigate its properties in the context of classical and quantum canonical transformations. Furthermore, we will see that there is an analogy between the trivializing map and the map associated to the universal covering in the uniformization theory of Riemann surfaces.

Another topic of this section, concerns the relations between the CSHJE and QSHJE. In particular, we will show that there is a transformation reducing the QSHJE to the CSHJE. Next, we will show that both the QSHJE and the SE can be considered in the framework of projective geometry. In particular, we will use the fact that the Schwarzian derivative can be interpreted as an invariant of an equivalence problem for curves in $\mathbb{P}^1$. This fact will shed light on the geometrical nature of the trivializing map. We will conclude this section by showing that Heisenberg commutation relations actually arise from the so called area function of projective geometry.

A central point of our investigation is that the existence of the trivializing coordinate is closely related to the existence of the self-dual states. A related issue concerns the structure of the QSHJE. We have seen that this equation is equivalent to the Schwarzian equation $\{e^{\frac{2i}{\hbar}\mathcal{S}_0}, q\} = -4m\mathcal{W}/\hbar^2$. On the other hand, since $e^{\frac{2i}{\hbar}\mathcal{S}_0}$ is a Möbius transformation of the trivializing coordinate, we see that the QSHJE (9.44) can be interpreted as the equation determining q^0 , namely

$$\{q^0, q\} = -\frac{4m}{\hbar^2}(V(q) - E). \quad (13.1)$$

In this context the SE can be seen as the linearization of the problem of finding the trivializing map. Since under $q \longrightarrow q^0$ we have $\mathcal{W} \longrightarrow \mathcal{W}^0$, there is the following correspondence between the derivation of the classical and quantum HJ equations

As the existence of the classical trivializing conjugate variables (Q, P) , defined by the canonical transformation

$$q \longrightarrow Q, \quad p \longrightarrow P = \text{cnst} = -\partial_Q \mathcal{S}_0^{\text{cl}}(q, Q)|_{Q=\text{cnst}}, \quad (13.2)$$

implies the CSHJE $H(q, p = \partial_q \mathcal{S}_0^{cl}) - E = \tilde{H}(Q, P) = 0$, also Eq.(13.1) is a consequence of the existence of the trivializing map

$$q \longrightarrow q^0 = \gamma_q^{-1} e^{\frac{2i}{\hbar} \mathcal{S}_0(q)} = \frac{i\hbar}{2p_0}, \quad p \longrightarrow p_0 = (\partial_q q^0)^{-1} p = \frac{i\hbar}{2q^0}, \quad (13.3)$$

under which $\mathcal{W} \longrightarrow \mathcal{W}^0$.

Even if in (13.3) we considered non-real functions, later on we will see that the reality condition will fix a different form which is related to (13.3) by a Möbius transformation. We now observe that there is a property of the approach which allows to make the correspondence between the classical and quantum cases rather stringent. The point is that due to the fact that $\mathcal{S}_0^v(q^v) = \mathcal{S}_0(q^0)$, the relation $p_0 = \partial_{q^0} \mathcal{S}_0^0(q^0)$ can be equivalently written in the form

$$p_0 = \frac{\partial \mathcal{S}_0(q)}{\partial q^0}. \quad (13.4)$$

As a consequence there is the classical-quantum correspondence

$$p = \frac{\partial \mathcal{S}_0^{cl}}{\partial q} \quad \Longleftrightarrow \quad p = \frac{\partial \mathcal{S}_0}{\partial q}, \quad (13.5)$$

$$P = cnst = -\frac{\partial \mathcal{S}_0^{cl}}{\partial Q} \Big|_{Q=cnst} \quad \Longleftrightarrow \quad p_0 = \frac{\partial \mathcal{S}_0}{\partial q^0}. \quad (13.6)$$

Let us show the effect of this transformation on the SE. First of all note that as $\mathcal{S}_0^v(q^v) = \mathcal{S}_0(q)$, it follows by (10.4) that if $\psi^v(q^v)$ solves the SE with $\mathcal{W}^v(q^v) = V^v(q^v) - E^v$, then $\psi(q)$ defined by

$$\psi^v(q^v)(dq^v)^{-1/2} = \psi(q)(dq)^{-1/2}, \quad (13.7)$$

is solution of the SE with $\mathcal{W}(q) = V(q) - E$. In other words, under v -maps the solutions of the SE transform as $-1/2$ -differentials. In particular

$$\psi^0(q^0) = (\partial_q q^0)^{1/2} \psi(q). \quad (13.8)$$

The SE for $\mathcal{W}^0$ is $\partial_{q^0}^2 \psi^0(q^0) = 0$, which is equivalent to

$$q^{0'3/2} \frac{\partial^2 \psi^0(q^0)}{\partial q^{02}} = 0, \quad (13.9)$$

where $q^{0'} \equiv \partial_q q^0$. Therefore, it follows from Eqs.(10.34)(13.8)(13.9) that

$$q^{0'3/2} \frac{\partial^2 \psi^0(q^0)}{\partial q^{02}} = q^{0'1/2} \frac{\partial}{\partial q} q^{0'-1} \frac{\partial}{\partial q} q^{0'1/2} \psi(q) = \left(\frac{\partial^2}{\partial q^2} + \frac{1}{2} \{e^{\frac{2i}{\hbar} \mathcal{S}_0}, q\} \right) \psi(q) = 0, \quad (13.10)$$

that by (9.42) is equivalent to the SE (10.2).

13.1 On the quantum canonical transformations

It follows by (13.7) that the VTs do not preserve the transition amplitudes. This was expected as the VTs, which connect different physical systems, are quite different from the transformations one usually considers in QM. In this context it is useful to recall some facts concerning quantum canonical transformations (we refer to [48] for further interesting observations). A first important step was made by Dirac [49] and Weyl [50] who observed that unitary transformations are canonical. In [51] it was proposed to define the quantum canonical transformations as the transformations in the quantum phase space such that the commutator remains invariant

$$[q, p] = i\hbar = [Q(q, p), P(q, p)]. \quad (13.11)$$

This implies that $Q(q, p) = CqC^{-1}$, $P(q, p) = CpC^{-1}$ with C dependent on q and p . Hence, the quantum canonical transformations may be non-unitary. As observed in [48], progress on the quantum canonical transformations has been inhibited because of the mistaken belief that such transformations must be unitary. In this context we recall that we derived the QSHJE just by posing a similar question to that considered in deriving the CSHJE from a canonical transformation. Then, the VTs seem closely related to a quantum analogue of the canonical transformations. In particular, observe that in order to closely follow the analogy with the classical case, in QM one should start with q and p considered as *independent* variables and then perform a transformation to a new set of *independent* variables Q and P . Suppose that the formalism has been defined. Then, in the case in which Q and P correspond to the state $\mathcal{W}^0$, the corresponding equation would relate q and p so that they will become *dependent* variables, that is $p = \partial_q \mathcal{S}_0$ with $\mathcal{S}_0$ solution of the QSHJE (9.44) which we obtained from the EP.

13.2 Analogy with uniformization theory

As we said, the main feature distinguishing the QSHJE from the CSHJE is that while the latter is a first-order non-linear differential equation, the QSHJE is a third-order one. This implies that in the quantum case one should specify more initial conditions than in the classical case. Let us show how this reflects in finding the trivializing coordinate.

The concept of trivializing coordinate is reminiscent of that arising in uniformization theory of Riemann surfaces. We now show the analogy between our formulation and uniformization theory (see [52][53][54] for an introduction to uniformization theory, including some physical applications.). According to uniformization theory, a negatively curved Riemann surface Σ can be represented as

$$\Sigma \cong H/\Gamma, \quad (13.12)$$

$$J_{\mathbb{H}} : \mathbb{H} \longrightarrow \Sigma, \quad (13.13)$$

where $\mathbb{H} = \{w | \text{Im } w > 0\}$ is the upper half plane, Γ is a discrete subgroup of $SL(2, \mathbb{R})$ (more precisely a Fuchsian group), and $J_{\mathbb{H}}$ is the uniformizing map satisfying

$$J_{\mathbb{H}} \left(\frac{Aw + B}{Cw + D} \right) = J_{\mathbb{H}}(w), \quad (13.14)$$

$\begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \Gamma$. Let us consider the trivial equation on $\mathbb{H}$

$$\partial_w^2 \phi^0 = 0, \quad (13.15)$$

and observe that since the ratio of two linearly independent solutions of (13.15) corresponds to a Möbius transformation of w , we have

$$w = \frac{A\phi_2^0 + B\phi_1^0}{C\phi_2^0 + D\phi_1^0}. \quad (13.16)$$

Eq.(13.15) can be seen as the analogue of Eq.(13.9). Let us set

$$w = J_{\mathbb{H}}^{-1}(z), \quad (13.17)$$

where $J_{\mathbb{H}}^{-1} : \Sigma \longrightarrow \mathbb{H}$ is the inverse of the uniformizing map (13.13). Then, since $\partial_w = \frac{1}{J_{\mathbb{H}}^{-1}} \partial_z$, we have that Eq.(13.15) is equivalent to

$$\left(\frac{\partial^2}{\partial z^2} + \frac{1}{2} \{J_{\mathbb{H}}^{-1}, z\} \right) (\partial_z J_{\mathbb{H}}^{-1})^{-1/2} \phi^0 = 0. \quad (13.18)$$

Suppose that $T(z) = \frac{1}{2} \{J_{\mathbb{H}}^{-1}, z\}$, known as Liouville stress tensor or Fuchsian projective connection, is given. Then, taking the ratio of two linearly independent solutions of the *uniformizing equation*

$$\left(\frac{\partial^2}{\partial z^2} + T(z) \right) \phi(z) = 0, \quad (13.19)$$

would give the inverse of the uniformizing map. Actually, first observe that by (13.18) and (13.19)

$$\phi^0(w) = (\partial_z w)^{1/2} \phi(z), \quad (13.20)$$

which is the analogue of (13.8). Then (13.15)(13.16) and (13.20) yield

$$J_{\mathbb{H}}^{-1}(z) = \frac{A\phi_2 + B\phi_1}{C\phi_2 + D\phi_1}. \quad (13.21)$$

Therefore, the trivializing coordinate of QM is the analogue of the inverse map of uniformization

$$q^0 \sim J_{\mathbb{H}}^{-1}. \quad (13.22)$$

While the map $q \longrightarrow q^0$ corresponds to $\mathcal{W} \longrightarrow \mathcal{W}^0$, in the case of uniformization theory we have that $z \longrightarrow J_{\mathbb{H}}^{-1}$ maps a Riemann surface to a simply connected domain (its universal covering).

Going further with the analogy between uniformization theory and our approach, we note that also in uniformization theory the manifest covariance is due to the conventional choice of considering ϕ as a $-1/2$ -differential. This is exactly the same transformation property of the wave-function: a direct consequence of choosing the VTs to represent the functional change on the reduced action induced by the coordinate transformations. As in our formulation, also in the case of uniformization theory, choosing a different convention would make the formalism far more cumbersome.

There is another analogy between our formulation and uniformization theory. Namely, we saw that $e^{\frac{2i}{\hbar}\mathcal{S}_0}$ may be chosen real in the case in which $\bar{\psi} \propto \psi$, while when $\bar{\psi} \not\propto \psi$ we can first choose $e^{\frac{2i}{\hbar}\mathcal{S}_0}$ to be a phase and then transform it to a real trivializing map q^0 by considering the Cayley map. In this case we can use the fact that a phase can be transformed to the real axis by any real Möbius transformation of the Cayley map¹¹

$$q^0 = l \frac{e^{\frac{2i}{\hbar}\mathcal{S}_0} + i}{ie^{\frac{2i}{\hbar}\mathcal{S}_0} + 1} = l \frac{\cos(2\mathcal{S}_0/\hbar)}{1 - \sin(2\mathcal{S}_0/\hbar)}, \quad (13.23)$$

where l is a constant with the dimension of a length. This means that in general, given two linearly independent solutions of the SE, we have to solve the constraint on A, B, C, D

$$\text{Im } X = 0, \quad (13.24)$$

where

$$X \equiv \frac{A\psi^D + B\psi}{C\psi^D + D\psi}. \quad (13.25)$$

Once the constraint is solved we have

$$q^0 = \gamma(X), \quad (13.26)$$

where $\gamma(X) \in \hat{\mathbb{R}}$ is a $PGL(2, \mathbb{R})$ Möbius transformation of X . Note that in writing (13.21) we did not specify the possible values of A, B, C, D . On the other hand, by definition we should require

$$\text{Im } J_{\mathbb{H}}^{-1} > 0. \quad (13.27)$$

Therefore, similarly to the identification of the trivializing coordinate q^0 , also in this case, for a given pair of linearly independent solutions of the uniformizing equation, we have to find the constraint on the generally *complex* coefficients A, B, C, D , such that Eq.(13.27) is satisfied. Since $PSL(2, \mathbb{R})$ maps $\mathbb{H}$ into itself, once that A, B, C, D are determined, any other $PSL(2, \mathbb{R})$ -transformation would correspond to a possible determination of $J_{\mathbb{H}}^{-1}$. As in Eq.(13.26), which follows from the reality of the VTs and corresponds to a residual symmetry of the original Möbius one, also in the case of uniformization we have that the original complex Möbius symmetry is restricted by (13.27) to a residual $PSL(2, \mathbb{R})$ -symmetry. However, this symmetry corresponds to the fact that $PSL(2, \mathbb{R})$ maps $\mathbb{H}$ into itself. Acting with $\Gamma \subset PSL(2, \mathbb{R})$ on $J_{\mathbb{H}}^{-1}$, one obtains points in $\mathbb{H}$ with the same image in Σ .

Similarly to the case of QM, where the trivializing map reduces the system to the state $\mathcal{W}^0$, in the case of uniformization theory, the polymorphic function $J_{\mathbb{H}}^{-1}$ maps a non-trivial topology to a simply connected domain. More explicitly, similarly to the reduction $\mathcal{W} \longrightarrow \mathcal{W}^0$, also the Liouville stress tensor T in the uniformizing equation for Σ reduces to the trivial one under the $J_{\mathbb{H}}^{-1}$ map. In particular, in the context of this analogy, connecting different $\mathcal{W}$ states corresponds to connecting Riemann

¹¹The Cayley transformation maps $\Delta = \{z ||z| < 1\}$ into $\mathbb{H}$. Then, if $J_{\Delta} : \Delta \longrightarrow \Sigma$ is the uniformizing map, we have

$$J_{\mathbb{H}}^{-1} = \frac{J_{\Delta}^{-1} + i}{iJ_{\Delta}^{-1} + 1}.$$

surfaces that may be different not only in their complex structures but already at the topological level. A suitable space to describe any possible Riemann surface is the famous Bers universal Teichmüller space $T(1)$ [55]. This can be represented in the form

$$T(1) = QS(S^1)/Möb(S^1), \quad (13.28)$$

where $QS(S^1)$ is the space of *quasisymmetric* maps of the unit circle and $Möb(S^1)$ denotes the boundary transformations induced by the conformal automorphisms of the Poincaré disc. A quasisymmetric map f is an increasing self-homeomorphism of the real axis that can be extended to a quasiconformal mapping of $\mathbb{H}$ that fixes the point at infinity. According to a basic result by Beurling and Ahlfors [56] f is quasisymmetric if for some constant K , $1 \leq K < \infty$

$$\frac{1}{K} \leq \frac{f(q+t) - f(q)}{f(q) - f(q-t)} \leq K, \quad (13.29)$$

where $q \in \mathbb{R}$ and $t > 0$ (for details see *e.g.* [57]). Eq.(13.29) and its duality $K \longleftrightarrow 1/K$ indicate that these aspects are related to the theory of univalent functions and to distortion theorems. The role of this fascinating field in basic physical topics, and the related complex dynamics theory, is still to be developed. For example, the Koebe 1/4-theorem and the Schwarz lemma determine inequalities both in two-dimensional quantum gravity [54] and $N = 2$ SYM theories [58]. This would suggest that the theory of univalent functions, including related topics such as uniformization theory and Teichmüller theory, should play a role in considering basic quantum field theoretical aspects such as possible dualities between infrared and ultraviolet behavior. A related topic has been investigated by Callan and Wilczek [59]. We also observe that the extension to $\mathbb{H}$ of maps defined on $\mathbb{R}$ is reminiscent of the Holographic Principle [60]. Actually, recently ultraviolet – infrared dualities have been investigated in the framework of the Holographic Principle and AdS dualities [61] by Susskind and Witten [62]. The physical relevance of the theory of univalent functions is related to the role of the Möbius transformations in uniformization theory. The deepest one being Poincaré’s discovery that $PSL(2, \mathbb{R})$ Möbius transformations are both automorphisms of $\mathbb{H}$ and isometries of the Poincaré metric. The fact that the theory of univalent functions, and in particular $T(1)$, is of physical interest, was first realized by Bers in [63] who noticed its relation with the DeWitt superspace. It is worth noting that the universal Teichmüller space $T(1)$ also appears in string theory.

An interesting point concerning the VTs is that these correspond to real maps. We have seen that, thanks to the Möbius symmetry of the Schwarzian derivative, for any state one can define a real trivializing map. Now, suppose we are interested in finding the map

$$q^a \longrightarrow q^b = v^{ba}(q^a), \quad (13.30)$$

such that $\mathcal{W}^a(q^a)$ maps to $\mathcal{W}^b(q^b)$. To find it we can choose the pattern

$$\mathcal{W}^a \longrightarrow \mathcal{W}^0 \longrightarrow \mathcal{W}^b. \quad (13.31)$$

Denoting by $q^0 = v^x(q^x)$ the trivializing map reducing $\mathcal{W}^x(q^x)$ to $\mathcal{W}^0(q^0)$, we have

$$v^{ba} = v^{b^{-1}} \circ v^a, \quad (13.32)$$

so that, since the trivializing maps can be chosen to be real, we have that v^{ba} is a real map.

13.3 Quantum transformations

A property of the QSHJE is that it can be seen as a deformation of the CSHJE (2.15) by a “conformal factor”. In fact, noting that

$$\{\mathcal{S}_0, q\} = -(\partial_q \mathcal{S}_0)^2 \{q, \mathcal{S}_0\}, \quad (13.33)$$

we have that the QSHJE (9.44) is equivalent to

$$\frac{1}{2m} \left(\frac{\partial \mathcal{S}_0}{\partial q} \right)^2 [1 - \hbar^2 \mathcal{U}(\mathcal{S}_0)] + V(q) - E = 0, \quad (13.34)$$

where $\mathcal{U}(\mathcal{S}_0)$ is just the canonical potential (4.10) we introduced in the framework of p - q duality. Note that (13.33) also implies

$$2mQ + \hbar^2 p^2 \mathcal{U} = 0. \quad (13.35)$$

In the case of the state $\mathcal{W}^0(q^0)$, Eq.(13.34) becomes

$$\frac{1}{2m} \left(\frac{\partial \mathcal{S}_0^0}{\partial q^0} \right)^2 [1 - \hbar^2 \mathcal{U}(\mathcal{S}_0)] = 0, \quad (13.36)$$

whose solution is

$$\mathcal{U} \left(\frac{\hbar}{2i} \ln \gamma(q^0) \right) = \frac{1}{\hbar^2}, \quad (13.37)$$

where

$$\gamma(q^0) = \frac{Aq^0 + B}{Cq^0 + D}, \quad (13.38)$$

$AD - BC \neq 0$. Eq.(13.37) shows the important role of the purely quantum mechanical self-dual state (10.23). Eq.(13.34) can be written in the form

$$\frac{1}{2m} \left(\frac{\partial \mathcal{S}_0}{\partial \hat{q}} \right)^2 + V(q) - E = 0, \quad (13.39)$$

where

$$\left(\frac{\partial q}{\partial \hat{q}} \right)^2 = [1 - \hbar^2 \mathcal{U}(\mathcal{S}_0)], \quad (13.40)$$

or equivalently (we omit the solution with the minus sign)

$$d\hat{q} = \frac{dq}{\sqrt{1 - \beta^2(q)}}, \quad (13.41)$$

with $\beta^2(q) = \hbar^2 \mathcal{U}(\mathcal{S}_0) = \hbar^2 \{q, \mathcal{S}_0\}/2$. Integrating (13.40) yields

$$\hat{q} = \int^q \frac{dx}{\sqrt{1 - \beta^2(x)}}. \quad (13.42)$$

Observe that

$$\lim_{\hbar \rightarrow 0} \hat{q} = q. \quad (13.43)$$

Eq.(13.42) indicates that in considering the differential structure one should take into account the effect of the potential on space geometry. In this context the deformation of the classical HJ equation amounts to replacing the standard derivative with respect to the classical coordinate q with the derivative with respect to the quantum coordinate $\hat{q}$. This allows to put the QSHJE in the classical form. Namely, setting $\hat{\mathcal{W}}(\hat{q}) = \mathcal{W}(q(\hat{q}))$ and $\hat{\mathcal{S}}_0(\hat{q}) = \mathcal{S}_0(q(\hat{q}))$, we have that Eq.(13.34) is equivalent to

$$\frac{1}{2m} \left(\frac{\partial \hat{\mathcal{S}}_0(\hat{q})}{\partial \hat{q}} \right)^2 + \hat{\mathcal{W}}(\hat{q}) = 0. \quad (13.44)$$

The fact that the QSHJE admits the classical representation (13.44) suggests that classically forbidden regions correspond to critical regions for the quantum coordinate. Actually, writing Eq.(13.42) in the equivalent form ($\mathbf{s} = \mathcal{S}_0(q)$)

$$\hat{q} = \int^q dx \frac{\partial_x \mathcal{S}_0}{\sqrt{-2m\mathcal{W}}} = \int^{\mathcal{S}_0(q)} \frac{ds}{\sqrt{-2m\mathcal{W}}}, \quad (13.45)$$

we see that the integrand is purely imaginary in the classically forbidden regions $\mathcal{W} > 0$. Furthermore, since according to (13.36), for the state $\mathcal{W}^0$ the conformal factor vanishes, it follows by (13.42) that the quantum coordinate for the free particle state with vanishing energy is divergent.

13.4 Classical and quantum potentials and equivalence of curves

We now consider a result obtained by Flanders [64] who showed that the Schwarzian derivative can be interpreted as an invariant (curvature) of an equivalence problem for curves in $\mathbb{P}^1$. Let us introduce a frame for $\mathbb{P}^1$, that is a pair $\mathbf{x}, \mathbf{y}$ of points in affine space $\mathbf{A}^2$ such that $[\mathbf{x}, \mathbf{y}] = 1$, where

$$[\mathbf{x}, \mathbf{y}] = \mathbf{x}^t \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \mathbf{y} = x_1 y_2 - x_2 y_1, \quad (13.46)$$

is the area function. Observe that the area function has the $SL(2, \mathbb{R})$ -symmetry

$$[\tilde{\mathbf{x}}, \tilde{\mathbf{y}}] = [\mathbf{x}, \mathbf{y}], \quad (13.47)$$

where $\tilde{\mathbf{x}} = R\mathbf{x}$ and $\tilde{\mathbf{y}} = R\mathbf{y}$ with $R \in SL(2, \mathbb{R})$. Considering the moving frame $s \longrightarrow \{\mathbf{x}(s), \mathbf{y}(s)\}$ and differentiating $[\mathbf{x}, \mathbf{y}] = 1$ yields the structure equations

$$\mathbf{x}' = a\mathbf{x} + b\mathbf{y}, \quad \mathbf{y}' = c\mathbf{x} - a\mathbf{y}, \quad (13.48)$$

where a, b , and c depend on s . Given a map $\phi = \phi(s)$ from a domain to $\mathbb{P}^1$, one can choose a moving frame $\mathbf{x}(s), \mathbf{y}(s)$ in such way that $\phi(s)$ is represented by $\mathbf{x}(s)$. Observe that this map can be seen as a curve in $\mathbb{P}^1$. Two mappings ϕ and ψ are said to be equivalent if $\psi = \pi \circ \phi$, with π a projective transformation on $\mathbb{P}^1$.

Flanders considered two extreme situations. The first case corresponds to $b(s) = 0, \forall s$. In this case ϕ is constant. To see this observe that taking the derivative of $\lambda\mathbf{x}$, for some $\lambda(s) \neq 0$, we have by

(13.48) that $(\lambda \mathbf{x})' = (\lambda' + a\lambda)\mathbf{x}$. Choosing $\lambda \propto \exp[-\int_{s_0}^s dt a(t)] \neq 0$, we have $(\lambda \mathbf{x})' = 0$, so that $\lambda \mathbf{x}$ is a constant representative of ϕ .

The other case is for b never vanishing. There are only two inequivalent situations. The first one is when b is either complex or positive. It turns out that it is always possible to choose the following “natural moving frame” for ϕ [64]

$$\mathbf{x}' = \mathbf{y}, \quad \mathbf{y}' = -k\mathbf{x}. \quad (13.49)$$

In the other case, corresponding to b real and negative, the natural moving frame for ϕ is

$$\mathbf{x}' = -\mathbf{y}, \quad \mathbf{y}' = k\mathbf{x}. \quad (13.50)$$

A characterizing property of the natural moving frame is that it is determined up to a sign with k an invariant. Thus, for example, suppose that for a given ϕ there is, besides (13.49), the natural moving frame $\mathbf{x}'_1 = \mathbf{y}_1$, $\mathbf{y}'_1 = -k_1\mathbf{x}_1$. Since both $\mathbf{x}$ and $\mathbf{x}_1$ are representatives of ϕ , we have $\mathbf{x} = \lambda\mathbf{x}_1$, so that $\mathbf{y} = \mathbf{x}' = \lambda'\mathbf{x}_1 + \lambda\mathbf{y}_1$ and $1 = [\mathbf{x}, \mathbf{y}] = \lambda^2$. Therefore, $\mathbf{x}_1 = \pm\mathbf{x}$, $\mathbf{y}_1 = \pm\mathbf{y}$ and $k_1 = k$ [64].

Let us now review the derivation of Flanders formula for k . Consider $s \longrightarrow \mathbf{z}(s)$ to be an affine representative of ϕ and let $\mathbf{x}(s), \mathbf{y}(s)$ be a natural frame. Then $\mathbf{z} = \lambda\mathbf{x}$ where $\lambda(s)$ is never vanishing. Now note that, since $\mathbf{z}' = \lambda'\mathbf{x} + \lambda\mathbf{y}$, we have that λ can be written in terms of the area function $[\mathbf{z}, \mathbf{z}'] = \lambda^2$. Computing the relevant area functions, one can check that k has the following expression

$$2k = \frac{[\mathbf{z}, \mathbf{z}'''] + 3[\mathbf{z}', \mathbf{z}'']}{[\mathbf{z}, \mathbf{z}']} - \frac{3}{2} \left(\frac{[\mathbf{z}, \mathbf{z}']}{[\mathbf{z}, \mathbf{z}']} \right)^2. \quad (13.51)$$

Given a function $z(s)$, this can be seen as the non-homogeneous coordinate of a point in $\mathbb{P}^1$. Therefore, we can associate to z the map ϕ defined by $s \longrightarrow (1, z(s)) = \mathbf{z}(s)$. In this case we have $[\mathbf{z}, \mathbf{z}'] = z'$, $[\mathbf{z}, \mathbf{z}'] = z''$, $[\mathbf{z}, \mathbf{z}'''] = z'''$, $[\mathbf{z}', \mathbf{z}''] = 0$, and the curvature becomes [64]

$$k = \frac{1}{2}\{z, s\}. \quad (13.52)$$

Let us now consider an arbitrary state $\mathcal{W}$. We have

$$\mathcal{W} = -\frac{\hbar^2}{4m}\{e^{\frac{2i}{\hbar}\mathcal{S}_0}, q\} = -\frac{\hbar^2}{2m}k_{\mathcal{W}}. \quad (13.53)$$

Similarly, for the quantum potential

$$Q = \frac{\hbar^2}{4m}\{\mathcal{S}_0, q\} = \frac{\hbar^2}{2m}k_Q, \quad (13.54)$$

where $k_{\mathcal{W}}$ is the curvature associated to the map

$$q \longrightarrow (1, e^{\frac{2i}{\hbar}\mathcal{S}_0(q)}), \quad (13.55)$$

while the curvature k_Q is associated to the map

$$q \longrightarrow (1, \mathcal{S}_0(q)). \quad (13.56)$$

Remarkably, the function defining the map (13.55) coincides with the trivializing map. Observe that the SE takes the geometrical form

$$\left(\frac{\partial^2}{\partial q^2} + k_{\mathcal{W}}\right)\psi = 0. \quad (13.57)$$

Furthermore, the identity (9.7) can be now seen as difference of curvatures

$$(\partial_q \mathcal{S}_0)^2 = \hbar^2 k_{\mathcal{W}} - \hbar^2 k_Q, \quad (13.58)$$

and the QSHJE (9.44) can be written in the form

$$\frac{1}{2m} \left(\frac{\partial \mathcal{S}_0(q)}{\partial q}\right)^2 + \mathcal{W}(q) + \frac{\hbar^2}{2m} k_Q = 0. \quad (13.59)$$

Let us now consider the meaning of the natural moving frame in the framework of the QSHJE. First observe that the structure equations imply that

$$\mathbf{x}'' = -k\mathbf{x}. \quad (13.60)$$

In the case of $k = k_{\mathcal{W}}$, this equation is the SE, so that

$$\mathbf{x} = (\psi^D, \psi), \quad \mathbf{y} = (\psi^{D'}, \psi'), \quad (13.61)$$

and the frame condition is the statement that the Wronskian W of the SE is a constant

$$[\mathbf{x}, \mathbf{y}] = \psi' \psi^D - \psi^{D'} \psi = W = 1. \quad (13.62)$$

Hence, the SE determines the natural moving frame associated to the curve in $\mathbb{P}^1$ given by the representative (13.55) with $-2m\mathcal{W}/\hbar^2$ denoting the invariant associated to the map. In other words, the Schrödinger problem corresponds to finding the natural moving frame such that $-2m\mathcal{W}/\hbar^2$ be the invariant curvature. In the $k = k_Q$ case Eq.(13.60) becomes

$$\left(\frac{\hbar^2}{2m} \frac{\partial^2}{\partial q^2} + Q\right)\phi = 0, \quad (13.63)$$

so that if ϕ^D and ϕ are solutions of (13.63), then

$$\mathcal{S}_0 = \frac{A\phi^D + B\phi}{C\phi^D + D\phi}. \quad (13.64)$$

13.5 Area function and commutator

The area function has a structure which is reminiscent of the commutator. This feature is emphasized in Eq.(13.62) that contains the derivative operator. In fact, Eq.(13.62) can be written using the commutator between ∂_q and the ratio of solutions ψ^D/ψ . We have

$$[\mathbf{x}, \mathbf{y}] = [\partial_q, \psi^D/\psi]_c \psi^2 = 1, \quad (13.65)$$

where $[A, B]_c \equiv AB - BA$. Introducing the momentum operator $p = -i\hbar\partial_q$, (13.65) becomes

$$[\psi^D/\psi, p]_c \psi^2 = i\hbar. \quad (13.66)$$

Note that the $SL(2, \mathbb{R})$ -symmetry (13.47) implies

$$\left[\frac{A\psi^D/\psi + B}{C\psi^D/\psi + D}, p \right]_c (C\psi^D + D\psi)^2 = i\hbar. \quad (13.67)$$

Since, as we will see, the trivializing map is a Möbius transformation of the ψ^D/ψ , we have that the area function is equivalent to

$$[q^0, p]_c (C\psi^D + D\psi)^2 = i\hbar. \quad (13.68)$$

This reproduces the Heisenberg commutation relation

$$[q^0, p_0]_c = i\hbar, \quad (13.69)$$

$p_0 = -i\hbar\partial_{q^0}$, with (13.68) representing the Jacobian

$$\partial_q q^0 = (C\psi^D + D\psi)^{-2}, \quad (13.70)$$

of the trivializing transformation.

14 Canonical variables and Möbius transformations

We now discuss the role of Möbius transformations in considering the canonical variables p and q . Once again, we will use the fact that, according to (10.25), $\mathcal{W}$ remains invariant under a Möbius transformation of $e^{\frac{2i}{\hbar}\mathcal{S}_0}$. While under this transformation both $(\partial_q \mathcal{S}_0)^2$ and Q get transformed, we have the invariance

$$\left(\frac{\partial \tilde{\mathcal{S}}_0(q)}{\partial q} \right)^2 + \frac{\hbar^2}{2} \{ \tilde{\mathcal{S}}_0, q \} = \left(\frac{\partial \mathcal{S}_0(q)}{\partial q} \right)^2 + \frac{\hbar^2}{2} \{ \mathcal{S}_0, q \}, \quad (14.1)$$

where $\tilde{\mathcal{S}}_0$ is given in (10.26). We will begin this section by considering the explicit dependence of the conjugate momentum on the coordinate in the case of the state $\mathcal{W}^0$. Next, we will make some preliminary observation on the possible definition of time parameterization. Then the general expression for p in terms of q will be derived for an arbitrary state $\mathcal{W}$. We will conclude this section by a detailed investigation of the symmetries of the wave-function. By this, we mean the transformations on the initial conditions for the QSHJE which leave invariant a given eigenfunction of the Hamiltonian.

14.1 The canonical variables of the state $\mathcal{W}^0$

Let us consider the conjugate momentum in the case of the state $\mathcal{W}^0$. By (10.24) we have

$$p_0 = \frac{i(BC - AD)\hbar}{2(Aq^0 + B)(Cq^0 + D)}. \quad (14.2)$$

On the other hand, since $AD - BC \neq 0$, by a common rescaling of A, B, C, D , that by projectivity of the Möbius transformations do not change (10.24), we can set

$$AD - BC = \pm i, \quad (14.3)$$

so that

$$p_0 = \pm \frac{\hbar}{2[ACq^{02} + (AD + BC)q^0 + BD]}. \quad (14.4)$$

Observe that if $AC = 0$, then either BC or AD vanish and by (14.3) $p_0 = \pm\hbar/2(\pm iq^0 + BD)$. Similarly, if $BD = 0$, then $p_0 = \pm\hbar/2(ACq^{02} \pm iq^0)$. It follows that

p_0 can take real values only if $AC \neq 0$ and $BD \neq 0$, that is $p_0 = \pm\hbar(\alpha q^{02} + \beta q^0 + \gamma)^{-1}$, $4\alpha\gamma - \beta^2 = 4$, $\beta \in \mathbb{R}$ with both α and γ taking values in $\mathbb{R} \setminus \{0\}$.

By (14.3) it follows that the poles of p_0 are never on the real q^0 -axis. Actually, the poles of (14.4) are

$$q^0 = -\frac{BC + AD \pm i}{2AC}. \quad (14.5)$$

Since reality of (14.4) implies $(BC + AD) \in \mathbb{R}$ and $AC \in \mathbb{R} \setminus \{0\}$, it follows that the solutions (14.5) cannot be real. By (14.3) we can rewrite Eq.(14.4) in the form

$$p_0 = \pm \frac{\hbar \ell_1}{(q^0 + \ell_2)^2 + \ell_1^2}, \quad (14.6)$$

where

$$\ell_1 = \frac{1}{2AC} \neq 0, \quad \ell_2 = \frac{BC + AD}{2AC}. \quad (14.7)$$

Observe that both ℓ_1 and ℓ_2 are real constants. A property of p_0 is that it vanishes only at infinity

$$\lim_{q^0 \rightarrow \pm\infty} p_0 = 0. \quad (14.8)$$

Furthermore, for $q^0 = -\ell_2$, $|p_0|$ reaches its maximum

$$|p_0(-\ell_2)| = \frac{\hbar}{\ell_1}. \quad (14.9)$$

Since $\ell_1 \neq 0$, we have

$$|p_0| < \infty. \quad (14.10)$$

14.2 Time and elliptic curve as moduli of the state $\mathcal{W}^0$

Let us consider time parameterization. In CM, this is derived from the CSHJE by identifying the conjugate momentum with $m\dot{q}^0$. Since $p_0 = \partial_{q^0}\mathcal{S}_0^0 = m\dot{q}^0$, this identification would imply a non-vanishing velocity associated to the state $\mathcal{W}^0$. Later on, we will see that a natural derivation of time parameterization, suggested by Floyd [15], implies that, generally, one has $\partial_q\mathcal{S}_0 \neq m\dot{q}$. We will see that with Floyd's time parameterization one has $\dot{q}^0 = 0$.

Let us then investigate the structure of $q^0(\tau)$ with τ defined by $\partial_q \mathcal{S}_0 = m \partial_\tau q$, that is

$$\tau - \tau_0 = m \int_{q_0}^q dx \frac{1}{\partial_x \mathcal{S}_0(x)}, \quad (14.11)$$

so that

$$p_0 = m \frac{dq^0}{d\tau}. \quad (14.12)$$

Eq.(14.6) becomes $m\dot{q}^0[(q^0 + \ell_2)^2 + \ell_1^2] \pm \hbar \ell_1 = 0$, which can be integrated to

$$\frac{m}{3} q^{03} + m \ell_2 q^{02} + m(\ell_1^2 + \ell_2^2) q^0 + c \pm \hbar \ell_1 \tau = 0, \quad (14.13)$$

where c is an integration constant. Setting $y^2 = \pm 3\hbar \ell_1 \tau / m$, and $\alpha_0 = 3\ell_2$, $\alpha_1 = 3(\ell_1^2 + \ell_2^2)$, $\alpha_2 = 3c/m$, we have that Eq.(14.13) becomes the equation of an elliptic curve

$$y^2 = q^{03} + \alpha_0 q^{02} + \alpha_1 q^0 + \alpha_2. \quad (14.14)$$

As we noted, this time parameterization, made by identifying the conjugate momentum with the mechanical one, gives $\dot{q}^0 \neq 0$. We will then consider Floyd's time parameterization which will imply the expected result $\dot{q}^0 = 0$. However, let us note that it is intriguing that the above identification of time, borrowed from CM, leads to an underlying non-trivial geometrical structure. This suggests that time defined through $\partial_q \mathcal{S}_0 = m\dot{q}$ plays some non-trivial role. Even if we will not consider further this “ τ -parameterization” here, its analogy with the classical case deserves further exploration.

14.3 The canonical variables of arbitrary states

The above investigation indicates that there is a structure which generalizes to arbitrary $\mathcal{W}$ states. In order to determine p as function of q , we use the fact that the SE can be seen as the linearization of the Schwarzian equation

$$\{e^{\frac{2i}{\hbar} \mathcal{S}_0}, q\} = -\frac{4m}{\hbar^2} \mathcal{W}(q). \quad (14.15)$$

We have seen that this implies that

$$e^{\frac{2i}{\hbar} \mathcal{S}_0} = \frac{A\psi^D + B\psi}{C\psi^D + D\psi}, \quad (14.16)$$

so that

$$p = \frac{i}{2} \frac{\hbar(AD - BC)W}{AC\psi^{D^2} + (AD + BC)\psi^D\psi + BD\psi^2}, \quad (14.17)$$

where W denotes the Wronskian

$$W = \psi' \psi^D - \psi^{D'} \psi, \quad (14.18)$$

which is a constant. Note that p can be written in the form

$$p = \frac{i}{2} \frac{\hbar \tilde{W}}{\tilde{\psi}^D \tilde{\psi}}, \quad (14.19)$$

where

$$\begin{pmatrix} \tilde{\psi}^D \\ \tilde{\psi} \end{pmatrix} = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \begin{pmatrix} \psi^D \\ \psi \end{pmatrix}, \quad (14.20)$$

and $\tilde{W} = \tilde{\psi}'\tilde{\psi}^D - \tilde{\psi}^{D'}\tilde{\psi} = (AD - BC)W$. Setting

$$\alpha = \frac{2iAC}{(BC - AD)W}, \quad \beta = \frac{2i(AD + BC)}{(BC - AD)W}, \quad \gamma = \frac{2iBD}{(BC - AD)W}, \quad (14.21)$$

Eq.(14.17) becomes

$$p = \frac{\hbar}{\alpha\psi^{D^2} + \beta\psi^D\psi + \gamma\psi^2}. \quad (14.22)$$

Since $(AD + BC)^2 - 4(AC)(BD) = (AD - BC)^2$, it follows by (14.21) that

$$\beta^2 - 4\alpha\gamma = -\frac{4}{W^2}. \quad (14.23)$$

Note that since ψ^D and ψ are two arbitrary linearly independent solutions of the SE, we can always choose them to take real values

$$(\psi^D, \psi) \in \mathbb{R}^2. \quad (14.24)$$

Actually, since W is real, it follows that if ϕ is a solution of the SE, then the real function $\psi = \phi + \bar{\phi}$ is still a solution. Any other solution has the form

$$\psi^D(q) = c\psi(q) \int_{q_0}^q dx \psi^{-2}(x) + d\psi(q), \quad (14.25)$$

for some constants c and d . Observe that $W = -c$. Note that a rescaling of ψ and a change of c and d in (14.25) is equivalent to a transformation of the coefficients A, B, C, D in (14.16). By (14.22) it follows that reality of (ψ^D, ψ) and (p, q) implies $W \in \mathbb{R} \setminus \{0\}$, and $\alpha, \beta, \gamma \in \mathbb{R}$, which in turn, by (14.23), imply $\alpha \neq 0, \gamma \neq 0$. Since $\alpha\psi^{D^2} + \beta\psi^D\psi + \gamma\psi^2$ vanishes for

$$\psi^D(q) = -\frac{\psi(q)}{2\alpha} \left(\beta \pm \frac{2i}{W} \right), \quad (14.26)$$

and being $W \in \mathbb{R} \setminus \{0\}$, it follows that p has no poles for real q . The possibility that ψ^D and ψ vanish at the same point is easily ruled out. For, if $\psi^D(q_0) = 0 = \psi(q_0)$ then $W = \psi'(q_0)\psi^D(q_0) - \psi^{D'}(q_0)\psi(q_0) = 0$, which is not the case as ψ^D and ψ are linearly independent so that $W \neq 0$.

Let us consider again Eq.(14.22). Note that writing it in the form

$$p = \frac{\hbar\alpha^{-1}}{\left(\psi^D + \frac{\beta}{2\alpha}\psi\right)^2 + \frac{1}{\alpha^2 W^2}\psi^2}, \quad (14.27)$$

and setting

$$\ell_1 = \frac{1}{\alpha W} = i\frac{AD - BC}{2AC}, \quad \ell_2 = \frac{\beta}{2\alpha} = \frac{AD + BC}{2AC}, \quad (14.28)$$

we have

$$p = \pm \frac{\hbar W \ell_1}{(\psi^D + \ell_2 \psi)^2 + \ell_1^2 \psi^2}. \quad (14.29)$$

Since $AD - BC \neq 0$ we have¹²

$$\ell_1 \in \mathbb{R} \setminus \{0\}. \quad (14.30)$$

From the general expression (14.29) we can directly derive the solution Eq.(14.6) for the state $\mathcal{W}^0$. Namely, since two linearly independent solutions of the SE associated with the state $\mathcal{W}^0$ are

$$\psi^{D^0} = q^0, \quad \psi^0 = 1, \quad (14.31)$$

we have $W = -1$ and Eq.(14.29) reproduces Eq.(14.6). Note that Eq.(14.29) is invariant under a common rescaling of ψ^D and ψ . More generally, any $GL(2, \mathbb{R})$ -transformation of the vector

$$v = \begin{pmatrix} \psi^D \\ \psi \end{pmatrix}, \quad (14.32)$$

which in turn may imply also a rescaling of the Wronskian W , is always equivalent to a transformation of the ℓ_k 's. This can be seen also by expressing p in a different form. Let us introduce the matrix

$$\gamma = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \in GL(2, \mathbb{R}). \quad (14.33)$$

Setting

$$a_{11} = \frac{c}{2\ell_1 \hbar W}, \quad a_{12} = \frac{c\ell_2}{2\ell_1 \hbar W}, \quad a_{21} = \frac{1}{c}, \quad a_{22} = \frac{\ell_1^2 + \ell_2^2}{c\ell_2}, \quad (14.34)$$

where c is an arbitrary non-vanishing constant, we have

$$p^{-1} = \pm v^t \left[\gamma^t \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \gamma - \begin{pmatrix} 0 & \sigma \\ \sigma & 0 \end{pmatrix} \right] v, \quad (14.35)$$

where $\sigma \equiv \det \gamma$, that by (14.34) is

$$\sigma = \frac{\ell_1}{2\ell_2 \hbar W}. \quad (14.36)$$

Observe that

$$\gamma_p \equiv \gamma^t \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \gamma - \begin{pmatrix} 0 & \sigma \\ \sigma & 0 \end{pmatrix} = 2 \begin{pmatrix} a_{11}a_{21} & a_{12}a_{21} \\ a_{12}a_{21} & a_{12}a_{22} \end{pmatrix}, \quad (14.37)$$

so that with the identification (14.34), we have

$$\gamma_p = \frac{1}{\ell_1 \hbar W} \begin{pmatrix} 1 & \ell_2 \\ \ell_2 & \ell_1^2 + \ell_2^2 \end{pmatrix}. \quad (14.38)$$

¹²The ± 1 factor is due to the invariance of the QSHJE (9.44) under a change of sign of $\mathcal{S}_0$. This reflects the fact that the motion can be in two directions. Observe that one can fix ℓ_1 to be either positive or negative. Alternatively, one can consider $\ell_1 \in \mathbb{R} \setminus \{0\}$ and then dropping the ± 1 factor in (14.29).

Note that $\det \gamma_p = 1/\hbar^2 W^2$ does not depend on the ℓ_k 's. Let us write down p in terms of γ_p

$$p^{-1} = \pm v^t \gamma_p v. \quad (14.39)$$

Since under a $GL(2, \mathbb{R})$ -transformation of v

$$v \longrightarrow v_R = Rv, \quad (14.40)$$

$$R = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in GL(2, \mathbb{R}), \quad (14.41)$$

the Wronskian gets rescaled by $\rho \equiv \det R$, it follows that under (14.40) p transforms as

$$p^{-1} \longrightarrow p_R^{-1} = \pm v^t R^t \gamma_{p\rho} R v, \quad (14.42)$$

where $\gamma_{p\rho} = \rho^{-1} \gamma_p$. In particular, as we already noticed, since $\det \gamma_{p\rho} = (\det R)^{-2} \det \gamma_p$, it follows that p is invariant under dilatations of the vector v . Observe that the effect on p of any linear transformation of v is equivalent to the following transformation of γ_p

$$\gamma_p \longrightarrow \gamma'_p = R^t \gamma_{p\rho} R. \quad (14.43)$$

Since $\gamma_p^t = \gamma_p$ and $\det \gamma'_p = \det \gamma$, we have

$$\gamma'_p = \frac{1}{\ell'_1 \hbar W} \begin{pmatrix} 1 & \ell'_2 \\ \ell'_2 & \ell_1'^2 + \ell_2'^2 \end{pmatrix}, \quad (14.44)$$

where

$$\ell'_1 = \rho \frac{\ell_1}{(a + \ell_2 c)^2 + \ell_1^2 c^2}, \quad \ell'_2 = \frac{ab + (bc + ad)\ell_2 + (\ell_1^2 + \ell_2^2)cd}{(a + \ell_2 c)^2 + \ell_1^2 c^2}. \quad (14.45)$$

Any linear transformation of v is equivalent to keeping v fixed and making a transformation of ℓ_1 and ℓ_2 .¹³ In particular, since W can be chosen to be an arbitrary fixed value, we have that the degrees of freedom of p are described by the two real parameters ℓ_1 and ℓ_2 . Note that setting $W = \pm 1/\hbar$, yields

$$\gamma_p(\ell_1, \ell_2) = \begin{pmatrix} \ell_1^{-1} & \ell_2 \ell_1^{-1} \\ \ell_2 \ell_1^{-1} & \ell_1 + \ell_2^2 \ell_1^{-1} \end{pmatrix}, \quad (14.46)$$

so that $\gamma_p \in SL(2, \mathbb{R})$, and

$$|\text{tr } \gamma_p| = |\ell_1^{-1} + \ell_1 + \ell_2^2 \ell_1^{-1}| \geq 2. \quad (14.47)$$

¹³The transformed quantity ℓ_{kR} have some interesting properties. For example, there is the following “formal coincidence”. Namely, relaxing the condition for the elements of the R -matrix to be constants and setting

$$R = \begin{pmatrix} \psi^D & \hbar \psi^{D'} \\ \psi & \hbar \psi' \end{pmatrix},$$

it follows by (14.45) that $\ell_1 \longrightarrow \ell_{1R} = p$.

Since $|\text{tr } \gamma_p| = 2$ only in the case in which $\ell_1 = 1, \ell_2 = 0$, corresponding to the identity matrix, it follows that γ_p is a hyperbolic matrix.

An interesting feature of the above investigation is that we started with the general expression (14.16) in which $e^{\frac{2i}{\pi} S_0}$ is given in terms of Möbius transformations of the ratio

$$w = \frac{\psi^D}{\psi} \in \mathbb{R}, \quad (14.48)$$

and then arrived at hyperbolic matrices. Let us consider the transformation

$$\gamma_p(w) = \frac{\gamma_{p11}w + \gamma_{p12}}{\gamma_{p21}w + \gamma_{p22}} = \frac{w + \ell_2}{\ell_2 w + (\ell_1^2 + \ell_2^2)}, \quad (14.49)$$

and its associated fixed point equation

$$\gamma_p(w) = w, \quad (14.50)$$

in the case in which $\gamma_p \neq \mathbb{I}_{2 \times 2}$. We have

$$w_{\pm} = \frac{1 - \ell_1^2 - \ell_2^2 \pm \sqrt{(\ell_1^2 + \ell_2^2 - 1)^2 + 4\ell_2^2}}{2\ell_2}. \quad (14.51)$$

Since, according to (14.47), we have $(\ell_1^2 + \ell_2^2 - 1)^2 + 4\ell_2^2 > 0$, it follows that $\gamma_p \neq \mathbb{I}_{2 \times 2}$ has distinct fixed points lying on the real axis. This structure indicates that the ratio $w = \psi^D/\psi \in \mathbb{R}$ can be seen as belonging to the boundary of $\mathbb{H}$. Let us denote by Γ_H the set of all matrices of the kind (14.46). A property of Γ_H is that $\gamma \in \Gamma_H \longrightarrow \gamma^{-1} \in \Gamma_H$, in particular

$$\gamma_p^{-1}(\ell_1, \ell_2) = \gamma_p\left(\frac{\ell_1}{\ell_1^2 + \ell_2^2}, -\frac{\ell_2}{\ell_1^2 + \ell_2^2}\right). \quad (14.52)$$

Suppose that there are elements γ_k in Γ_H such that

$$\prod_{k=1}^h (\gamma_{2k-1} \gamma_{2k} \gamma_{2k-1}^{-1} \gamma_{2k}^{-1}) = \mathbb{I}_{2 \times 2}. \quad (14.53)$$

Doing this is equivalent to selecting a set of $2h$ pairs (ℓ_1, ℓ_2) . Then, denoting by $\Gamma_{p\{\ell_k\}}$ the group generated by the γ_k 's, we have that with the $2h$ values of p determined by the ℓ_k 's moduli, there is associated the genus h Riemann surface

$$\Sigma_{p\{\ell_k\}} \cong H/\Gamma_{p\{\ell_k\}}. \quad (14.54)$$

Therefore, we are posing the question of determining the possible sets $\{\ell_k\}$ such that $\Sigma_{p\{\ell_k\}}$ is a Riemann surface. Since different ℓ_k 's correspond to different p 's, studying these surfaces would shed light on the geometry of the possible paths. It would be interesting to understand whether a Riemann surface of infinite genus may appear.

The generators γ_k 's of a given $\Gamma_{p\{\ell_k\}}$ would represent the handles of the Riemann surface. These elements can be represented in the form

$$\frac{\gamma_k(w) - w_+}{\gamma_k(w) - w_-} = e^{\lambda} \frac{w - w_+}{w - w_-}, \quad (14.55)$$

where

$$e^\lambda = \frac{1 + \ell_1^2 + \ell_2^2 - \sqrt{(\ell_1^2 + \ell_2^2 - 1)^2 + 4\ell_2^2}}{1 + \ell_1^2 + \ell_2^2 + \sqrt{(\ell_1^2 + \ell_2^2 - 1)^2 + 4\ell_2^2}}. \quad (14.56)$$

This means that any hyperbolic element is conjugate to a dilatation. The scale factor λ has the following geometrical meaning. Consider a point $w \in H$ and its hyperbolic transformed $\gamma_k(w)$. Then $\lambda(\gamma_k)$ corresponds to the minimal hyperbolic distance between w and $\gamma_k(w)$. This minimum is reached for w lying in the geodesic intersecting the real axis at w_- and w_+ .

14.4 The symmetries of the wave-function

The fact that the QSHJE (9.44) is a third-order differential equation implies that there are three integration constants which specify $\mathcal{S}_0$. In particular, we have seen that two real constants specify p . Concerning $\mathcal{S}_0$ there is one more constant which arises by integrating (14.29) ($w = \psi^D/\psi \in \mathbb{R}$)

$$e^{\frac{2i}{\hbar}\mathcal{S}_0\{\delta\}} = e^{i\alpha} \frac{w + i\bar{\ell}}{w - i\ell}, \quad (14.57)$$

where $\delta = \{\alpha, \ell\}$, with α a real integration constant and $\ell = \ell_1 + i\ell_2$. Observe that the condition $\ell_1 \neq 0$ is equivalent to having $\mathcal{S}_0 \neq \text{const}$ which is a necessary condition to define the term $\{\mathcal{S}_0, q\}$ in the QSHJE. We also note that changing sign of ℓ_1 corresponds to changing the sign of $\mathcal{S}_0$; more precisely we have

$$\mathcal{S}_0\{\alpha, -\bar{\ell}\} = -\mathcal{S}_0\{\alpha, \ell\} + \hbar\alpha = -\mathcal{S}_0\{-\alpha, \ell\}. \quad (14.58)$$

Note that for the conjugate momentum we have

$$p = \frac{\hbar W(\ell + \bar{\ell})}{2|\psi^D - i\ell\psi|^2}. \quad (14.59)$$

Let ψ_E be the wave-function of a state of energy E . Since ψ_E solves the SE, according to (10.4), for any fixed set of integration constants δ , there are coefficients A and B such that

$$\psi_E\{\delta\} = \frac{1}{\sqrt{\mathcal{S}_0\{\delta\}}} \left(A e^{-\frac{i}{\hbar}\mathcal{S}_0\{\delta\}} + B e^{\frac{i}{\hbar}\mathcal{S}_0\{\delta\}} \right). \quad (14.60)$$

An interesting quantity to consider is the Ermakov invariant [65]. This has been rediscovered by Lewis [66] and further investigated also in connection with the Milne equation [67] (see [68] for related investigations). In the context of the SE, the Ermakov invariant has been considered by Floyd in [43]

$$I = (2m)^{-1/2} \left[p\psi_E^2 + \hbar^2(\psi_E\partial_q p^{-1/2} - p^{-1/2}\partial_q\psi_E)^2 \right]. \quad (14.61)$$

This invariant can be constructed in terms of any solution of the SE. Once the form (14.60) of ψ_E is given, any complex Möbius transformation of $e^{\frac{2i}{\hbar}\mathcal{S}_0}$, which according to (14.60) will in general correspond to a linear transformation $\psi_E \longrightarrow \tilde{\psi}_E = a\psi_E^D + b\psi_E$, still solution of the SE, leaves I

invariant. The Ermakov invariant was evaluated in [43] in terms of the parameters defining p and of the coefficients in the expression of ψ_E in terms of ψ^D and ψ . Here we have

$$p\psi_E^2 - A^2 e^{-\frac{2i}{\hbar}S_0} - B^2 e^{\frac{2i}{\hbar}S_0} = 2AB = \hbar^2 (\psi_E \partial_q p^{-1/2} - p^{-1/2} \partial_q \psi_E)^2 + A^2 e^{-\frac{2i}{\hbar}S_0} + B^2 e^{\frac{2i}{\hbar}S_0}, \quad (14.62)$$

so that

$$I = (2m)^{-1/2} 4AB. \quad (14.63)$$

In the case in which either A or B vanish, we have $I = 0$. Let us set (recall that $W \in \mathbb{R} \setminus \{0\}$)

$$\epsilon = \frac{W(\ell + \bar{\ell})}{|W(\ell + \bar{\ell})|} = \text{sgn}[W(\ell + \bar{\ell})], \quad (14.64)$$

and define

$$\phi = \sqrt{2} \frac{e^{-i\frac{\alpha}{2}}(\psi^D - i\ell\psi)}{\hbar^{1/2}|W(\ell + \bar{\ell})|^{1/2}}. \quad (14.65)$$

It follows from (14.57) and (14.65) that

$$e^{\frac{2i}{\hbar}S_0\{\delta\}} = \frac{\bar{\phi}}{\phi}, \quad (14.66)$$

and since $\phi'\bar{\phi} - \phi\bar{\phi}' = -2i\epsilon/\hbar$, we have

$$p = \frac{\hbar}{2i} \partial_q \ln \frac{\phi}{\phi} = \epsilon |\phi|^{-2}, \quad (14.67)$$

so that $\epsilon = \pm 1$ fixes the direction of motion. By (14.66) and (14.67) we have

$$\phi = \frac{\epsilon^{1/2}}{\sqrt{S_0'\{\delta\}}} e^{-\frac{i}{\hbar}S_0\{\delta\}}, \quad (14.68)$$

so that we obtain the expression of the wave-function in terms of ϕ

$$\psi_E\{\delta\} = A\epsilon^{-1/2}\phi + B\epsilon^{1/2}\bar{\phi}, \quad (14.69)$$

where we used $\overline{\epsilon^{-1/2}} = \epsilon^{1/2}$. Note that by $\bar{\phi} \not\propto \phi$ and $\phi'\bar{\phi} - \phi\bar{\phi}' \neq 0$, it follows that ϕ never vanishes. Therefore, a necessary condition for a solution of the SE to have zeroes is that it must be proportional to a real function. Hence, if ψ_E has zeroes, then $A\epsilon^{-1/2}\phi + B\epsilon^{1/2}\bar{\phi} \propto \bar{A}\epsilon^{1/2}\bar{\phi} + \bar{B}\epsilon^{-1/2}\phi$, that is $|A| = |B|$, in agreement with the fact that $\psi_E'\bar{\psi}_E - \bar{\psi}_E'\psi_E = -2i\epsilon(|A|^2 - |B|^2)/\hbar$ can be seen as a Wronskian only if $\bar{\psi}_E \not\propto \psi_E$. We also note that in the case in which $\psi_E = \psi$, we have

$$A = i \left[\frac{e^{i\alpha}\hbar W}{2(\ell + \bar{\ell})} \right]^{1/2}, \quad B = -\epsilon e^{-i\alpha} A = \bar{A}, \quad (14.70)$$

where the identity $-\epsilon e^{-i\alpha} A = \bar{A}$ follows from $\text{sgn}[W/(\ell + \bar{\ell})] = \text{sgn}[W(\ell + \bar{\ell})]$. An interesting question is to find the transformations $\delta \longrightarrow \delta'$ leaving the state described by $\psi_E\{\delta\}$ invariant. To this end it

is useful to write $e^{\frac{2i}{\hbar}\mathcal{S}_0}$ in a different form. First of all note that the expression of $e^{\frac{2i}{\hbar}\mathcal{S}_0}$ in (14.57) can be seen as the composition of two maps. The first one is the Cayley transformation

$$w \longrightarrow z = \sigma(w) = \frac{w+i}{w-i} = \frac{\psi^D + i\psi}{\psi^D - i\psi} \in S^1, \quad (14.71)$$

where σ is the matrix

$$\sigma = \frac{1}{(-2i)^{1/2}} \begin{pmatrix} 1 & i \\ 1 & -i \end{pmatrix}. \quad (14.72)$$

Then $e^{\frac{2i}{\hbar}\mathcal{S}_0}$ is obtained as the Möbius transformation

$$e^{\frac{2i}{\hbar}\mathcal{S}_0} = \gamma_{\mathcal{S}_0}(z) = \frac{az+b}{cz+d}, \quad (14.73)$$

where, by (14.57), the entries a, b, c, d of $\gamma_{\mathcal{S}_0}$ are

$$\gamma_{\mathcal{S}_0} = \frac{1}{2^{1/2}|\ell + \bar{\ell}|^{1/2}} \begin{pmatrix} e^{\frac{i}{2}\alpha}(1 + \bar{\ell}) & e^{\frac{i}{2}\alpha}(1 - \bar{\ell}) \\ e^{-\frac{i}{2}\alpha}(1 - \ell) & e^{-\frac{i}{2}\alpha}(1 + \ell) \end{pmatrix}. \quad (14.74)$$

Note that $d = \bar{a}$, $c = \bar{b}$ and $\mu \equiv \det \gamma_{\mathcal{S}_0} = \text{sgn Re } \ell$, so that, if $\text{Re } \ell > 0$, then $\gamma_{\mathcal{S}_0} \in SU(1, 1)$. By (14.60) and (14.73) we can write ψ_E in the form

$$\psi_E\{\delta\} = \left(\frac{2i}{\hbar \partial_q \gamma_{\mathcal{S}_0}(z)} \right)^{1/2} (A + B \gamma_{\mathcal{S}_0}(z)). \quad (14.75)$$

Under the transformation

$$\delta \longrightarrow \delta' = \{\alpha', \ell'\}, \quad (14.76)$$

we have

$$e^{\frac{2i}{\hbar}\mathcal{S}_0\{\delta\}} \longrightarrow e^{\frac{2i}{\hbar}\mathcal{S}_0\{\delta'\}} = \gamma'_{\mathcal{S}_0}(z) = e^{i\alpha'} \frac{w + i\bar{\ell}'}{w - i\ell'}, \quad (14.77)$$

where $\gamma'_{\mathcal{S}_0}$ is the matrix (14.74) with α and ℓ replaced by α' and ℓ' respectively. For our purpose it is useful to write $e^{\frac{2i}{\hbar}\mathcal{S}_0\{\delta'\}}$ as the Möbius transformation of $e^{\frac{2i}{\hbar}\mathcal{S}_0\{\delta\}}$, that is

$$e^{\frac{2i}{\hbar}\mathcal{S}_0\{\delta'\}} = \tilde{\gamma}_{\mathcal{S}_0}(\gamma_{\mathcal{S}_0}(z)), \quad (14.78)$$

where

$$\tilde{\gamma}_{\mathcal{S}_0} = \gamma'_{\mathcal{S}_0} \gamma_{\mathcal{S}_0}^{-1}. \quad (14.79)$$

We can now determine the transformations (14.76) such that

$$\psi_E\{\delta'\} = \left(\frac{2i}{\hbar \partial_q \gamma'_{\mathcal{S}_0}(z)} \right)^{1/2} (A + B \gamma'_{\mathcal{S}_0}(z)), \quad (14.80)$$

describes the same state described by $\psi_E\{\delta\}$. In other words, we are considering the transformations of the integration constants of the QSHJE (9.44), corresponding to real p , such that $\psi_E\{\delta\}$ remains unchanged up to some multiplicative constant c , that is

$$\psi_E\{\delta\} \longrightarrow \psi_E\{\delta'\} = c \psi_E\{\delta\}. \quad (14.81)$$

In order to compute $\psi_E\{\delta'\}$ we observe that by (14.79)

$$\partial_q \gamma'_{S_0} = \partial_q \tilde{\gamma}_{S_0}(\gamma_{S_0}(z)) = \frac{\partial \tilde{\gamma}_{S_0}}{\partial \gamma_{S_0}} \partial_q \gamma_{S_0} = \tilde{\mu} \frac{\partial_q \gamma_{S_0}}{(\tilde{c} \gamma_{S_0} + \tilde{d})^2}, \quad (14.82)$$

where $\tilde{\mu} \equiv \det \tilde{\gamma}_{S_0} = \text{sgn Re } \tilde{\ell}$ and

$$\tilde{\gamma}_{S_0}(\gamma_{S_0}) = \frac{\tilde{a} \gamma_{S_0} + \tilde{b}}{\tilde{c} \gamma_{S_0} + \tilde{d}}, \quad (14.83)$$

with

$$\tilde{a} = \bar{\tilde{d}} = e^{\frac{i}{2}\tilde{\alpha}} \frac{1 + \tilde{\ell}}{2^{1/2}|\tilde{\ell} + \bar{\tilde{\ell}}|^{1/2}}, \quad \tilde{b} = \bar{\tilde{c}} = e^{\frac{i}{2}\tilde{\alpha}} \frac{1 - \tilde{\ell}}{2^{1/2}|\tilde{\ell} + \bar{\tilde{\ell}}|^{1/2}}, \quad (14.84)$$

which are the matrix elements of $\tilde{\gamma}_{S_0}$ given by (14.74) with the δ -moduli replaced by $\tilde{\delta} = \{\tilde{\alpha}, \tilde{\ell}\}$. Therefore, by (14.80) and (14.82)

$$\psi_E\{\delta'\} = \left(\frac{2i}{\tilde{\mu} \hbar \partial_q \gamma_{S_0}} \right)^{1/2} [A\tilde{d} + B\tilde{b} + (A\tilde{c} + B\tilde{a})\gamma_{S_0}], \quad (14.85)$$

and Eq.(14.81) gives

$$A^2 \bar{\tilde{b}} + AB \tilde{a} = AB \bar{\tilde{a}} + B^2 \tilde{b}. \quad (14.86)$$

Note that if either $A = 0$ or $B = 0$, then $\tilde{b} = 0$, *i.e.* $\tilde{\ell} = 1$, and the only transformation leaving invariant the unit ray Ψ_E associated to ψ_E corresponds, as obvious, to adding a constant to $\mathcal{S}_0$. In the case in which ψ_E is proportional to a real function, one has

$$B = e^{-i\gamma} A, \quad (14.87)$$

for some $\gamma \in \mathbb{R}$. Eq.(14.86) and (14.87) imply that the matrix elements of the transformations leaving the unit ray invariant satisfy

$$\text{Im } \tilde{a} = \text{Im } (e^{-i\gamma} \tilde{b}). \quad (14.88)$$

In the case in which $\psi_E = \psi$, by (14.70) and (14.87) we have $\gamma = \alpha - (\epsilon + 1 + 4k)\pi/2$, $k \in \mathbb{Z}$.

While Ψ_E is invariant under the transformations satisfying (14.86)

$$\Psi_E\{\delta\} \longrightarrow \Psi_E\{\delta'\} = \Psi_E\{\delta\}, \quad (14.89)$$

in the case of the conjugate momentum we have

$$p\{\delta\} = \frac{\hbar}{2i} \partial_q \ln \gamma_{S_0} \longrightarrow p\{\delta'\} = \frac{\hbar}{2i} \partial_q \ln \gamma'_{S_0} = \frac{\tilde{\mu} \gamma_{S_0}}{(\tilde{a} \gamma_{S_0} + \tilde{b})(\tilde{b} \gamma_{S_0} + \tilde{a})} p\{\delta\}. \quad (14.90)$$

15 Trajectories and the Equivalence Principle

We now start considering the conditions on $\mathcal{S}_0$ coming from the EP. In other words, until now we did not care about the differential properties of $\mathcal{S}_0$; we just implicitly assumed that $\mathcal{S}_0$ had such properties that the EP could be implemented. In other words, instead of defining the properties $\mathcal{S}_0$ should have in order for the relevant equations to be defined, we did not mention them reserving part of this section to this basic issue. This approach is particularly convenient as the conditions for the implementation of the EP reduce to the conditions that $\mathcal{S}_0$ should satisfy in order for the QSHJE to exist. As we will see, deriving these conditions will lead to the quantization of the energy without making use of the conventional axiomatic interpretation of the wave-function. This is an important result of our approach as we derive a basic fact directly from the EP. In this section we will also investigate in detail the symmetry properties of the trivializing map. As we will see, the role of the Möbius transformation is that of making the generally projective transformation, connecting the trivializing map and the ratio $w = \psi^D/\psi$, equivalent to an affine transformation.

The conditions concerning the existence of the QSHJE are closely related to the nature of the trivializing map. In particular, we will see that this corresponds to a local homeomorphism of the extended real line into itself. In this context we will see that there is an underlying Möbius symmetry which is related to the EP. Next, we will consider the case of the free particle and will define the time parameterization by following Floyd's suggestion of using Jacobi's theorem. This will lead to a dynamical equation corresponding to the quantum analogue of Newton's law $F = ma$.

15.1 Differential properties of the reduced action

We saw that implementing the EP the CSHJE is modified by the additive term $\hbar^2\{\mathcal{S}_0, q\}/4m$. In order to discuss the conditions for the existence of the QSHJE (9.44), it is convenient to consider its form (9.42) that by the Möbius symmetry of the Schwarzian derivative and (14.57) is equivalent to

$$\{w, q\} = -\frac{4m}{\hbar^2}\mathcal{W}(q). \quad (15.1)$$

Existence of this equation requires some conditions on the continuity properties of w and its derivatives. Since the QSHJE is the consequence of the EP, we can say that the EP imposes some constraints on $w = \psi^D/\psi$. These constraints are nothing but the existence of QSHJE (9.44) or, equivalently, of Eq.(15.1). That is, implementation of the EP imposes that $\{w, q\}$ exists, so that

$$w \neq \text{cnst}, w \in C^2(\mathbb{R}), \text{ and } \partial_q^2 w \text{ differentiable on } \mathbb{R}. \quad (15.2)$$

Note that requiring w to be of class $C^3(\mathbb{R})$ would be an unjustified restriction as the Schwarzian derivative is defined also in the case in which $\partial_q^3 w$ is discontinuous (in this case $\mathcal{W}$ is discontinuous).

We now show that the conditions (15.2) are not complete. The reason is that, as we have seen, the implementation of the EP requires that the properties of the Schwarzian derivative be satisfied. Actually, its very properties, derived from the EP, led to the identification $(q^a; q^b) = -\hbar^2\{q^a, q^b\}/4m$. Therefore, in order to implement the EP it is of basic importance that the transformation properties

of the Schwarzian derivative and its symmetries be satisfied. In deriving the transformation properties of $(q^a; q^b)$ we noticed how, besides dilatations and translations there is a highly non-trivial symmetry such as that under inversion. Therefore, we have that (15.1) must be equivalent to

$$\{w^{-1}, q\} = -\frac{4m}{\hbar^2} \mathcal{W}(q). \quad (15.3)$$

A property of the Schwarzian derivative is duality between its entries

$$\{w, q\} = -\left(\frac{\partial w}{\partial q}\right)^2 \{q, w\}. \quad (15.4)$$

This shows that the invariance under inversion of w reflects in the invariance, up to a Jacobian factor, under inversion of q . That is $\{w, q^{-1}\} = q^4 \{w, q\}$, so that the QSHJE (15.1) can be written in the equivalent form

$$\{w, q^{-1}\} = -\frac{4m}{\hbar^2} q^4 \mathcal{W}(q). \quad (15.5)$$

In other words, starting from the EP one can arrive to either Eq.(15.1) or Eq.(15.5). The consequence of this fact is that since under

$$q \longrightarrow \frac{1}{q}, \quad (15.6)$$

$0^\pm$ maps to $\pm\infty$, we have to extend the continuity conditions (15.2) to the point at infinity. In other words, the continuity conditions (15.2) should hold on the extended real line $\hat{\mathbb{R}} = \mathbb{R} \cup \{\infty\}$. This aspect is related to the fact that the Möbius transformations, under which the Schwarzian derivative transforms as a quadratic differential, map circles to circles. We stress that we are considering the systems defined on $\mathbb{R}$ and not $\hat{\mathbb{R}}$. What happens is that the existence of QSHJE forces us to impose smoothly joining conditions even at $\pm\infty$, that is (15.2) must be extended to

$$w \neq \text{const}, \quad w \in C^2(\hat{\mathbb{R}}), \quad \text{and } \partial_q^2 w \text{ differentiable on } \hat{\mathbb{R}}. \quad (15.7)$$

As we will see, w is a Möbius transformation of the trivializing map. Therefore, Eq.(15.4), which is defined if and only if $w(q)$ can be inverted, that is if $\partial_q w \neq 0, \forall q \in \mathbb{R}$, is a consequence of the cocycle condition (8.14). By (15.5) we see that also local univalence should be extended to $\hat{\mathbb{R}}$. This implies the following joining condition at spatial infinity

$$w(-\infty) = \begin{cases} w(+\infty), & \text{for } w(-\infty) \neq \pm\infty, \\ -w(+\infty), & \text{for } w(-\infty) = \pm\infty. \end{cases} \quad (15.8)$$

As illustrated by the non-univalent function $w = q^2$, the apparently natural choice $w(-\infty) = w(+\infty)$, one would consider also in the $w(-\infty) = \pm\infty$ case, does not satisfy local univalence.

15.2 The Equivalence Principle and the trivializing map

We can now derive the explicit expression of the trivializing map $q \longrightarrow q^0$ under which a state $\mathcal{W}$ reduces to $\mathcal{W}^0$. A first remark is that, as we have seen, for a given $\mathcal{W}$ there are different possible $\mathcal{S}_0$'s

which are parameterized by the δ -moduli. We already derived the structure of the trivializing map in (10.34). Here we derive the explicit expression of the trivializing map as function of the δ -moduli. This expression follows directly from $\mathcal{S}_0^0(q^0) = \mathcal{S}_0(q)$. We simply observe that for the state $\mathcal{W}^0$ we can choose, as in (14.31), $\psi^{D^0} = q^0$ and $\psi^0 = 1$. Then by (14.57) and $\mathcal{S}_0^0(q^0) = \mathcal{S}_0(q)$ we have

$$e^{i\alpha_0} \frac{q^0 + i\bar{\ell}_0}{q^0 - i\ell_0} = e^{i\alpha} \frac{w + i\bar{\ell}}{w - i\ell}. \quad (15.9)$$

Therefore, the trivializing map transforming the state $\mathcal{W}$ with moduli $\delta = \{\alpha, \ell\}$ into the state $\mathcal{W}^0$ with moduli $\delta_0 = \{\alpha_0, \ell_0\}$ is given by the real map

$$q^0 = \frac{(\ell_0 e^{i\beta} + \bar{\ell}_0 e^{-i\beta})w + i\ell_0 \bar{\ell} e^{i\beta} - i\bar{\ell}_0 \ell e^{-i\beta}}{2w \sin \beta + \ell e^{-i\beta} + \bar{\ell} e^{i\beta}}, \quad (15.10)$$

where $\beta = (\alpha - \alpha_0)/2$. Let us consider the case in which the functional structure of two reduced actions differs only by a constant, that is

$$\mathcal{S}_0^a(q^a) = \mathcal{S}_0^b(q^a) + \hbar(\alpha_a - \alpha_b)/2. \quad (15.11)$$

It is clear that these two reduced actions lead to the same conjugate momentum, that is to the same physical state. In fact, by (15.11) we have

$$p_a(q^a) = \frac{\partial \mathcal{S}_0^a(q^a)}{\partial q^a} = \frac{\partial \mathcal{S}_0^b(q^a)}{\partial q^a} = \frac{\partial \mathcal{S}_0^b(q^b)}{\partial q^b} \Big|_{q^b=q^a} = p_b(q^a). \quad (15.12)$$

Therefore, while for general reduced actions $\mathcal{S}_0^a$ and $\mathcal{S}_0^b$ one has $p_a(q) \neq p_b(q)$, in the case (15.11) the functional dependence of p_a and p_b on their arguments coincides and the only effect of the identification $\mathcal{S}_0^a(q^a) = \mathcal{S}_0^b(q^b)$ is a trivial re-labeling of the coordinate. Thus, even if $p_a(q^a) \neq p_b(q^b)$, we have $p_a(q) = p_b(q)$. Therefore, since the transformations (15.10) that differ only by the value of β connect the same pair of states, we can set $\alpha = \alpha_0 + 2k\pi$, and (15.10) becomes

$$q^0 = \frac{(\ell_0 + \bar{\ell}_0)w + i\ell_0 \bar{\ell} - i\bar{\ell}_0 \ell}{\ell + \bar{\ell}}. \quad (15.13)$$

We will call Möbius states the states parameterized by ℓ associated with a given $\mathcal{W}$.

15.3 Equivalence Principle and local homeomorphicity of v -maps

We now discuss some basic properties of the v -maps. First we note that consistency of the EP requires continuity of the v -maps: since both q and q^v take values continuously on $\mathbb{R}$, it is clear that full equivalence between the two systems requires that the v -maps should be continuous. Next note that the EP implies that there always exists the trivializing map $q \longrightarrow q^0$ under which $\mathcal{W} \longrightarrow \mathcal{W}^0$. Furthermore, since the transformation should exist for any pair of $\mathcal{W}$ states, we have that v -maps, including the trivializing map, should be locally invertible. This provides the pseudogroup property.

We saw that all the above continuity properties hold. In fact these are the properties we derived for w as a condition for the existence of the QSHJE. On the other hand, it follows by (15.10) that the

properties of w reflect on the properties of the trivializing map. As for w , even in considering v -maps we should consider $\hat{\mathbb{R}}$. In particular, from the properties of w and by (15.10) it follows that the v -maps are local self-homeomorphisms of $\hat{\mathbb{R}}$. Let us map $\hat{\mathbb{R}}$ to S^1 by means of a Cayley transformation and then consider the case of the trivializing map. While

$$z = \frac{q - i}{1 - iq}, \quad (15.14)$$

spans S^1 once, we have that

$$z^0 = \frac{q^0 - i}{1 - iq^0}, \quad (15.15)$$

should run continuously around S^1 . Since the Cayley transformation is a global univalent transformation, we have that the v -maps induce local self-homeomorphisms of S^1 . We also note that since local homeomorphisms are closed under composition, it follows that local homeomorphicity of any v -map also follows from local homeomorphicity of the trivializing map.

15.4 The Möbius symmetry of the trivializing map

Let us further illustrate the properties of the trivializing map. Choosing a pair of real linearly independent solutions of the SE different from (ψ^D, ψ) , reflects in a real Möbius transformation of w

$$w \longrightarrow \tilde{w} = \frac{aw + b}{cw + d}. \quad (15.16)$$

By (14.57), the effect of this transformation on $\mathcal{S}_0$ is

$$e^{\frac{2i}{\hbar}\mathcal{S}_0\{\delta\}} \longrightarrow e^{\frac{2i}{\hbar}\mathcal{S}_0\{\delta'\}} = e^{i\alpha} \left(\frac{a + i\bar{\ell}c}{a - i\ell c} \right) \frac{w + i(\bar{\ell}d - ib)/(a + i\bar{\ell}c)}{w - i(\ell d + ib)/(a - i\ell c)} = e^{i\alpha'} \frac{w + i\bar{\ell}'}{w - i\ell'}, \quad (15.17)$$

where

$$e^{i\alpha'} = e^{i\alpha} \frac{a + i\bar{\ell}c}{a - i\ell c}, \quad \ell' = \frac{\ell d + ib}{a - i\ell c}. \quad (15.18)$$

Considering the fixed point equation

$$\ell' = \ell, \quad (15.19)$$

we see that the transformations (15.16) satisfying

$$\ell = i \frac{d - a \pm \sqrt{(d - a)^2 + 4bc}}{2c}, \quad (15.20)$$

correspond to changing only the phase α and do not affect the conjugate momentum. Since the transformation (15.16) corresponds to

$$\tilde{\psi}^D = a\psi^D + b\psi, \quad \tilde{\psi} = c\psi^D + d\psi, \quad (15.21)$$

we have that replacing (ψ^D, ψ) in (14.29) with $(\tilde{\psi}^D, \tilde{\psi})$, has no effect on p if the transformation (15.21) satisfies (15.20). This analysis also shows that the transformation of w connecting (15.10) and (15.13)

has the structure (15.16)(15.20). This is an interesting property of the trivializing map. In particular, (15.10) and (15.13) show that there is a symmetry underlying the trivializing map, and therefore, by the pseudogroup property, the v -maps. To see this observe that (15.19) is equivalent to

$$d = a - i\ell c - i\ell^{-1}b. \quad (15.22)$$

On the other hand, reality of a, b, c, d implies the condition

$$c = -\frac{b^2}{|\ell|^2}. \quad (15.23)$$

This means that all the transformations of the kind

$$\begin{pmatrix} \psi^D \\ \psi \end{pmatrix} \longrightarrow \begin{pmatrix} \tilde{\psi}^D \\ \tilde{\psi} \end{pmatrix} = \begin{pmatrix} a & b \\ -b|\ell|^{-2} & a + ib(\bar{\ell}^{-1} - \ell^{-1}) \end{pmatrix} \begin{pmatrix} \psi^D \\ \psi \end{pmatrix}, \quad (15.24)$$

induce a transformation of q^0 under which $\mathcal{W}^0$ remains invariant. Hence, the trivializing map is defined up to a group of transformations that we now derive. First note that the effect of (15.24) on w is

$$w \longrightarrow \tilde{w} = \tilde{\gamma}(w) = \frac{a|\ell|^2 w + b|\ell|^2}{-bw + a|\ell|^2 + ib(\ell - \bar{\ell})}, \quad (15.25)$$

where $\tilde{\gamma}$ is the matrix in (15.24). From Eq.(15.13) we have

$$w = \gamma(q^0), \quad (15.26)$$

where

$$\gamma = (\ell_0 + \bar{\ell}_0)^{-1} \begin{pmatrix} \ell + \bar{\ell} & i\bar{\ell}_0\ell - i\ell_0\bar{\ell} \\ 0 & \ell_0 + \bar{\ell}_0 \end{pmatrix}. \quad (15.27)$$

By (15.25) and (15.26) we have

$$\tilde{w} = \tilde{\gamma}(w) = \tilde{\gamma}(\gamma(q^0)) = \tilde{\gamma} \cdot \gamma(q^0). \quad (15.28)$$

The effect on q^0 induced by the transformation (15.24) is defined by

$$\tilde{w} = \gamma(\hat{\gamma}(q^0)) = \gamma \cdot \hat{\gamma}(q^0), \quad (15.29)$$

that compared with (15.28) gives

$$\hat{\gamma} = \gamma^{-1} \cdot \tilde{\gamma} \cdot \gamma. \quad (15.30)$$

Therefore, given the trivializing map $q \longrightarrow q^0 = v^0(q)$ connecting a Möbius state of $\mathcal{W}$ to a Möbius state of $\mathcal{W}^0$, we have that any other map defined by

$$\hat{q}^0 = \hat{\gamma}(q^0), \quad (15.31)$$

still connects the same Möbius states. These transformations can be parameterized by β in (15.10). Let us denote q^0 in (15.10) by q_β^0 , so that q^0 in (15.13) corresponds to $q_{k\pi}^0$. By (15.10) we have

$$q_\beta^0 = \gamma_1(w), \quad (15.32)$$

where

$$\gamma_1 = \begin{pmatrix} \ell_0 e^{i\beta} + \bar{\ell}_0 e^{-i\beta} & i\ell_0 \bar{\ell} e^{i\beta} - i\bar{\ell}_0 \ell e^{-i\beta} \\ 2 \sin \beta & \ell e^{-i\beta} + \bar{\ell} e^{i\beta} \end{pmatrix}. \quad (15.33)$$

On the other hand, by (15.26) and (15.32) we have $q_\beta^0 = \gamma_1 \cdot \gamma(q^0)$, so that

$$q_\beta^0 = \gamma_\beta(q^0), \quad (15.34)$$

where

$$\gamma_\beta = \gamma_1 \cdot \gamma = \frac{\ell + \bar{\ell}}{\ell_0 + \bar{\ell}_0} \begin{pmatrix} \ell_0 e^{i\beta} + \bar{\ell}_0 e^{-i\beta} & -2|\ell_0|^2 \sin \beta \\ 2 \sin \beta & \ell_0 e^{-i\beta} + \bar{\ell}_0 e^{i\beta} \end{pmatrix}. \quad (15.35)$$

Therefore, we have that the trivializing map is defined up to the Möbius transformation

$$q_\beta^0 = \frac{(\ell_0 e^{i\beta} + \bar{\ell}_0 e^{-i\beta})q^0 - 2|\ell_0|^2 \sin \beta}{2q^0 \sin \beta + \ell_0 e^{-i\beta} + \bar{\ell}_0 e^{i\beta}}. \quad (15.36)$$

Note that for $\ell_0 = |\ell_0| \exp -i[\beta + (2k+1)\pi/2]$, $k \in \mathbb{Z}$, we have

$$q_\beta^0 = -\frac{|\ell_0|^2}{q^0 + 2e^{k\pi i}|\ell_0| \cos \beta}, \quad (15.37)$$

that for $\beta = (2n+1)\pi/2$, $n \in \mathbb{Z}$, becomes

$$q_{(2n+1)\pi/2}^0 = -\frac{|\ell_0|^2}{q^0}. \quad (15.38)$$

Note that Eq.(15.36) can be also directly derived from (15.10) by considering the case in which $w = q^0$ and $\ell = \ell_0$, corresponding to

$$e^{i\alpha_0} \frac{q_\beta^0 + i\bar{\ell}_0}{q_\beta^0 - i\ell_0} = e^{i\alpha} \frac{q^0 + i\bar{\ell}_0}{q^0 - i\ell_0}. \quad (15.39)$$

This represents the transformation connecting the same Möbius state but with the action having a different α . Since this constant has no dynamical effects, the Möbius transformations (15.36) correspond to a one-parameter group of symmetries of the trivializing map.

It is worth stressing another property of the trivializing map. Namely, we have seen that it is a Möbius, or projective, transformation (15.10) of the ratio w . On the other hand, this can be reduced to the affine mapping (15.13). Recall that in proving that the relation Eq.(8.14) implies that $(q^a, q^b) \propto \{q^a, q^b\}$, we mentioned the fact that any element g in the family of one-dimensional complex analytic local homeomorphisms, such that either $g''(x)/g'(x) = 0$ or $\{g(x), x\} = 0$, are the unique solutions of a system of differential equations involving only the first and higher derivatives, satisfying the pseudogroup property [35][36]. Then, in this context it is interesting to observe that according to (15.10) we have $\{w, q^0\} = 0$ which by (15.13) reduces to $\partial_{q^0}^2 w / \partial_{q^0} w = 0$.

Let us denote by q_a and q_b the coordinates of two Möbius states associated to the same $\mathcal{W}$. While in the case of the state $\mathcal{W}^0$, q_a^0 and q_b^0 are related by the affine transformation

$$q_b^0 = \frac{(\ell_{0,b} + \bar{\ell}_{0,b})q_a^0 + i\ell_{0,b}\bar{\ell}_{0,a} - i\bar{\ell}_{0,b}\ell_{0,a}}{\ell_{0,a} + \bar{\ell}_{0,a}}, \quad (15.40)$$

in the case of Möbius states associated to arbitrary $\mathcal{W}$ states, the v -map $q_a \longrightarrow q_b = v^{ba}(q_a)$ corresponds to the affine transformation of the w 's

$$w_b = \frac{(\ell_b + \bar{\ell}_b)w_a + i\ell_b\bar{\ell}_a - i\bar{\ell}_b\ell_a}{\ell_a + \bar{\ell}_a}, \quad (15.41)$$

where $w_k \equiv w(q_k)$, $k = a, b$. The v -map connecting two different states, say $\mathcal{W}^a(q^a)$ and $\mathcal{W}^b(q^b)$, is still defined by (15.41) where now $w_k \equiv w_k(q_k) = \psi_k^D(q_k)/\psi_k(q_k)$, with (ψ_k^D, ψ_k) a pair of linearly independent real solutions of the SE defined by $\mathcal{W}^k(q^k)$.

15.5 The free particle

A feature of our formulation is that no use of the usual axioms of QM is made. In particular, the SE arises as mathematical tool to linearize the QSHJE (9.44). We now start studying the physical consequences of the QSHJE which is seen as an equation for trajectories. While the concept of trajectories in QM is reminiscent of Bohm's theory, we will see that there are basic differences. A first important aspect is that the conjugate momentum $p = \partial_q \mathcal{S}_0$, which is a real quantity even in the classically forbidden regions, has a well-defined and natural classical limit. Furthermore, besides the fact that there is no pilot-wave guide, we will also see that $p \neq m\dot{q}$. This is a consequence of the definition of time parameterization which follows from Jacobi's theorem as done by Floyd in [15].

A basic request to formulating a trajectory interpretation of the QSHJE would be the derivation of the main features of QM without assuming its axioms. Since tunnelling and energy quantization are basic features distinguishing QM from CM, we have to understand whether the QSHJE reproduces such phenomena. We will give a detailed proof that this is in fact the case: both the tunnel effect and energy quantization are predictions of the EP and in particular of the QSHJE which follows from it. In this respect let us recall that in the conventional formulation of QM the tunnel effect is a consequence of the wave-function interpretation in terms of probability amplitude of finding the particle in a measurement process. Since in the classically forbidden regions one generally has $\psi \neq 0$, there is a non-zero probability of finding the particle in the region where $V - E > 0$. On the other hand, we have seen that $p = \epsilon|\phi|^{-2}$, which is a real quantity even in the classically forbidden regions. As we will see, the reality property inside the classically forbidden region is a property which holds also for the velocity field.

Another feature, which is a consequence of the wave-function interpretation, concerns the quantization of the energy spectrum. In particular, since $|\psi|^2 dq$ is the probability of finding the particle in the interval $[q, q + dq]$, it follows that there are situations (bound states) in which the SE admits a solution ψ , identified with the wave-function, belonging to $L^2(\mathbb{R})$. We will see that in our formulation this condition follows from the local homeomorphicity of v -maps, a condition implied by the EP! This condition directly follows from (15.7) which concerns the existence of the QSHJE.

The above aspects will be discussed in detail later on. We now consider the solutions of the QSHJE for the free particle of energy E . In order to find the conjugate momentum, we first have to choose two real linearly independent solutions of the SE. Let us set

$$\psi^D = a \sin(kq), \quad \psi = b \cos(kq), \quad (15.42)$$

where $k = \sqrt{2mE}/\hbar$. For the Wronskian we have $W = -abk$. By (14.29) and (15.42) we have

$$p_E = \pm \frac{\hbar(\ell_E + \bar{\ell}_E)abk}{2|a \sin(kq) - i\ell_E b \cos(kq)|^2}. \quad (15.43)$$

An important topic of our approach that will be discussed later, concerns the $E \rightarrow 0$ and $\hbar \rightarrow 0$ limits. In doing this we have to consider $\mathcal{S}_0$ or p . This implies that these limits should be performed by considering ψ^D , ψ and ℓ simultaneously. However, since any linear transformation of ψ^D and ψ reflects in a transformation of ℓ and α , in considering the $E \rightarrow 0$ limit of ψ^D and ψ , we can also require that the corresponding functions reduce to the one of the state $\mathcal{W}^0$

$$\lim_{E \rightarrow 0} a\psi^D = q, \quad \lim_{E \rightarrow 0} b\psi = 1. \quad (15.44)$$

A possible solution is $a = k^{-1}$, $b = 1$, so that $W = -1$ and Eq.(15.43) becomes

$$p_E = \pm \frac{\hbar(\ell_E + \bar{\ell}_E)}{2|k^{-1} \sin(kq) - i\ell_E \cos(kq)|^2}. \quad (15.45)$$

15.6 Time parameterization

It is well-known that while in CM the regions where $V - E > 0$ are forbidden this is not the case in QM. In our approach the tunnel effect is due to the fact that p is a real function even in the classically forbidden regions, a consequence of the quantum potential which, unlike in Bohm theory, is never trivial. This is the case also for the state $\mathcal{W}^0$. As stressed in [13], this property is reminiscent of the relativistic rest energy.¹⁴ As a consequence of the quantum potential, the dynamics of the system is rather different from the classical one. In order to specify it, we have to introduce time parameterization. In [15] Floyd made the interesting proposal of using Jacobi's theorem. According to this theorem, time parameterization is given by the partial derivative of the reduced action with respect to E , that is

$$t - t_0 = \frac{\partial \mathcal{S}_0}{\partial E}. \quad (15.46)$$

Using the QSHJE (9.44), we can write (15.46) in the form

$$t - t_0 = \frac{\partial}{\partial E} \int_{q_0}^q dx \frac{\partial \mathcal{S}_0}{\partial x} = \left(\frac{m}{2}\right)^{1/2} \int_{q_0}^q dx \frac{1 - \partial_E Q}{(E - V - Q)^{1/2}}, \quad (15.47)$$

where $q_0 \equiv q(t_0)$. The velocity and acceleration are [15]

$$\frac{dq}{dt} = \left(\frac{dt}{dq}\right)^{-1} = \frac{\partial_q \mathcal{S}_0}{m(1 - \partial_E \mathcal{V})}, \quad (15.48)$$

and

$$\ddot{q} = -\dot{q}^3 \frac{d^2 t}{dq^2} = \frac{2(E - \mathcal{V})\partial_q \partial_E \mathcal{V}}{m(1 - \partial_E \mathcal{V})^3} - \frac{\partial_q \mathcal{V}}{m(1 - \partial_E \mathcal{V})^2}, \quad (15.49)$$

where $\mathcal{V}$ denotes the “effective potential”

$$\mathcal{V}(q) \equiv V(q) + Q(q). \quad (15.50)$$

¹⁴In a different context the usual quantum potential was considered in [14] as an internal potential.

15.7 Quantum mass field: dynamical equation

Observe that the deformation of the classical relation between conjugate momentum and velocity is provided by the quantum potential through the relation

$$\dot{q} = \frac{p}{m(1 - \partial_E Q)} \neq \frac{p}{m}. \quad (15.51)$$

The relationship between p and $\dot{q}$ can be written in an alternative way by using the QSHJE (9.44). In fact, the partial derivative of (9.44) with respect to E gives

$$m \frac{\partial Q}{\partial E} = m - p \frac{\partial p}{\partial E}, \quad (15.52)$$

so that (15.51) becomes

$$\dot{q} = \frac{1}{\partial_E p}, \quad (15.53)$$

which is satisfied also classically. The fact that $\dot{q} \neq p/m$ is somehow reminiscent of special relativity in which there is the correction due to the γ factor. In particular, for the free particle we have

$$E = \frac{m}{2} \dot{q}^2 (1 - \partial_E Q)^2 + Q. \quad (15.54)$$

This suggests defining the quantum mass field

$$m_Q = m(1 - \partial_E Q), \quad (15.55)$$

so that

$$p = m_Q \dot{q}. \quad (15.56)$$

This form of the conjugate momentum allows us to write the QSHJE (9.44) in terms of $\dot{q}, \ddot{q}, \dddot{q}$. Using

$$p' = m'_Q \dot{q} + m_Q \frac{\ddot{q}}{\dot{q}}, \quad p'' = m''_Q \dot{q} + 2m'_Q \frac{\ddot{q}}{\dot{q}} + m_Q \left(\frac{\dddot{q}}{\dot{q}^2} - \frac{\ddot{q}^2}{\dot{q}^3} \right), \quad (15.57)$$

and recalling that

$$Q = \frac{\hbar^2}{4m} \{ \mathcal{S}_0, q \} = \frac{\hbar^2}{4m} \left(\frac{p''}{p} - \frac{3p'^2}{2p^2} \right), \quad (15.58)$$

we have that the QSHJE (9.44) is equivalent to the dynamical equation

$$\frac{m_Q^2}{2m} \dot{q}^2 + V(q) - E + \frac{\hbar^2}{4m} \left(\frac{m''_Q}{m_Q} - \frac{3m'_Q{}^2}{2m_Q^2} - \frac{m'_Q}{m_Q} \frac{\ddot{q}}{\dot{q}^2} + \frac{\ddot{q}}{\dot{q}^3} - \frac{5\ddot{q}^2}{2\dot{q}^4} \right) = 0. \quad (15.59)$$

Note that taking the time derivative of this equation yields the quantum analogue of Newton's law. We also observe that the solution of the equation of motion depends on the integration constants $\text{Re } \ell$ and $\text{Im } \ell$ which play the role of hidden variables.

16 Equivalence Principle and fundamental constants

We have seen that the EP implied the QSHJE in which the Planck constant plays the role of covariantizing parameter. The fact that a fundamental constant follows from the EP suggests that other fundamental constants as well may be related to such a principle. One hint in this direction is to observe that the EP we formulated is reminiscent of Einstein's EP [3]. The similarity between our formulation and the basis of GR suggests that the long-standing problem of quantizing gravity is related to a deep relationship between gravity and the foundations of QM. This indicates that the EP should be seen as a principle underlying all possible interactions. In particular, not only gravity but any physical system under the action of an arbitrary potential should be equivalent to the free system. We have seen that this implied the introduction of the quantum potential. Unlike in the conventional approach to QM, this function is never trivial. In particular, by (14.6) it follows that in the case of the state $\mathcal{W}^0$ we have

$$Q^0 = -\frac{p_0^2}{2m} = -\frac{\hbar^2(\ell_0 + \bar{\ell}_0)^2}{8m|q^0 - i\ell_0|^4}. \quad (16.1)$$

We note that, unlike the relativistic rest energy, which depends on the mass only, the quantum potential depends on $V - E$ itself. It is just this peculiar property of Q which makes the tunnel effect possible: the intrinsic “self-energy” Q is the particle reaction to any external potential. This intrinsic effect makes p a real field also in the classically forbidden regions.

The connection between our formulation and GR would suggest the appearance of other fundamental constants besides the Planck constant. Since our formulation concerns QM, it is clear that a connection with GR would suggest the appearance of the Newton constant, in particular one should expect that the Planck length should arise in a natural way. We now follow [13] to show that in considering the classical and $E \rightarrow 0$ limits in the case of the free particle, one has to introduce fundamental lengths. The first condition is that in the $\hbar \rightarrow 0$ limit Eq.(15.45) reduces to the classical solution

$$\lim_{\hbar \rightarrow 0} p_E = \pm \sqrt{2mE}. \quad (16.2)$$

On the other hand, we should also have

$$\lim_{E \rightarrow 0} p_E = p_0 = \pm \frac{\hbar(\ell_0 + \bar{\ell}_0)}{2|q - i\ell_0|^2}, \quad (16.3)$$

in agreement with (14.6). Let us first consider Eq.(16.2). Due to the $\ell_E \cos(kq)$ term in (15.45), we see that existence of the classical limit would imply some condition on ℓ_E . In particular, in order to reach the classical value $\sqrt{2mE}$ in the $\hbar \rightarrow 0$ limit, the quantity ℓ_E should depend on E . Let us set

$$\ell_E \sim k^{-1} + \lambda_E, \quad (16.4)$$

for some λ_E . Recalling that $p = \epsilon|\phi|^{-2}$, we have

$$p_E = \pm \frac{\hbar(\ell_E + \bar{\ell}_E)}{2|k^{-1} \sin(kq) - i\ell_E \cos(kq)|^2} = \pm \frac{\sqrt{2mE} + mE(\lambda_E + \bar{\lambda}_E)/\hbar}{|e^{ikq} + \lambda_E k \cos(kq)|^2}. \quad (16.5)$$

Hence, if $\lambda_E \sim \hbar^n$, $n > 1$, then p_E would satisfy (16.2). However, we now show that the condition (16.3) implies that (16.4), although correct as λ_E is still undetermined, is not the natural one. In fact, to get the limit (16.3) one observes that q in the right hand side arises from the $E \rightarrow 0$ limit of $k^{-1} \sin(kq)$. This shows that to get this limit we cannot manipulate the expression (15.45) by moving k^{-1} from the $k^{-1} \sin(kq)$ term. On the other hand, with the position (16.4), we would have

$$p_E \underset{E \rightarrow 0}{\sim} \frac{2\hbar^2(2mE)^{-1/2} + \hbar(\lambda_E + \bar{\lambda}_E)}{2|q - i\hbar(2mE)^{-1/2} - i\lambda_E|^2}, \quad (16.6)$$

implying a rather involved cancellation of the divergent $E^{-1/2}$ terms which should come from λ_E . This suggests considering the position

$$\ell_E = k^{-1}f(E, \hbar) + \lambda_E, \quad (16.7)$$

where f is dimensionless. Since λ_E is arbitrary, we choose f to be real. By (16.5) and (16.7) we have

$$p_E = \pm \frac{\sqrt{2mE}f(E, \hbar) + mE(\lambda_E + \bar{\lambda}_E)/\hbar}{|e^{ikq} + (f(E, \hbar) - 1 + \lambda_E k) \cos(kq)|^2}, \quad (16.8)$$

so that, forgetting for a moment λ_E , in order to recover the classical result we have that in the $\hbar \rightarrow 0$ limit f has to satisfy the condition

$$\lim_{\hbar \rightarrow 0} f(E, \hbar) = 1. \quad (16.9)$$

On the other hand, cancellation of the divergent term $E^{-1/2}$ in

$$p_E \underset{E \rightarrow 0}{\sim} \frac{\hbar^2(2mE)^{-1/2}f(E, \hbar) + \hbar(\lambda_E + \bar{\lambda}_E)}{|q - i\hbar(2mE)^{-1/2}f(E, \hbar) - i\lambda_E|^2}, \quad (16.10)$$

yields

$$\lim_{E \rightarrow 0} E^{-1/2}f(E, \hbar) = 0. \quad (16.11)$$

This analysis shows that the request that both the $\hbar \rightarrow 0$ and $E \rightarrow 0$ limits exist, is a strong constraint as we have to introduce the dimensionless quantity f which depends both on E and $\hbar$ and satisfies the conditions (16.9) and (16.11). We now see that this suggests introducing the Planck length.

16.1 $\hbar \rightarrow 0$, $E \rightarrow 0$ and Planck length

The limit (16.3) can be seen as the limit in which the trivializing map reduces to the identity. Actually, the trivializing map connecting $\mathcal{W} = -E$ and $\mathcal{W}^0$, reduces to the identity map in the $E \rightarrow 0$ limit. In the above investigation we considered q as an independent variable, however one can also consider $q_E(q^0)$ so that $\lim_{E \rightarrow 0} q_E = q^0$ and in the above formulas one can replace q with q_E .

In considering the two limits $E \rightarrow 0$ and $\hbar \rightarrow 0$, one has to introduce basic lengths. We know from (16.11) that k must enter in the expression of $f(E, \hbar)$. Since f is a dimensionless constant, we

need at least one more constant with the dimension of a length. Two fundamental lengths one can consider are the Compton length

$$\lambda_c = \frac{\hbar}{mc}, \quad (16.12)$$

and the Planck length

$$\lambda_p = \sqrt{\frac{\hbar G}{c^3}}. \quad (16.13)$$

Two dimensionless quantities depending on E are

$$x_c = k\lambda_c = \sqrt{\frac{2E}{mc^2}}, \quad (16.14)$$

and

$$x_p = k\lambda_p = \sqrt{\frac{2mEG}{\hbar c^3}}. \quad (16.15)$$

On the other hand, since x_c does not depend on $\hbar$, this combination is ruled out by (16.9). Then we see that a natural expression for f is a function of the Planck length times k . Let us set

$$f(E, \hbar) = e^{-\alpha(x_p^{-1})}, \quad (16.16)$$

where

$$\alpha(x_p^{-1}) = \sum_{k \geq 1} \alpha_k x_p^{-k}. \quad (16.17)$$

Eqs.(16.9)(16.11) correspond to conditions on the α_k 's. For example, in the case in which one considers α to be the function $\alpha(x_p^{-1}) = \alpha_1 x_p^{-1}$, then by (16.11) we have $\alpha_1 > 0$. Therefore, we have

$$\ell_E = k^{-1} e^{-\alpha(x_p^{-1})} + \lambda_E. \quad (16.18)$$

In order to consider the structure of λ_E , we note that although $e^{-\alpha(x_p^{-1})}$ cancelled the $E^{-1/2}$ divergent term, we still have some conditions to be satisfied. To see this note that

$$p_E = \pm \frac{\sqrt{2mE} e^{-\alpha(x_p^{-1})} + mE(\lambda_E + \bar{\lambda}_E)/\hbar}{\left| e^{ikq} + (e^{-\alpha(x_p^{-1})} - 1 + k\lambda_E) \cos(kq) \right|^2}, \quad (16.19)$$

so that the condition (16.2) implies

$$\lim_{\hbar \rightarrow 0} \frac{\lambda_E}{\hbar} = 0. \quad (16.20)$$

To discuss this limit, we first note that by (16.5) and (16.18)

$$p_E = \pm \frac{2\hbar k^{-1} e^{-\alpha(x_p^{-1})} + \hbar(\lambda_E + \bar{\lambda}_E)}{2 \left| k^{-1} \sin(kq) - i \left(k^{-1} e^{-\alpha(x_p^{-1})} + \lambda_E \right) \cos(kq) \right|^2}. \quad (16.21)$$

So that, since

$$\lim_{E \rightarrow 0} k^{-1} e^{-\alpha(x_p^{-1})} = 0, \quad (16.22)$$

we have by (16.3) and (16.21) that

$$\lim_{E \rightarrow 0} \lambda_E = \lim_{E \rightarrow 0} \ell_E = \ell_0. \quad (16.23)$$

Consider now the limit

$$\lim_{\hbar \rightarrow 0} p_0 = 0. \quad (16.24)$$

Writing p_0 in the equivalent form

$$p_0 = \pm \frac{\hbar(\ell_0 + \bar{\ell}_0)}{2|q^0 - i\ell_0|^2}, \quad (16.25)$$

we see that to get the classical limit, we have to consider both cases $q^0 \neq 0$ and $q^0 = 0$. In particular, we see that the behavior of ℓ_0 should be

$$\ell_0 \sim \hbar^\gamma, \quad -1 < \gamma < 1. \quad (16.26)$$

Lengths having the term $\hbar^\gamma$, $-1 < \gamma < 1$, can be constructed by means of λ_c and λ_p . We also note that a constant length independent from $\hbar$ is given by $\lambda_e = e^2/mc^2$ with e the electric charge.¹⁵ Then ℓ_0 can be considered as a suitable function of λ_c , λ_p and λ_e satisfying the constraint (16.26). Consistency between (16.20) and (16.26) implies that the condition (16.20) is given by the λ_E/λ_0 term of λ_E . In particular, the above investigation indicates that a natural way to express λ_E is

$$\lambda_E = e^{-\beta(x_p)} \lambda_0, \quad (16.27)$$

where

$$\beta(x_p) = \sum_{k \geq 1} \beta_k x_p^k. \quad (16.28)$$

Any possible choice should satisfy the conditions (16.20) and (16.23). For example, for ℓ_E built with $\beta(x_p) = \beta_1 x_p$, one should have $\beta_1 > 0$. Summarizing, we have

$$\ell_E = k^{-1} e^{-\alpha(x_p^{-1})} + e^{-\beta(x_p)} \ell_0, \quad (16.29)$$

where

$$\ell_0 = \ell_0(\lambda_c, \lambda_p, \lambda_e), \quad (16.30)$$

and for the conjugate momentum of the state $\mathcal{W} = -E$, we have

$$p_E = \pm \frac{2k^{-1} \hbar e^{-\alpha(x_p^{-1})} + \hbar e^{-\beta(x_p)} (\ell_0 + \bar{\ell}_0)}{2 \left| k^{-1} \sin(kq) - i \left(k^{-1} e^{-\alpha(x_p^{-1})} + e^{-\beta(x_p)} \ell_0 \right) \cos(kq) \right|^2}. \quad (16.31)$$

We observe that the above investigation answers the question posed at the end of subsect.10.4, where we noticed non-triviality of the classical limit. In particular, one can check that the coefficients in (10.29), which are defined up to a global scaling factor, are

$$A = e^{\frac{i}{2}\alpha}(k^{-1} - \bar{\ell}_E), \quad B = -e^{\frac{i}{2}\alpha}(\bar{\ell}_E + k^{-1}), \quad C = e^{-\frac{i}{2}\alpha}(k^{-1} + \ell_E), \quad D = e^{-\frac{i}{2}\alpha}(\ell_E - k^{-1}), \quad (16.32)$$

¹⁵Note that the ratio λ_e/λ_c is the fine-structure constant $\alpha = e^2/\hbar c$.

where we used

$$\frac{w + i\bar{\ell}_E}{w - i\ell_E} = \frac{k^{-1} \tan kq + i\bar{\ell}_E}{k^{-1} \tan kq - i\ell_E} = \frac{(k^{-1} - \bar{\ell}_E)e^{2ikq} - \bar{\ell}_E - k^{-1}}{(k^{-1} + \ell_E)e^{2ikq} + \ell_E - k^{-1}}. \quad (16.33)$$

The limit (10.29) is a direct consequence of the fact that the classical limit of (16.31) is $\pm\sqrt{2mE}$.

A feature related to the above investigation concerns the analogue of de Broglie's wave-length

$$\lambda(q) = \frac{h}{|p|} = h|\phi|^2, \quad (16.34)$$

where h denotes the Planck constant. Let us denote by q_R a solution of the equation

$$\psi^D(q) = -\psi(q)\text{Im } \ell, \quad (16.35)$$

that determines the points where $\psi^D - i\ell\psi$ is purely imaginary. We have

$$\lambda(q_R) = 2\pi\psi^2(q_R)|W|^{-1}\text{Re } \ell, \quad (16.36)$$

and

$$|p(q_R)| = \frac{\hbar|W|}{\psi^2(q_R)\text{Re } \ell}. \quad (16.37)$$

In the case of the state $\mathcal{W}^0$, we have

$$\lambda^0(q^0) = 4\pi \frac{|q^0 - i\ell_0|^2}{\ell_0 + \bar{\ell}_0}, \quad (16.38)$$

that reaches its minimum precisely at $q^0 = q_R^0 = -\text{Im } \ell_0$

$$\lambda^0(-\text{Im } \ell_0) = 2\pi\text{Re } \ell_0. \quad (16.39)$$

Under the trivializing map all the solutions of (16.35) map to q_R^0 . In fact, by (15.13) and (16.35)

$$-\text{Im } \ell_0 = \frac{-(\ell_0 + \bar{\ell}_0)\text{Im } \ell + i\ell_0\bar{\ell} - i\bar{\ell}_0\ell}{\ell + \bar{\ell}}. \quad (16.40)$$

More generally, by (15.41)

$$-\text{Im } \ell_b = \frac{-(\ell_b + \bar{\ell}_b)\text{Im } \ell_a + i\ell_b\bar{\ell}_a - i\bar{\ell}_b\ell_a}{\ell_a + \bar{\ell}_a}, \quad (16.41)$$

which can be also seen as a consequence of (16.40) and of the pseudogroup property of v -maps. Note that since

$$p_0 dq^0 = pdq, \quad (16.42)$$

we have the transformation property

$$\lambda^0(q^0) = \left| \frac{\partial q^0}{\partial q} \right| \lambda(q), \quad (16.43)$$

which is equivalent to

$$\lambda^0(q^0) = \left| \frac{\phi^0(q^0)}{\phi(q)} \right|^2 \lambda(q), \quad (16.44)$$

where ϕ^0 is the function given in (14.65) in the case of $\mathcal{W}^0$.

We have seen that in considering the $\hbar \rightarrow 0$ and $E \rightarrow 0$ limits, one has to introduce some fundamental lengths. The structure of ℓ_E indicates a possible way in which fundamental constants and related interactions may arise. In particular, Eq.(16.29) suggests that in a suitable context these quantities may play the role of a natural cut-off. In this respect we observe that in the expression of ℓ_E both x_p and x_p^{-1} appears, a fact suggesting a possible ultraviolet-infrared duality. In this context we recall that the basic Möbius symmetry (15.36) is intrinsically related to the structure of the QSHJE and then of the trivializing map. One may consider the state $\mathcal{W}^0$ as a sort of vacuum state with all the other states connected by the trivializing map. Then, thanks to the pseudogroup property of v -maps, the Möbius symmetry of the state $\mathcal{W}^0$, reflects in symmetry properties of arbitrary states. In this respect we stress that the symmetry associated to the state $\mathcal{W}^0$ includes the inversion (15.38). In particular, observe that after the rescaling of the space coordinate

$$q_\beta^0 \longrightarrow X_\beta^0 = \frac{q_\beta^0}{|\ell_0|}, \quad (16.45)$$

we have that (15.38) precisely corresponds to the S -duality transformation

$$X_{(2n+1)\pi/2}^0 = -\frac{1}{X^0}, \quad (16.46)$$

which makes a correspondence between long and short distances.

We have seen that the ℓ_E 's naturally appear in the framework of the Möbius transformations whose origin traces back to the p - q duality we studied at the beginning of our investigation. Actually, p - q duality is connected to the ψ^D - ψ duality. In particular, while in considering the self-dual states we were forced to introduce the dimensional constants γ , γ_p and γ_q , in the case discussed above we introduced the dimensional constants λ_p , λ_c and λ_e . As for p - q duality, in which the dimensionality of the constants is related to the Möbius symmetry and to the different dimensional properties of p and q , in the case of ψ^D and ψ we have that any linear combination of ψ^D and ψ corresponds to a Möbius transformation of $w = \psi^D/\psi$. On the other hand, a linear combination between ψ^D and ψ forces us, as in the case of p and q , to introduce dimensional constants. The origin of this is, once again, the existence of the self-dual state, which actually coincides with the state $\mathcal{W}^0$. In fact, since for the state $\mathcal{W}^0$ the functions ψ^D and ψ must solve the equation $\partial_q^2 \psi^D = 0 = \partial_q^2 \psi$, we necessarily have to introduce a constant length in considering linear combinations of $\psi^D = q$ and $\psi = cnst$. Since these solutions correspond to $\mathcal{W}^0 \equiv 0$, we do not have any constant length provided by the problem. This leads considering fundamental constants. This is the basic reason of the appearance of λ_p , λ_c and λ_e . Since these constants appear in ℓ , we have that in our formulation of QM there are hidden variables depending on the Planck length.

16.2 Time, energy and trajectories

We now consider an aspect concerning Floyd's proposal of using Jacobi's theorem to define time parameterization. The point is that in considering $\partial_E \mathcal{S}_0$, one has to understand whether the E dependence of ℓ , as in the case of ℓ_E for the state with $\mathcal{W} = -E$, should be derived. In other words, one should fix which terms in the expression of $\mathcal{S}_0$ one has to consider as explicit. For example, in the case of the free particle of energy E , one can absorb the multiplicative k^{-1} factor in $w = k^{-1} \tan kq$ just by a rescaling of ℓ_E in the expression of the reduced action that we denote by $\mathcal{S}_0^E$. If in evaluating the partial derivative $\partial_E \mathcal{S}_0^E$ one considers the dependence of ℓ_E on E as the unique implicit one, then the value of $\partial_E \mathcal{S}_0^E$ is not well-defined: a common E -dependent rescaling of $w + i\ell_E$ and $w - i\ell_E$ in (14.57) would change $\partial_E w$, and therefore $t - t_0 = \partial_E \mathcal{S}_0^E$, leaving $\mathcal{S}_0^E$ invariant. In order to better understand these aspects, let us consider time parameterization in the case of the state $\mathcal{W}^0$. Since this state corresponds to the $E \rightarrow 0$ limit of the state $\mathcal{W} = -E$, we have

$$t - t_0 = \lim_{E \rightarrow 0} \frac{\partial \mathcal{S}_0^E}{\partial E}. \quad (16.47)$$

On the other hand, for any fixed ℓ_0 any different choice of the functions $\alpha(x_p^{-1})$ and $\beta(x_p)$ corresponds by construction to a set of ℓ_E 's with the common property of having the same limit ℓ_0 . This means that for any choice of the functions $\alpha(x_p^{-1})$ and $\beta(x_p)$ one obtains always $\mathcal{S}_0^0$ with the same ℓ_0 . Hence, different choices of $\alpha(x_p^{-1})$ and $\beta(x_p)$ will correspond to the same Möbius state in the $E \rightarrow 0$ limit. This implies that the limit cannot depend on $\alpha(x_p^{-1})$ and $\beta(x_p)$. What remains to understand is whether the k^{-1} factor in $w = k^{-1} \tan kq$ and in $\ell_E = k^{-1} e^{-\alpha(x_p^{-1})} + e^{-\beta(x_p)} \ell_0$, should be derived. A natural possibility that would allow a non ambiguous definition of time parameterization, is to assume that all the terms depending on E that can be absorbed in a redefinition of ℓ should not be considered in evaluating $\partial_E \mathcal{S}_0$. In other words, the only E terms in $\mathcal{S}_0$ that should be considered in evaluating $\partial_E \mathcal{S}_0$ are those for which the transformation $E \rightarrow E' \neq E$ does not correspond to a Möbius transformation of $e^{\frac{2i}{\hbar} \mathcal{S}_0}$, that is such that $\mathcal{W} \rightarrow \mathcal{W}' \neq \mathcal{W}$. Let us consider again the case of the free particle. We have $w = k^{-1} \tan kq$. While changing E in E' in the k^{-1} factor would give $\{k'^{-1} \tan kq, q\} = \{w, q\} = -4mE/\hbar^2$, where $k' = \sqrt{2mE'}/\hbar$, in the case in which we change the E appearing in $\tan kq$ we have $\{k^{-1} \tan k'q, q\} = -4mE'/\hbar^2 \neq -4mE/\hbar^2$. Hence, the k^{-1} term in w should not be derived. The constant corresponding to the initial conditions for $\mathcal{S}_0^E$ are α and $k\ell_E$ and the only term in $\mathcal{S}_0^E$ to be derived with respect to E is $kw = \tan kq$. It follows that the equation for the trajectories of the free particle of energy E is

$$t - t_0 = \frac{\partial \mathcal{S}_0^E}{\partial E} = \frac{k(\ell_E + \bar{\ell}_E)}{2|\sin(kq) - ik\ell_E \cos(kq)|^2} \sqrt{\frac{m}{2E}} q. \quad (16.48)$$

Note that since by (16.7)(16.9) and (16.20)

$$\lim_{\hbar \rightarrow 0} k\ell_E = 1, \quad (16.49)$$

we have that the trajectories (16.48) precisely collapse to the classical one in the $\hbar \rightarrow 0$ limit

$$t - t_0 = \sqrt{\frac{m}{2E}} q. \quad (16.50)$$

Hence, the QSHJE correctly leads to trajectories whose classical limit arises in a natural way. Note that the quantum trajectory (16.48) can be written as the multiplicative correction of the classical one

$$t - t_0 = F_{qu}(q) \sqrt{\frac{m}{2E}} q, \quad (16.51)$$

where

$$F_{qu}(q) = \frac{k(\ell_E + \bar{\ell}_E)}{2|\sin(kq) - ik\ell_E \cos(kq)|^2}. \quad (16.52)$$

It is easy to see that the right hand side of (16.48) diverges in the $E \rightarrow 0$ limit. This means that the state $\mathcal{W}^0$ corresponds, as in the classical limit, to a particle at rest, that is

$$\dot{q}^0 = 0. \quad (16.53)$$

Therefore, in the formulation derived from the EP a free particle with vanishing energy is at rest, as in the classical case. However, while in CM one has $\partial_q \mathcal{S}_0^{cl} = m\dot{q}$, in QM we have $\partial_q \mathcal{S}_0 \neq m\dot{q}$. In particular, while for the state $\mathcal{W}^0$ we have $\dot{q} = 0$, the conjugate momentum is non-vanishing. This can be considered as the effect of the quantum mass field. Let us denote by m_{Q_E} the quantum mass field corresponding to the free particle of energy E . It follows from (14.6)(15.56) and (16.53) that

$$\lim_{E \rightarrow 0} m_{Q_E} = \pm\infty. \quad (16.54)$$

Hence, the fact that $p_0 \neq 0$ is due to the divergence of $\partial_E Q_E$ in the $E \rightarrow 0$ limit.

While the state $\mathcal{W}^0$ corresponds, as in the classical limit, to a particle at rest, in the $\mathcal{W} = -E$ case the velocity is not a constant. In fact, deriving (16.48) with respect to q we obtain

$$\dot{q} = \frac{2|\sin(kq) - ik\ell_E \cos(kq)|^4}{k(\ell_E + \bar{\ell}_E) \left\{ |\sin(kq) - ik\ell_E \cos(kq)|^2 + kq[|k\ell_E|^2 - 1] \sin(2kq) + ik(\ell_E - \bar{\ell}_E) \cos(2kq) \right\}} \sqrt{\frac{2E}{m}}, \quad (16.55)$$

which is a constant in the classical limit only

$$\lim_{\hbar \rightarrow 0} \dot{q} = \sqrt{2E/m}. \quad (16.56)$$

Therefore, for the trajectory (16.48) we have $\mathcal{S}_0^E \neq \sqrt{2mE}q$ unless $k\ell_E = 1$, which holds in the classical limit only. Now recall that in discussing p - q duality, we observed that since this is based on the existence of the Legendre transform of $\mathcal{S}_0$, it would break down when either $\mathcal{S}_0 = cst$ or $\mathcal{S}_0 \propto q + cst$. Hence, requiring a formulation with manifest p - q duality for all the possible states forces one to discard CM. On the other hand, we derived the QSHJE (9.44) from the EP without assuming any p - q duality. Then we noticed that, due to the $\{\mathcal{S}_0, q\}$ term, the QSHJE could not be defined for $\mathcal{S}_0 = cst$. Hence, a condition for the existence of p - q duality was a simple consequence of the implementation of the EP. It remains to understand whether also the case $\mathcal{S}_0^E \propto q + cst$, for which the Legendre transform of $\mathcal{S}_0$ degenerates, could be discarded for reasons related to the implementation of the EP. This would give an even deeper relationship between the EP and p - q duality. We have seen

above that this is in fact the case: this came out as a consistency condition on the limits. In particular, the solution $\mathcal{S}_0^E = \sqrt{2mE}q$, would imply

$$\mathcal{S}_0^0 = \lim_{E \rightarrow 0} \mathcal{S}_0^E = 0, \quad (16.57)$$

that we already discarded. Therefore, the EP guarantees p - q duality.

17 Equivalence Principle, tunnelling and quantized spectra

In this section we show how tunnelling arises in the context of the QSHJE. We also show, following [2], the basic fact that when $\lim_{q \rightarrow \pm\infty} \mathcal{W} > 0$ the QSHJE is defined only in the case in which the corresponding SE has square summable solutions. This implies the standard result on the quantized energy spectra without making use of any axiomatic interpretation of the wave-function. We will also discuss the structure of the spaces $\mathcal{H}$ and $\mathcal{K}$ of the admissible $\mathcal{W}$'s and $\mathcal{S}_0$'s respectively.

17.1 Tunnelling

It is well-known that it may happen that there are space regions which are forbidden to classical trajectories. From the CSHJE one has

$$p = \pm \sqrt{2m(E - V)}, \quad (17.1)$$

so that the classically forbidden region is $\Omega = \{q \in \mathbb{R} | V(q) - E > 0\}$. The situation is completely different in the case of the QSHJE where

$$p = \pm \sqrt{2m(-V - Q + E)}. \quad (17.2)$$

Hence, due to the Q term, we have that even if $q \in \Omega$, p may be real. In fact, since $p = \epsilon|\phi|^{-2}$, we have $p \in \mathbb{R}$, $\forall q \in \mathbb{R}$. As we will see the exception arises just for the case of the infinitely deep potential well. In other words, we have the tunnel effect for the trajectories described by the QSHJE. In the usual formulation, tunnelling is essentially a consequence of the axiomatic interpretation of the wave-function in terms of probability amplitude. Note that we considered reality of p . However, to have the tunnel effect one should check that $\dot{q}$, like p , is a real quantity for any q . It can be seen, for example by (15.53), that this is in fact the case. This can be also seen by noticing that

$$t - t_0 = \frac{\partial \mathcal{S}_0}{\partial E} = \frac{W_E}{W} p, \quad W_E \equiv \psi^D \partial_E \psi - \psi \partial_E \psi^D, \quad (17.3)$$

implies $t \in \mathbb{R}$, $\forall q \in \mathbb{R}$. The reason underlying the tunnel effect resides in the fact that it is always possible to choose a pair of linearly independent solutions of the SE which are real for any $q \in \mathbb{R}$. Deriving (17.3) with respect to q we obtain the expression of the velocity

$$\dot{q} = \frac{W}{\partial_q(pW_E)}. \quad (17.4)$$

Hence, there are no forbidden regions associated to the QSHJE. We note that this can be also seen as a direct consequence of the EP. In fact, since the EP implies that (2.37) is a local self-homeomorphism of $\hat{\mathbb{R}}$, even the existence of only one $\mathcal{S}_0$, $\forall q \in \mathbb{R}$, implies that any reduced action will be real for any real value of the coordinate. On the other hand, if the reduced actions take real values for all $q \in \mathbb{R}$, then $p = \partial_q \mathcal{S}_0 \in \mathbb{R}$, $\forall q \in \mathbb{R}$. This implies that $t \in \mathbb{R}$, $\forall q \in \mathbb{R}$. The exception arises for the infinitely deep potential well. We will see that this situation will come as a particular limiting case.

17.2 Quantized spectra from the Equivalence Principle

We saw that the EP implied the QSHJE (9.44). However, although this equation implies the SE, we saw that there are aspects concerning the canonical variables which arise in considering the QSHJE rather than the SE. In this respect a natural question is whether the basic facts of QM also arise in our formulation. A basic point concerns a property of many physical systems such as energy quantization. This is a matter of fact beyond any interpretational aspect of QM. Then, as we used the EP to get the QSHJE, it is important to understand how energy quantization arises in our approach. According to the EP, the QSHJE contains all the possible information on a given system. Then, the QSHJE itself should be sufficient to recover the energy quantization including its structure. In the usual approach the quantization of the spectrum arises from the basic condition that in the case in which $\lim_{q \rightarrow \pm\infty} \mathcal{W} > 0$, the wave-function should vanish at infinity. Once the possible solutions are selected, one also imposes the continuity conditions whose role in determining the possible spectrum is particularly transparent in the case of discontinuous potentials. For example, in the case of the potential well, besides the restriction on the spectrum due to the $L^2(\mathbb{R})$ condition for the wave-function (a consequence of the probabilistic interpretation of the wave-function), the spectrum is further restricted by the smoothly joining conditions. Since the SE contains the term $\partial_q^2 \psi$, the continuity conditions correspond to an existence condition for this equation. On the other hand, also in this case, the physical reason underlying this request is the interpretation of the wave-function in terms of probability amplitude. Actually, strictly speaking, the continuity conditions come from the continuity of the probability density $\rho = |\psi|^2$. This density should also satisfy the continuity equation $\partial_t \rho + \partial_q j = 0$, where $j = i\hbar(\psi \partial_q \bar{\psi} - \bar{\psi} \partial_q \psi)/2m$. Since for stationary states $\partial_t \rho = 0$, it follows that in this case $j = \text{const}$. Therefore, in the usual formulation, it is just the interpretation of the wave-function in terms of probability amplitude, with the consequent meaning of ρ and j , which provides the physical motivation for imposing the continuity of the wave-function and of its first derivative.

Now observe that in our formulation the continuity conditions arise from the QSHJE. In fact, (15.7) implies continuity of ψ^D , ψ , with $\partial_q \psi^D$ and $\partial_q \psi$ differentiable, that is

$$\textit{Equivalence Principle} \longrightarrow \textit{continuity of } (\psi^D, \psi), \text{ and } (\partial_q \psi^D, \partial_q \psi) \text{ differentiable.} \quad (17.5)$$

Let us now consider the condition about the existence of an $L^2(\mathbb{R})$ solution in the case in which $\lim_{q \rightarrow \pm\infty} \mathcal{W} > 0$. Let us first note that in the interesting paper [15], the possible values of E for which the SE has no $L^2(\mathbb{R})$ solution were a priori discarded. Apparently, in considering the QSHJE there are no reasons to make this assumption. Thus, even if in [15] the axiomatic interpretation of the wave-function is refused, and a basic trajectory interpretation of QM based on the QSHJE is provided, we

note that imposing the $L^2(\mathbb{R})$ condition appears as an unjustified assumption. Actually, this is what essentially happens also in Bohm's theory, where the $L^2(\mathbb{R})$ condition is essentially assumed. To a great extent this seems to happen also in the stochastic formulation of QM. On the other hand, in our approach we have a basic natural principle which motivates both the QSHJE and then the continuity condition. It is therefore of basic importance to investigate whether the existence of an $L^2(\mathbb{R})$ solution is also a consequence of the EP. We now show that this is in fact the case!

In the following we will derive a result concerning the energy spectra. In this context we will see that if $V(q) > E$, $\forall q \in \mathbb{R}$, then there are no solutions such that the ratio of two real linearly independent solutions of the SE corresponds to a local self-homeomorphism of $\hat{\mathbb{R}}$. The fact that this is an unphysical situation can be also seen from the fact that the case $V > E$, $\forall q \in \mathbb{R}$, has no classical limit. Therefore, if $V > E$ both at $-\infty$ and $+\infty$, a physical situation requires that there are at least two points in which $V - E = 0$. More generally, if the potential is not continuous, we should have at least two turning points for which $V(q) - E$ changes sign. Let us denote by q_- (q_+) the lowest (highest) turning point. We will prove the following basic fact

If

$$V(q) - E \geq \begin{cases} P_-^2 > 0, & q < q_-, \\ P_+^2 > 0, & q > q_+, \end{cases} \quad (17.6)$$

then w is a local self-homeomorphism of $\hat{\mathbb{R}}$ if and only if the corresponding SE has an $L^2(\mathbb{R})$ solution.

Note that by (17.6) we have

$$\int_{q_-}^{-\infty} dx \kappa(x) = -\infty, \quad \int_{q_+}^{+\infty} dx \kappa(x) = +\infty, \quad (17.7)$$

where $\kappa = \sqrt{2m(V - E)}/\hbar$. Before going further, let us stress that what we actually need to prove is that in the case (17.6), the joining condition (15.8) requires that the corresponding SE has an $L^2(\mathbb{R})$ solution. Observe that while (15.7), which however follows from the EP, can be recognized as the standard condition (17.5), the other condition (15.8), which still follows from the existence of the QSHJE, and therefore from the EP, is not directly recognized in the standard formulation. Since this leads to energy quantization, while in the usual approach one needs one more assumption, we see that there is a quite fundamental difference between the QSHJE and the SE. We stress that (15.7) and (15.8) guarantee that w is a local self-homeomorphism of $\hat{\mathbb{R}}$.

Let us first show that the request that the corresponding SE has an $L^2(\mathbb{R})$ solution is a sufficient condition for w to satisfy (15.8). Let $\psi \in L^2(\mathbb{R})$ and denote by ψ^D a linearly independent solution. As we will see, the fact that $\psi^D \not\propto \psi$ implies that if $\psi \in L^2(\mathbb{R})$, then $\psi^D \notin L^2(\mathbb{R})$. In particular, ψ^D is divergent both at $q = -\infty$ and $q = +\infty$. Let us consider the real ratio

$$w = \frac{A\psi^D + B\psi}{C\psi^D + D\psi}, \quad (17.8)$$

where $AD - BC \neq 0$. Since $\psi \in L^2(\mathbb{R})$, we have

$$\lim_{q \rightarrow \pm\infty} w = \lim_{q \rightarrow \pm\infty} \frac{A\psi^D + B\psi}{C\psi^D + D\psi} = \frac{A}{C}, \quad (17.9)$$

that is $w(-\infty) = w(+\infty)$. In the case in which $C = 0$ we have

$$\lim_{q \rightarrow \pm\infty} w = \lim_{q \rightarrow \pm\infty} \frac{A\psi^D}{D\psi} = \pm\epsilon \cdot \infty, \quad (17.10)$$

where $\epsilon = \pm 1$. The fact that $\lim_{q \rightarrow \pm\infty} A\psi^D/D\psi$ diverges follows from the mentioned properties of ψ^D and ψ . It remains to check that if $\lim_{q \rightarrow -\infty} A\psi^D/D\psi = -\infty$, then $\lim_{q \rightarrow +\infty} A\psi^D/D\psi = +\infty$, and vice versa. This can be seen by observing that

$$\psi^D(q) = c\psi(q) \int_{q_0}^q dx \psi^{-2}(x) + d\psi(q), \quad (17.11)$$

$c \in \mathbb{R} \setminus \{0\}$, $d \in \mathbb{R}$. Since $\psi \in L^2(\mathbb{R})$ we have $\psi^{-1} \notin L^2(\mathbb{R})$ and $\int_{q_0}^{+\infty} dx \psi^{-2}(x) = +\infty$, $\int_{q_0}^{-\infty} dx \psi^{-2}(x) = -\infty$, so that $\psi^D(-\infty)/\psi(-\infty) = -\epsilon \cdot \infty = -\psi^D(+\infty)/\psi(+\infty)$, where $\epsilon = \text{sgn } c$.

We now show that the existence of an $L^2(\mathbb{R})$ solution of the SE is a necessary condition to satisfy the joining condition (15.8). We give two different proofs of this, one is based on the WKB approximation while the other one uses Wronskian arguments. In the WKB approximation, we have

$$\psi = \frac{A_-}{\sqrt{\kappa}} e^{-\int_{q_-}^q dx \kappa(x)} + \frac{B_-}{\sqrt{\kappa}} e^{\int_{q_-}^q dx \kappa(x)}, \quad q \ll q_-, \quad (17.12)$$

and

$$\psi = \frac{A_+}{\sqrt{\kappa}} e^{-\int_{q_+}^q dx \kappa(x)} + \frac{B_+}{\sqrt{\kappa}} e^{\int_{q_+}^q dx \kappa(x)}, \quad q \gg q_+. \quad (17.13)$$

In the same approximation, a linearly independent solution has the form

$$\psi^D = \frac{A_-^D}{\sqrt{\kappa}} e^{-\int_{q_-}^q dx \kappa(x)} + \frac{B_-^D}{\kappa} e^{\int_{q_-}^q dx \kappa(x)}, \quad q \ll q_-. \quad (17.14)$$

Similarly, in the $q \gg q_+$ region we have

$$\psi^D = \frac{A_+^D}{\sqrt{\kappa}} e^{-\int_{q_+}^q dx \kappa(x)} + \frac{B_+^D}{\sqrt{\kappa}} e^{\int_{q_+}^q dx \kappa(x)}, \quad q \gg q_+. \quad (17.15)$$

Note that (17.12)–(17.13) are derived by solving the differential equations corresponding to the WKB approximation for $q \ll q_-$ and $q \gg q_+$, so that the coefficients of $\kappa^{-1/2} \exp \pm \int_{q_{\pm}}^q dx \kappa(x)$, *e.g.* A_- and B_- in (17.12), cannot be simultaneously vanishing. In particular, the fact that $\psi^D \not\propto \psi$ yields

$$A_- B_-^D - A_-^D B_- \neq 0, \quad A_+ B_+^D - A_+^D B_+ \neq 0. \quad (17.16)$$

Let us now consider the case in which, for a given E satisfying (17.6), any solution of the corresponding SE diverges at least at one of the two spatial infinities, that is

$$\lim_{q \rightarrow +\infty} (|\psi(-q)| + |\psi(q)|) = +\infty. \quad (17.17)$$

This implies that there is a solution diverging both at $q = -\infty$ and $q = +\infty$. In fact, if two solutions ψ_1 and ψ_2 satisfy $\psi_1(-\infty) = \pm\infty$, $\psi_1(+\infty) \neq \pm\infty$ and $\psi_2(-\infty) \neq \pm\infty$, $\psi_2(+\infty) = \pm\infty$, then $\psi_1 + \psi_2$

diverges at $\pm\infty$. On the other hand, (17.16) rules out the case in which all the solutions in their WKB approximation are divergent only at one of the two spatial infinities, say $-\infty$. Since, in the case (17.6), a solution which diverges in the WKB approximation is itself divergent (and vice versa), we have that in the case (17.6), the fact that all the solutions of the SE diverge only at one of the two spatial infinities cannot occur.

Let us denote by ψ a solution which is divergent both at $-\infty$ and $+\infty$. In the WKB approximation this means that both A_- and B_+ are non-vanishing, so that

$$\psi \underset{q \rightarrow -\infty}{\sim} \frac{A_-}{\sqrt{\kappa}} e^{-\int_{q_-}^q dx \kappa}, \quad \psi \underset{q \rightarrow +\infty}{\sim} \frac{B_+}{\sqrt{\kappa}} e^{\int_{q_+}^q dx \kappa}. \quad (17.18)$$

The asymptotic behavior of the ratio ψ^D/ψ is given by

$$\lim_{q \rightarrow -\infty} \frac{\psi^D}{\psi} = \frac{A_-^D}{A_-}, \quad \lim_{q \rightarrow +\infty} \frac{\psi^D}{\psi} = \frac{B_+^D}{B_+}. \quad (17.19)$$

Note that since in the case at hand any divergent solution also diverges in the WKB approximation, we have that (17.17) rules out the case $A_-^D = B_+^D = 0$. Let us then suppose that either $A_-^D = 0$ or $B_+^D = 0$. If $A_-^D = 0$, then $w(-\infty) = 0 \neq w(+\infty)$. Similarly, if $B_+^D = 0$, then $w(+\infty) = 0 \neq w(-\infty)$. Hence, in this case w , and therefore the trivializing map, cannot satisfy (15.8). On the other hand, also in the case in which both A_-^D and B_+^D are non-vanishing, w cannot satisfy Eq.(15.8). For, if $A_-^D/A_- = B_+^D/B_+$, then

$$\phi = \psi - \frac{A_-}{A_-^D} \psi^D = \psi - \frac{B_+}{B_+^D} \psi^D, \quad (17.20)$$

would be a solution of the SE whose WKB approximation has the form

$$\phi = \frac{B_-}{\sqrt{\kappa}} e^{\int_{q_-}^q dx \kappa(x)}, \quad q \ll q_-, \quad (17.21)$$

and

$$\phi = \frac{A_+}{\sqrt{\kappa}} e^{-\int_{q_+}^q dx \kappa(x)}, \quad q \gg q_+. \quad (17.22)$$

Hence, if $A_-^D/A_- = B_+^D/B_+$, then there is a solution whose WKB approximation vanishes both at $-\infty$ and $+\infty$. On the other hand, we are considering the values of E satisfying Eq.(17.6) and for which any solution of the SE has the property (17.17). This implies that no solutions can vanish both at $-\infty$ and $+\infty$ in the WKB approximation. Hence

$$\frac{A_-^D}{A_-} \neq \frac{B_+^D}{B_+}, \quad (17.23)$$

so that $w(-\infty) \neq w(+\infty)$. We also note that not even the case $w(-\infty) = \pm\infty = -w(+\infty)$ can occur, as this would imply that $A_- = B_+ = 0$, which in turn would imply, against the hypothesis, that there are solutions vanishing at $q = \pm\infty$. Hence, if for a given E satisfying (17.6), any solution of the corresponding SE diverges at least at one of the two spatial infinities, we have that the trivializing

map has a discontinuity at $q = \pm\infty$. As a consequence, the EP cannot be implemented in this case so that this value E cannot belong to the physical spectrum.

Therefore, the physical values of E satisfying (17.6) are those for which there are solutions which are divergent neither at $-\infty$ nor at $+\infty$. On the other hand, from the WKB approximation and (17.6), it follows that the non-divergent solutions must vanish both at $-\infty$ and $+\infty$. It follows that the only energy levels satisfying the property (17.6), which are compatible with the EP, are those for which there exists the solution vanishing both at $\pm\infty$. On the other hand, solutions vanishing as $\kappa^{-1/2} \exp \int_{q_-}^q dx \kappa$ at $-\infty$ and $\kappa^{-1/2} \exp - \int_{q_+}^q dx \kappa$ at $+\infty$, with $P_{\pm}^2 > 0$, cannot contribute with an infinite value to $\int_{-\infty}^{+\infty} dx \psi^2(x)$. The reason is that existence of the QSHJE requires that $\{e^{\frac{2i}{\hbar} S_0}, q\}$ be defined and this, in turn, implies that any solution of the SE must be continuous. On the other hand, since ψ is continuous, and therefore finite also at finite values of q , we have $\int_{q_a}^{q_b} dx \psi^2(x) < +\infty$ for all finite q_a and q_b . In other words, the only possibility for a continuous function to have a divergent value of $\int_{-\infty}^{+\infty} dx \psi^2(x)$ comes from its behavior at $\pm\infty$. Therefore, since the implementation of the EP in the case (17.6) requires that the corresponding E should admit a solution with the behavior

$$\psi_{q \rightarrow -\infty} \sim \frac{A_-}{\sqrt{\kappa_-}} e^{\int_{q_-}^q dx \kappa}, \quad \psi_{q \rightarrow +\infty} \sim \frac{B_+}{\sqrt{\kappa_+}} e^{-\int_{q_+}^q dx \kappa}, \quad (17.24)$$

we have the following basic fact

The values of E satisfying

$$V(q) - E \geq \begin{cases} P_-^2 > 0, & q < q_-, \\ P_+^2 > 0, & q > q_+, \end{cases} \quad (17.25)$$

are physically admissible if and only if the corresponding SE has an $L^2(\mathbb{R})$ solution.

We now give another proof of the fact that if $\mathcal{W}$ is of the type (17.25), then the corresponding SE must have an $L^2(\mathbb{R})$ solution in order to satisfy (15.8). In particular, we will show that this is a necessary condition. That this is sufficient has been already proved above.

Let us start by observing that Wronskian arguments, which can be found in Messiah's book [40], imply that if $V(q) - E \geq P_+^2 > 0$, $q > q_+$, then as $q \rightarrow +\infty$, we have ($P_+ > 0$)

- There is a solution of the SE that vanishes at least as $e^{-P_+ q}$.
- Any other linearly independent solution diverges at least as $e^{P_+ q}$.

Similarly, if $V(q) - E \geq P_-^2 > 0$, $q < q_-$, then as $q \rightarrow -\infty$, we have ($P_- > 0$)

- There is a solution of the SE that vanishes at least as $e^{P_- q}$.
- Any other linearly independent solution diverges at least as $e^{-P_- q}$.

These properties imply that if there is a solution of the SE in $L^2(\mathbb{R})$, then any solution is either in $L^2(\mathbb{R})$ or diverges both at $-\infty$ and $+\infty$. Let us show that the possibility that a solution vanishes only at one of the two spatial infinities is ruled out. Suppose that, besides the $L^2(\mathbb{R})$ solution, which we

denote by ψ_1 , there is a solution ψ_2 which is divergent only at $+\infty$. On the other hand, the above properties show that there exists also a solution ψ_3 which is divergent at $-\infty$. Since the number of linearly independent solutions of the SE is two, we have $\psi_3 = A\psi_1 + B\psi_2$. However, since ψ_1 vanishes both at $-\infty$ and $+\infty$, we see that $\psi_3 = A\psi_1 + B\psi_2$ can be satisfied only if ψ_2 and ψ_3 are divergent both at $-\infty$ and $+\infty$. This fact and the above properties imply that

If the SE has an $L^2(\mathbb{R})$ solution, then any solution has two possible asymptotics

- Vanishes both at $-\infty$ and $+\infty$ at least as e^{P-q} and e^{-P+q} respectively.
- Diverges both at $-\infty$ and $+\infty$ at least as e^{-P-q} and e^{P+q} respectively.

Similarly, we have

If the SE does not admit an $L^2(\mathbb{R})$ solution, then any solution has three possible asymptotics

- Diverges both at $-\infty$ and $+\infty$ at least as e^{-P-q} and e^{P+q} respectively.
- Diverges at $-\infty$ at least as e^{-P-q} and vanishes at $+\infty$ at least as e^{-P+q} .
- Vanishes at $-\infty$ at least as e^{P-q} and diverges at $+\infty$ at least as e^{P+q} .

Let us consider the ratio $w = \psi^D/\psi$ in the latter case. Since any different choice of linearly independent solutions of the SE corresponds to a Möbius transformation of w , we can choose¹⁶

$$\psi^D_{q \rightarrow -\infty} \sim a_- e^{P-q}, \quad \psi^D_{q \rightarrow +\infty} \sim a_+ e^{P+q}, \quad (17.26)$$

and

$$\psi_{q \rightarrow -\infty} \sim b_- e^{-P-q}, \quad \psi_{q \rightarrow +\infty} \sim b_+ e^{-P+q}. \quad (17.27)$$

Their ratio has the asymptotics

$$\frac{\psi^D}{\psi}_{q \rightarrow -\infty} \sim c_- e^{2P-q} \rightarrow 0, \quad \frac{\psi^D}{\psi}_{q \rightarrow +\infty} \sim c_+ e^{2P+q} \rightarrow \pm\infty, \quad (17.28)$$

so that w cannot satisfy Eq.(15.8). This concludes the alternative proof of the fact that, in the case (17.25), the existence of the $L^2(\mathbb{R})$ solution is a necessary condition in order (15.8) be satisfied. The fact that this is a sufficient condition has been proved previously in deriving Eq.(17.9).

The above results imply that the usual quantized spectrum arises as a consequence of the EP. As examples we will consider the potential well and the simple and double harmonic oscillators.

Let us note that we are considering real solutions of the SE. Thus, apparently, in requiring the existence of an $L^2(\mathbb{R})$ solution, one should specify the existence of a real $L^2(\mathbb{R})$ solution. However, if there is an $L^2(\mathbb{R})$ solution ψ , this is unique up to a constant, and since also $\bar{\psi} \in L^2(\mathbb{R})$ solves the SE, we have that an $L^2(\mathbb{R})$ solution of the SE is real up to a phase.

¹⁶Here by $\sim$ we mean that ψ^D and ψ either diverge or vanish “at least as”.

17.3 The index of the trivializing map

Let us consider some further properties of the trivializing map. Let n be the index $I[q^0]$ of the covering associated to the trivializing map (15.10). This is the number of times q^0 spans $\hat{\mathbb{R}}$ while q spans $\hat{\mathbb{R}}$. Since q^0 and w are related by a Möbius transformation, we have

$$I[q^0] = I[w]. \quad (17.29)$$

Another property of the trivializing map is that its index depends on $\mathcal{W}$ but not on the specific Möbius state. Note that since p does not vanish for finite values of w , it follows that $I[q^0]$ coincides with the number of zeroes of w . This fact may be also understood by recalling the Sturm theorem (see [69][39][70] for a simple geometrical interpretation), stating that given two linearly independent solutions ψ^D and ψ of $\psi'' = K\psi$, between any two zeroes of ψ^D there is one zero of ψ .¹⁷ This theorem, and the condition that the values of E satisfying (17.25) should correspond to a SE having an $L^2(\mathbb{R})$ solution, guarantees local homeomorphicity of the trivializing map. It remains to understand the case in which $V(q) > E, \forall q \in \mathbb{R}$. We already noticed that, since there is no classical limit in this case, these solutions are not admissible ones. This also follows, as it should, from the fact that these solutions do not satisfy (15.8). To see this, it is sufficient to note that if ψ decreases as $q \rightarrow -\infty$, then by $\psi''/\psi = 4m\mathcal{W}/\hbar^2 > 0, \forall q \in \mathbb{R}$, it follows that ψ is always convex, $\psi \notin L^2(\mathbb{R})$. Therefore, the absence of turning points does not modify the essence of the conclusions in the previous subsection and if $V(q) > E, \forall q \in \mathbb{R}$, then (15.8) cannot be satisfied. We refer to (1.48) for an explicit example.

17.4 Equivalence Principle and admissible potentials

We can now better define the spaces $\mathcal{H}$ and $\mathcal{K}$ that we introduced as the spaces of all possible $\mathcal{W}$ and $\mathcal{S}_0$ respectively. We introduced these spaces and then arrived to the conclusions that the implementation of the EP univocally implied the QSHJE (9.44). The appearance of $\{\mathcal{S}_0, q\}$ then implied that the trivializing map must be a local self-homeomorphism of $\hat{\mathbb{R}}$. This means that the possible $\mathcal{W}$'s which are compatible with the EP are all the functions which are $-\hbar^2/4m$ times the Schwarzian derivative of a local self-homeomorphism of $\hat{\mathbb{R}}$. This is nothing but a consequence of the cocycle condition (1.20) and therefore of the EP. Therefore, we have

The space $\mathcal{H}$ of admissible $\mathcal{W}$ states consists of the functions of the form

$$\mathcal{W} = -\frac{\hbar^2}{4m}\{w, q\}, \quad (17.30)$$

where w is an arbitrary local homeomorphism of $\hat{\mathbb{R}}$ into itself.

Similarly, we have

¹⁷This is another manifestation of ψ^D - ψ duality. In this context we stress that, while in the usual approach the wave-function plays the central role, our description contains both ψ^D and ψ .

The space $\mathcal{K}$ of admissible reduced actions $\mathcal{S}_0$, consists of the functions of the form

$$\mathcal{S}_0(q) = \frac{\hbar}{2i} \ln \left(\frac{w(q) + i\bar{\ell}}{w(q) - i\ell} \right) + \frac{\alpha\hbar}{2}, \quad (17.31)$$

where w is an arbitrary local homeomorphism of $\hat{\mathbb{R}}$ into itself, and¹⁸ $\alpha \in \mathbb{R}$, $\text{Re } \ell \neq 0$.

18 The potential well and the harmonic oscillators

In this section we will consider some relevant examples of the general fact we proved in the previous section. In particular, we will consider the cases of the potential well and of the simple and double harmonic oscillators. We will explicitly see that in the case one considers the energy values for which the corresponding SE has not $L^2(\mathbb{R})$ solutions, the ratio ψ^D/ψ has a discontinuity at spatial infinity.

18.1 The potential well

We now show how the quantized spectrum and its structure arise in the case of the potential well

$$V(q) = \begin{cases} 0, & |q| \leq L, \\ V_0, & |q| > L. \end{cases} \quad (18.1)$$

According to (14.29), in order to find the expression for the conjugate momentum, we have to find two linearly independent solutions of the SE which should satisfy the continuity conditions. Let us set

$$k = \frac{\sqrt{2mE}}{\hbar}, \quad \kappa = \frac{\sqrt{2m(V_0 - E)}}{\hbar}. \quad (18.2)$$

Since $V(q)$ is an even function, we can choose solutions of definite parity. For $|q| \leq L$ we can choose either $\psi_1^1 = \cos(kq)$ or $\psi_2^1 = \sin(kq)$. For $q > L$ we can choose either $\psi_1^2 = Ae^{-\kappa q}$ or $\psi_2^2 = Be^{\kappa q}$ (or any their linear combination). Parity of $V(q)$ fixes the solutions for $q < -L$. In general choosing ψ_i^1 and $A\psi_1^2 + B\psi_2^2$ will not give quantized spectra. Before considering this general situation, we consider the four cases given by the joining conditions $\psi_i^1 = \psi_j^2$, $\partial_q \psi_i^1 = \partial_q \psi_j^2$ at $q = L$. We will denote such solutions by (i, j) . We will see that the cases $(1, 2)$ and $(2, 2)$ correspond to a trivializing map which is discontinuous at $q = \pm\infty$ so that, according to the EP, these solutions are not physical and must be discarded. Let us first consider the $(1, 1)$ case.

$$\underline{i = 1, j = 1}$$

Imposing the continuity conditions (17.5), we have

$$\psi = \begin{cases} \cos(kL) \exp[\kappa(q + L)], & q < -L, \\ \cos(kq), & |q| \leq L, \\ \cos(kL) \exp[-\kappa(q - L)], & q > L, \end{cases} \quad (18.3)$$

¹⁸To be precise, note that even if $\mathcal{S}_0$ takes complex values for $\alpha \in \mathbb{C}$, there are no specific reasons to forbid this as both p and $\dot{q}$ are independent of α .

and

$$k \tan(kL) = \kappa. \quad (18.4)$$

A linearly independent solution is given by

$$\psi^D = [2k \sin(kL)]^{-1} \cdot \begin{cases} \cos(2kL) \exp[\kappa(q+L)] - \exp[-\kappa(q+L)], & q < -L, \\ 2 \sin(kL) \sin(kq), & |q| \leq L, \\ \exp[\kappa(q-L)] - \cos(2kL) \exp[-\kappa(q-L)], & q > L. \end{cases} \quad (18.5)$$

Note that any other possible solution has the form $A\psi^D + B\psi$. Let us now consider the trivializing map associated to this solution. To do this it is sufficient to find w . In the $(1, 1)$ case we have

$$\frac{\psi^D}{\psi} = [k \sin(2kL)]^{-1} \cdot \begin{cases} \cos(2kL) - \exp[-2\kappa(q+L)], & q < -L, \\ \sin(2kL) \tan(kq), & |q| \leq L, \\ \exp[2\kappa(q-L)] - \cos(2kL), & q > L. \end{cases} \quad (18.6)$$

Observe that

$$\lim_{q \rightarrow \pm\infty} \frac{\psi^D}{\psi} = \pm\infty. \quad (18.7)$$

Hence, in the $(1, 1)$ case the trivializing map is a local self-homeomorphism of $\hat{\mathbb{R}}$ as required by the EP, so that the solutions of (18.4) are physical energy levels. In particular, if E_n is the $(n+1)^{th}$ solution of (18.4), then it follows from (18.6) that the map covers $\hat{\mathbb{R}}$ $(n+1)$ -times.

$$\underline{i = 2, j = 1}$$

In the $(2, 1)$ case we have

$$\psi = k^{-1} \cdot \begin{cases} -\sin(kL) \exp[\kappa(q+L)], & q < -L, \\ \sin(kq), & |q| \leq L, \\ \sin(kL) \exp[-\kappa(q-L)], & q > L. \end{cases} \quad (18.8)$$

where

$$k \cot(kL) = -\kappa. \quad (18.9)$$

A linearly independent solution is given by

$$\psi^D = [2 \cos(kL)]^{-1} \cdot \begin{cases} \exp[-\kappa(q+L)] + \cos(2kL) \exp[\kappa(q+L)] & q < -L, \\ 2 \cos(kL) \cos(kq), & |q| \leq L, \\ \exp[\kappa(q-L)] + \cos(2kL) \exp[-\kappa(q-L)], & q > L. \end{cases} \quad (18.10)$$

and the ratio $w = \psi^D/\psi$ is given by

$$\frac{\psi^D}{\psi} = k[\sin(2kL)]^{-1} \cdot \begin{cases} -\cos(2kL) - \exp[-2\kappa(q+L)], & q < -L, \\ \sin(2kL) \cot(kq), & |q| \leq L, \\ \cos(2kL) + \exp[2\kappa(q-L)], & q > L. \end{cases} \quad (18.11)$$

Even in this case we have

$$\lim_{q \rightarrow \pm\infty} \frac{\psi^D}{\psi} = \pm\infty, \quad (18.12)$$

so that the associated trivializing map is a local self-homeomorphism of $\hat{\mathbb{R}}$ and the corresponding spectrum given by the solutions of (18.9) is a physical one.

$$\underline{i = 1, j = 2}$$

The (1, 2) case is the one of the two cases in which there are not solutions of the SE which are vanishing both at $-\infty$ and $+\infty$. We have

$$\psi = \begin{cases} \cos(kL) \exp[-\kappa(q + L)], & q < -L, \\ \cos(kq), & |q| \leq L, \\ \cos(kL) \exp[\kappa(q - L)], & q > L. \end{cases} \quad (18.13)$$

Also in this case we would have a quantized spectrum. In fact, the continuity conditions give

$$k \tan(kL) = -\kappa. \quad (18.14)$$

However, we now show that these solutions do not satisfy the EP. Let us consider the dual solution

$$\psi^D = [2k \sin(kL)]^{-1} \cdot \begin{cases} \cos(2kL) \exp[-\kappa(q + L)] - \exp[\kappa(q + L)], & q < -L, \\ 2 \sin(kL) \sin(kq), & |q| \leq L, \\ \exp[-\kappa(q - L)] - \cos(2kL) \exp[\kappa(q - L)], & q > L. \end{cases} \quad (18.15)$$

These solutions provide an explicit example of the general case we discussed in considering the asymptotic conditions coming from the EP. Namely, observe that in the (1, 2) case, both ψ and ψ^D are divergent at $\pm\infty$. Also observe that there is no way to have a solution vanishing both at $\pm\infty$. To see this, note that for a given E , solution of (18.14), any solution can be written as $\phi = A\psi^D + B\psi$. Let us choose ϕ in such a way that it vanishes at $+\infty$ that is $\phi = k\psi^D + \cot(2kL)\psi$, so that

$$\phi = [2 \sin(kL)]^{-1} \cdot \begin{cases} 2 \cos(2kL) \exp[-\kappa(q + L)] - \exp[\kappa(q + L)], & q < -L, \\ 2 \sin(kL) [\sin(kq) + \cot(2kL) \cos(kq)], & |q| \leq L, \\ \exp[\kappa(q - L)], & q > L. \end{cases} \quad (18.16)$$

Hence, even if $\phi \underset{q \rightarrow +\infty}{\sim} 0$, we have that it diverges at $-\infty$

$$\phi \underset{q \rightarrow -\infty}{\sim} -\frac{\cos(2kL)}{\sin(kL)} \exp[-\kappa(q + L)]. \quad (18.17)$$

Note that the only case in which ϕ would vanish at $-\infty$ is for $\cos(2kL) = 0$. On the other hand, (18.14) and number theoretical arguments show that this is never the case. We have seen that, depending on $\mathcal{W}$, there are cases with a quantized spectrum even if the SE does not have $L^2(\mathbb{R})$ solutions. However, as discussed in the general case, in this case the EP cannot be satisfied and the corresponding values of E

cannot be in the physical spectrum. Let us explicitly see how the $(1, 2)$ case would give a discontinuous trivializing map. To see this it is sufficient to study the behavior of the ratio ψ^D/ψ . By (18.13) and (18.15), we have

$$\frac{\psi^D}{\psi} = [k \sin(2kL)]^{-1} \cdot \begin{cases} \cos(2kL) - \exp[2\kappa(q + L)], & q < -L, \\ \sin(2kL) \tan(kq), & |q| \leq L, \\ \exp[2\kappa(q - L)] - \cos(2kL), & q > L, \end{cases} \quad (18.18)$$

whose asymptotic behavior is

$$\lim_{q \rightarrow \pm\infty} \frac{\psi^D}{\psi} = \mp k^{-1} \cot(2kL). \quad (18.19)$$

It follows that the only possibility to have $w(-\infty) = w(+\infty)$ is that $k^{-1} \cot(2kL) = 0$. However, this equation is not compatible with the condition (18.14). Hence, we have

$$w(-\infty) \neq w(+\infty). \quad (18.20)$$

It follows that the E 's solutions of (18.14) must be discarded. We also note that the other possibility which would give a trivializing map, corresponding to a local self-homeomorphism of $\hat{\mathbb{R}}$, would be for $w(-\infty) = \pm\infty = -w(+\infty)$. However, this case would correspond to $k^{-1} \cot(2kL) = \pm\infty$, which has not solutions compatible with (18.14).

$$\underline{i = 2, j = 2}$$

In this case we have

$$\psi = k^{-1} \cdot \begin{cases} -\sin(kL) \exp[-\kappa(q + L)], & q < -L, \\ \sin(kq), & |q| \leq L, \\ \sin(kL) \exp[\kappa(q - L)], & q > L, \end{cases} \quad (18.21)$$

where

$$k \cot(kL) = \kappa. \quad (18.22)$$

For the dual solution, we have

$$\psi^D = [2 \cos(kL)]^{-1} \cdot \begin{cases} \exp[\kappa(q + L)] + \cos(2kL) \exp[-\kappa(q + L)], & q < -L, \\ 2 \cos(kL) \cos(kq), & |q| \leq L, \\ \exp[-\kappa(q - L)] + \cos(2kL) \exp[\kappa(q - L)], & q > L, \end{cases} \quad (18.23)$$

and the ratio of the solutions is

$$\frac{\psi^D}{\psi} = k[\sin(2kL)]^{-1} \cdot \begin{cases} -\cos(2kL) - \exp[2\kappa(q + L)], & q < -L, \\ \sin(2kL) \cot(kq), & |q| \leq L, \\ \cos(2kL) + \exp[-2\kappa(q - L)], & q > L. \end{cases} \quad (18.24)$$

The asymptotic behavior is

$$\lim_{q \rightarrow \pm\infty} \frac{\psi^D}{\psi} = \pm k \cot(2kL). \quad (18.25)$$

It follows that the only possibility to have either $w(-\infty) = w(+\infty)$ or $w(-\infty) = \pm\infty = -w(+\infty)$ is that either $k \cot(2kL) = 0$ or $k \cot(2kL) = \pm\infty$. However, also in this case this equation is not compatible with the condition (18.22). Hence, we have

$$w(-\infty) \neq w(+\infty), \quad (18.26)$$

and the values of E solutions of (18.22) must be discarded.

Let us make some remarks. First, recall that even in this case, for any admissible E , there are Möbius states parameterized by the values of ℓ . Secondly, we note once again the crucial role of the dual solution: while ψ is vanishing at $\pm\infty$, the fact that ψ^D is divergent implies that p and $\dot{q}$ vanish exponentially at infinity. This makes evident the crucial role of the dual solution, an aspect which is peculiar of the present formulation. We also note that even in this case the classical limit is completely under control. For example, in the $i = j = 1$ case by (14.59)(18.3) and (18.5) we have

$$p = -\frac{\hbar}{2}(\ell + \bar{\ell})k^2 \sin^2(kL) \cdot \begin{cases} 4 |-\exp[-\kappa(q+L)] + [\cos(2kL) - ik\ell \sin(2kL)] \exp[\kappa(q+L)]|^{-2}, & q < -L, \\ \sin^{-2}(kL) |\sin(kq) - ik\ell \cos(kq)|^{-2}, & |q| \leq L, \\ 4 |\exp[\kappa(q-L)] - [\cos(2kL) + ik\ell \sin(2kL)] \exp[-\kappa(q-L)]|^{-2}, & q > L. \end{cases} \quad (18.27)$$

While in the region $|q| \leq L$ the classical limit parallels the analysis we considered in the case of the free particle, in the regions $|q| > L$ we have

$$\lim_{\hbar \rightarrow 0} p = 0, \quad (18.28)$$

which is a consequence of the $\exp(\pm kq)$ terms.

Above we considered the four cases in which ψ_i^1 is smoothly joined to ψ_j^2 at $q = L$. However, one can equivalently consider ψ_i^1 and an arbitrary linear combination $A\psi_1^2 + B\psi_2^2$. This is the example in the Introduction. In particular, by (1.36) we see that (1.41) is the $(2, 1)$ case, with (18.8) corresponding to (1.35) and (18.10) to (1.37). Similarly, by (1.38) the case (1.42) is the $(1, 1)$ case, with (18.3) corresponding to (1.37) and (18.5) to (1.35).

In conclusion, we note that the joining condition (15.8) implied by the EP, leads, as follows from the general results we proved, to the usual spectrum. We also saw that there is a quantized spectrum associated to the case in which there are divergent solutions only. Even if these are not admissible solutions, we note that this implied a doubling of the relevant transcendental equations.

18.2 The infinitely deep potential well and the free particle

The case of the infinitely potential well can be studied as limiting case in which $V_0 \rightarrow \infty$. However, we observe that for $V_0 = \infty$, there is a peculiar situation. The reason is that, according to the EP the reduced action cannot be a constant in a finite (or infinite) region. In the case of the infinitely deep potential well, we have $\mathcal{S}_0 = \text{const}$ and $p = 0$ in the region $|q| > L$. However, according to the EP,

we have that $\mathcal{S}_0$, and therefore p , spreads on all the existent space. In this sense, the infinitely deep potential well can be seen as a restriction of $\mathbb{R}$ to a finite interval. Then, in this case, the boundary at $q = \pm L$ plays the role of the points $\pm\infty$ and the joining condition (15.8) is replaced by

$$w(-L) = \begin{cases} w(+L), & \text{for } w(-L) \neq \pm\infty, \\ -w(+L), & \text{for } w(-L) = \pm\infty, \end{cases} \quad (18.29)$$

which is in fact satisfied as¹⁹

$$\frac{\psi^D}{\psi} = \begin{cases} k^{-1} \tan(\frac{n\pi}{2L}q), & n \text{ even}, \\ k \cot(\frac{n\pi}{2L}q), & n \text{ odd}. \end{cases} \quad (18.30)$$

In this way, the discontinuity at $q = \pm L$ of ψ^D , and therefore of $\partial_q \psi$, which unavoidably arises when $V_0 = +\infty$, naturally disappears. We note that, while this solution is a consequence of the EP, in the conventional approach to QM one considers the self-adjoint extension of the operators (see *e.g.* [71]). Observe that the solutions (18.30) correspond to the cases $i = 1, j = 1$ and $i = 2, j = 1$ in the limit $V_0 \rightarrow +\infty$. For the energy levels we have

$$E_n = \frac{\hbar^2 \pi^2}{8mL^2} n^2, \quad n = 1, 2, 3, \dots \quad (18.31)$$

This way of considering the infinitely deep well is also consistent with the limit case $L \rightarrow +\infty$ under which $\mathcal{S}_0$ reduces to the one of the free particle. In doing this one has to consider the double scaling limit $L \rightarrow +\infty$ and $n \rightarrow +\infty$, with the ratio n/L taking continuous finite values

$$\lim_{(n,L) \rightarrow (+\infty, +\infty)} \frac{n}{L} = \frac{\sqrt{8mE}}{\hbar\pi}. \quad (18.32)$$

Observe that this way of obtaining the free particle shows that in this case the trivializing map satisfies the continuity at $\pm\infty$ obtained from (18.29) in the $L \rightarrow +\infty$ limit. Furthermore, the discontinuity of the ratio $w = \psi^D/\psi$ at $\pm\infty$ of the non-admissible solutions disappears in the $L \rightarrow +\infty$ limit with the consequence that these become physical solutions. Let us also note that in the case of the free particle, the ratio of solutions is a Möbius transformation of $w_E = k^{-1} \tan kq$, so that the index of the trivializing map is infinite in this case

$$I[w_E] = \infty. \quad (18.33)$$

Note that $\lim_{E \rightarrow 0} w_E = q^0$, so that in this limit the right hand side of (18.33) collapses to 1. Let n be the maximum number of zeroes that a solution of the SE associated to $\mathcal{W}$ may have and denote by w_n the ratio of two linearly independent solutions of the SE. Local homeomorphicity of the trivializing map and Sturm's theorem on zeroes of solutions of the SE, yield

$$I[w_n] = n. \quad (18.34)$$

¹⁹Note that with this choice we have $w(-L) = w(+L)$. Interchanging ψ^D and ψ , this becomes $w(-L) = -w(+L)$.

18.3 The simple and double harmonic oscillators

Another relevant system with quantized spectrum is the harmonic oscillator. We now consider the case of the double harmonic oscillator, with the simple one corresponding to a particular case. The Hamiltonian describing the relative motion of the reduced mass of the double harmonic oscillator is

$$H = \frac{p^2}{2m} + \frac{1}{2}m\omega^2(|q| - q_0)^2. \quad (18.35)$$

The reduced action is given by (14.57) with ψ^D and ψ real linearly independent solutions of the SE

$$\left(-\frac{\hbar^2}{2m}\frac{\partial^2}{\partial q^2} + \frac{1}{2}m\omega^2(|q| - q_0)^2\right)\psi = E\psi. \quad (18.36)$$

Let us set

$$E = \left(\mu + \frac{1}{2}\right)\hbar\omega, \quad (18.37)$$

and

$$z' = \alpha(q + q_0), \quad q \leq 0, \quad z = \alpha(q - q_0), \quad q \geq 0, \quad (18.38)$$

where $\alpha = \sqrt{2m\omega/\hbar}$. Note that $z'(-q) = -z(q)$ and that for $q_0 = 0$ the system reduces to the simple harmonic oscillator. We stress that since we consider (18.36) for any real E , at this stage μ is an arbitrary real number. The SE (18.36) is equivalent to

$$\frac{\partial^2 \psi}{\partial z'^2} + \left(\mu + \frac{1}{2} - \frac{z'^2}{4}\right)\psi = 0, \quad q \leq 0, \quad (18.39)$$

and

$$\frac{\partial^2 \psi}{\partial z^2} + \left(\mu + \frac{1}{2} - \frac{z^2}{4}\right)\psi = 0, \quad q \geq 0. \quad (18.40)$$

A solution of (18.40) is given for any μ by the parabolic cylinder function (see *e.g.* [72])

$$D_\mu(z) = 2^{\frac{\mu}{2}} e^{-\frac{z^2}{4}} \left[\frac{\Gamma(\frac{1}{2})}{\Gamma[\frac{1-\mu}{2}]} {}_1F_1\left(-\frac{\mu}{2}; \frac{1}{2}; \frac{z^2}{2}\right) + \frac{z}{\sqrt{2}} \frac{\Gamma(-\frac{1}{2})}{\Gamma(-\frac{\mu}{2})} {}_1F_1\left[\frac{1-\mu}{2}; \frac{3}{2}; \frac{z^2}{2}\right] \right], \quad (18.41)$$

where ${}_1F_1$ is the confluent hypergeometric function

$${}_1F_1(a; c; z) = 1 + \frac{a}{c} \frac{z}{1!} + \frac{a(a+1)}{c(c+1)} \frac{z^2}{2!} + \dots \quad (18.42)$$

We are interested in the behavior of w at $\pm\infty$. If $w(-\infty) \neq \pm\infty$ we have to impose $w(-\infty) = w(+\infty)$, while if $w(-\infty) = \pm\infty$, then we should have $w(-\infty) = -w(+\infty)$. To consider this aspect we need the behavior of D_μ for $|z| \gg 1$, $|z| \gg |\mu|$. For $\pi/4 < \arg z < 5\pi/4$ we have

$$D_\mu(z) \underset{|z| \gg 1}{\sim} -\frac{\sqrt{2\pi}}{\Gamma(-\mu)} e^{\mu\pi i} e^{z^2/4} z^{-\mu-1} \left[1 + \frac{(\mu+1)(\mu+2)}{2z^2} + \frac{(\mu+1)(\mu+2)(\mu+3)(\mu+4)}{2 \cdot 4z^4} + \dots \right]$$

$$+ e^{-z^2/4} z^\mu \left[1 - \frac{\mu(\mu-1)}{2z^2} + \frac{\mu(\mu-1)(\mu-2)(\mu-3)}{2 \cdot 4z^4} - \dots \right], \quad (18.43)$$

while for $|\arg z| < 3\pi/4$

$$D_\mu(z) \underset{|z| \gg 1}{\sim} e^{-z^2/4} z^\mu \left[1 - \frac{\mu(\mu-1)}{2z^2} + \frac{\mu(\mu-1)(\mu-2)(\mu-3)}{2 \cdot 4z^4} - \dots \right]. \quad (18.44)$$

A property of $D_\mu(z)$ is that also $D_\mu(-z)$ solves (18.40). If μ is a non-negative integer, then $D_\mu(z)$ and $D_\mu(-z)$ coincide.²⁰ Let us then consider the case in which μ is not a non-negative integer. Two sets of solutions of (18.36), which can be written in the form (18.39)(18.40), are

$$\psi_\pm = \begin{cases} \pm D_\mu(-\alpha(q+q_0)), & q \leq 0, \\ D_\mu(\alpha(q-q_0)), & q \geq 0, \end{cases} \quad (18.45)$$

where μ is solution of the continuity condition at $q = 0$. For the even solution ψ_+ , this gives

$$D'_\mu(-\alpha q_0) = 0, \quad (18.46)$$

whereas for the odd one ψ_- , we have

$$D_\mu(-\alpha q_0) = 0. \quad (18.47)$$

For each μ , solution of Eq.(18.46) (Eq.(18.47)), we have that a solution ψ_-^D (ψ_+^D) of the SE (18.36), linearly independent from ψ_+ (ψ_-), is given by

$$\psi_\mp^D = \begin{cases} \mp D_\mu(\alpha(q+q_0)) \mp a_\mp D_\mu(-\alpha(q+q_0)), & q \leq 0, \\ D_\mu(-\alpha(q-q_0)) + a_\mp D_\mu(\alpha(q-q_0)), & q \geq 0. \end{cases} \quad (18.48)$$

We have

$$a_- = -\frac{D_\mu(\alpha q_0)}{D_\mu(-\alpha q_0)}, \quad (18.49)$$

with μ solution of Eq.(18.46). Similarly

$$a_+ = \frac{D'_\mu(\alpha q_0)}{D'_\mu(-\alpha q_0)}, \quad (18.50)$$

where μ is solution of Eq.(18.47). Therefore, for a given μ solution of Eq.(18.46) (Eq.(18.47)), ψ_+ and ψ_-^D (ψ_- and ψ_+^D) are linearly independent solutions of the SE (18.36). The transcendental equations (18.46) and (18.47) determine, each one, a set of possible energy eigenvalues. Let us now consider the asymptotic behavior of ψ_-^D/ψ_+ and ψ_+^D/ψ_- . By (18.45) and (18.48), we have

$$\frac{\psi_-^D}{\psi_+} = \begin{cases} -a_- - D_\mu(\alpha(q+q_0))/D_\mu(-\alpha(q+q_0)), & q \leq 0, \\ a_- + D_\mu(-\alpha(q-q_0))/D_\mu(\alpha(q-q_0)), & q \geq 0, \end{cases} \quad (18.51)$$

²⁰Note that if μ is a non-negative integer, then $\Gamma^{-1}(-\mu) = 0$, so that in this case the first term in (18.43) cancels.

and

$$\frac{\psi_+^D}{\psi_-} = \begin{cases} -a_+ - D_\mu(\alpha(q + q_0)) / D_\mu(-\alpha(q + q_0)), & q \leq 0, \\ a_+ + D_\mu(-\alpha(q - q_0)) / D_\mu(\alpha(q - q_0)), & q \geq 0. \end{cases} \quad (18.52)$$

By (18.43) and (18.44), we have the asymptotics²¹

$$\frac{\psi_-^D}{\psi_+} \underset{q \rightarrow -\infty}{\sim} \frac{\sqrt{2\pi}}{\Gamma(-\mu)} e^{\mu\pi i} e^{z'^2/2} z'^{-\mu-1} (-z')^{-\mu} \longrightarrow -\infty, \quad (18.53)$$

and

$$\frac{\psi_-^D}{\psi_+} \underset{q \rightarrow +\infty}{\sim} -\frac{\sqrt{2\pi}}{\Gamma(-\mu)} e^{\mu\pi i} e^{z^2/2} z^{-\mu} (-z)^{-\mu-1} \longrightarrow +\infty. \quad (18.54)$$

By (18.51)(18.52) we see that ψ_+^D/ψ_- and ψ_-^D/ψ_+ have the same asymptotic limits. Hence, since

$$\lim_{q \rightarrow -\infty} \frac{\psi_-^D}{\psi_+} = -\infty = -\lim_{q \rightarrow +\infty} \frac{\psi_-^D}{\psi_+}, \quad (18.55)$$

and similarly for ψ_+^D/ψ_- , we have that the trivializing map associated to the double harmonic oscillator, with energy levels given by (18.46) and (18.47), is a local homeomorphism of $\hat{\mathbb{R}}$ into itself. Other possible solutions of the SE (18.36) are

$$\psi_\pm = \begin{cases} \pm D_\mu(\alpha(q + q_0)), & q \leq 0, \\ D_\mu(-\alpha(q - q_0)), & q \geq 0, \end{cases} \quad (18.56)$$

where μ is solution of the continuity condition at $q = 0$. For the even solution ψ_+ , this condition gives

$$D'_\mu(\alpha q_0) = 0, \quad (18.57)$$

whereas for the odd one ψ_- , we have

$$D_\mu(\alpha q_0) = 0. \quad (18.58)$$

For each μ solution of Eq.(18.57) (Eq.(18.58)) we have that a solution ψ_-^D (ψ_+^D) of (18.36), linearly independent from ψ_+ (ψ_-), is given by

$$\psi_\mp^D = \begin{cases} \mp D_\mu(-\alpha(q + q_0)) \mp b_\mp D_\mu(\alpha(q + q_0)), & q \leq 0, \\ D_\mu(\alpha(q - q_0)) + b_\mp D_\mu(-\alpha(q - q_0)), & q \geq 0, \end{cases} \quad (18.59)$$

where

$$b_- = -\frac{D_\mu(-\alpha q_0)}{D_\mu(\alpha q_0)}, \quad (18.60)$$

²¹The asymptotics (18.43) and (18.44) include $\arg z' = \pi$ and $\arg z = 0$ respectively, so that

$$z'^{-\mu-1}(-z')^{-\mu} = -e^{-\mu\pi i} |z'|^{-2\mu-1}, \quad z^{-\mu}(-z)^{-\mu-1} = -e^{-\mu\pi i} |z|^{-2\mu-1}.$$

with μ solution of Eq.(18.57). Similarly

$$b_+ = \frac{D'_\mu(-\alpha q_0)}{D'_\mu(\alpha q_0)}, \quad (18.61)$$

where μ is solution of Eq.(18.58). By (18.56) and (18.59), we have

$$\frac{\psi_-^D}{\psi_+} = \begin{cases} -b_- - D_\mu(-\alpha(q+q_0))/D_\mu(\alpha(q+q_0)), & q \leq 0, \\ b_- + D_\mu(\alpha(q-q_0))/D_\mu(-\alpha(q-q_0)), & q \geq 0, \end{cases} \quad (18.62)$$

and

$$\frac{\psi_+^D}{\psi_-} = \begin{cases} -b_+ - D_\mu(-\alpha(q+q_0))/D_\mu(\alpha(q+q_0)), & q \leq 0, \\ b_+ + D_\mu(\alpha(q-q_0))/D_\mu(-\alpha(q-q_0)), & q \geq 0. \end{cases} \quad (18.63)$$

By (18.43) and (18.44), we have the asymptotics

$$\frac{\psi_-^D}{\psi_+} \underset{q \rightarrow \pm\infty}{\sim} \pm b_-, \quad (18.64)$$

and

$$\frac{\psi_+^D}{\psi_-} \underset{q \rightarrow \pm\infty}{\sim} \pm b_+. \quad (18.65)$$

Since both b_- and b_+ are finite non-vanishing, we have that the ratios ψ_-^D/ψ_+ and ψ_+^D/ψ_- associated to the solutions (18.57) and (18.58) respectively, do not satisfy the joining condition (15.8). Therefore, according to the EP, the energy eigenvalues defined by (18.57) and (18.58) are not physical ones.

Thus, besides the potential well, also in this case, in agreement with the general theorem (17.25), the only physical solutions are (18.46)(18.47), with the $L^2(\mathbb{R})$ solutions given in (18.45).

Let us now consider the simple harmonic oscillator. In this case $q_0 = 0$, that is $z' = z$. By (18.46) and (18.47) it follows that the continuity conditions in the case of the harmonic oscillator are

$$D_\mu(0) = 0, \quad (18.66)$$

and

$$D'_\mu(0) = 0. \quad (18.67)$$

Let us write down the initial values of D_μ

$$D_\mu(0) = 2^{\mu/2} \frac{\Gamma(1/2)}{\Gamma[(1-\mu)/2]}, \quad D'_\mu(0) = 2^{(\mu-1)/2} \frac{\Gamma(-1/2)}{\Gamma(-\mu/2)}. \quad (18.68)$$

Since the poles of $\Gamma(\mu)$ are at the non-positive integer values of μ , the solutions of (18.66) are

$$\mu = 2n + 1, \quad n = 0, 1, 2, \dots, \quad (18.69)$$

while for Eq.(18.67) we have

$$\mu = 2n, \quad n = 0, 1, 2, \dots \quad (18.70)$$

For non-negative integer values of n we have

$$D_n(z) = (-1)^n e^{z^2/4} \frac{d^n}{dz^n} e^{-z^2/2} = e^{-z^2/4} H_n(z), \quad (18.71)$$

where the H_n 's are the Hermite polynomials. By (18.37)(18.69) and (18.70), we obtain the spectrum of the harmonic oscillator

$$E = \left(n + \frac{1}{2}\right) \hbar\omega, \quad n = 0, 1, 2, \dots \quad (18.72)$$

We already observed that since $\Gamma^{-1}(-n) = 0$ for $n = 0, 1, 2, \dots$, it follows that $D_n(z)$ and $D_n(-z)$ are linearly dependent. Besides Eq.(18.71), this can be also seen from the fact that $D_n(z)$ and $D_n(-z)$ are both convergent with the same leading terms, implies that $D_n(z) = (-1)^n D_n(-z)$ (where the $(-1)^n$ factor follows from (18.43) and (18.44)). This also follows from the relation

$$D_\mu(z) = e^{\mu\pi i} D_\mu(-z) + \frac{\sqrt{2\pi}}{\Gamma(-\mu)} e^{(\mu+1)\pi i/2} D_{-\mu-1}(-iz), \quad (18.73)$$

which implies that $D_{-n-1}(-iz)$ is a solution of the SE of the simple harmonic oscillator, which is linearly independent from $D_n(z)$. Then, in this case one has

$$\psi = D_n(\alpha q), \quad \psi^D = e^{-(n+1)\pi i/2} D_{-n-1}(-i\alpha q), \quad (18.74)$$

where the role of the factor $e^{-(n+1)\pi i/2}$ is that of making ψ^D real. The ratio

$$\frac{\psi^D}{\psi} = e^{-(n+1)\pi i/2} \frac{D_{-n-1}(-i\alpha q)}{D_n(\alpha q)}, \quad (18.75)$$

has the asymptotics

$$\frac{\psi^D}{\psi} \underset{q \rightarrow -\infty}{\sim} e^{\alpha^2 q^2/2} (\alpha q)^{-2n-1} \longrightarrow -\infty, \quad (18.76)$$

$$\frac{\psi^D}{\psi} \underset{q \rightarrow +\infty}{\sim} e^{\alpha^2 q^2/2} (\alpha q)^{-2n-1} \longrightarrow +\infty, \quad (18.77)$$

showing that, according to the general result we derived previously, also the standard spectrum of the harmonic oscillator defines a local homeomorphism of $\hat{\mathbb{R}}$ into itself.

18.4 The general case

In the previous investigation we considered solutions of the SE for which ψ contains either the vanishing or divergent solution. However, one may consider the following general form

$$\psi = \begin{cases} D_\mu(-z') + cD_\mu(z'), & q \leq 0, \\ D_\mu(z) + cD_\mu(-z), & q \geq 0, \end{cases} \quad (18.78)$$

where

$$c = \frac{D'_\mu(-\alpha q_0)}{D'_\mu(\alpha q_0)}. \quad (18.79)$$

For any given μ , a linearly independent solution is given by

$$\psi^D = \begin{cases} -D_\mu(-z') - dD_\mu(z'), & q \leq 0, \\ D_\mu(z) + dD_\mu(-z), & q \geq 0, \end{cases} \quad (18.80)$$

where

$$d = -\frac{D_\mu(-\alpha q_0)}{D_\mu(\alpha q_0)}. \quad (18.81)$$

The ratio

$$\frac{\psi_+^D}{\psi_-} = \begin{cases} -(D_\mu(-z') + dD_\mu(z'))/(D_\mu(-z') + cD_\mu(z')), & q \leq 0, \\ (D_\mu(z) + dD_\mu(-z))/(D_\mu(z) + cD_\mu(-z)), & q \geq 0, \end{cases} \quad (18.82)$$

has the asymptotics behavior

$$\lim_{q \rightarrow \pm\infty} \frac{\psi^D}{\psi} = \pm \frac{d}{c}. \quad (18.83)$$

This shows that Eq.(15.8) is satisfied in the case in which either

$$c = 0, \quad (18.84)$$

or

$$d = 0. \quad (18.85)$$

These cases correspond to the ones discussed in (18.45)–(18.55). Note that if $\mu = 0, 1, 2, \dots$, then ψ in (18.78) and its dual (18.80) are not linearly independent. In this case there is always a solution vanishing both at $-\infty$ and $+\infty$. If $q_0 = 0$, then this situation corresponds to the harmonic oscillator. Generally, for an arbitrary q_0 and for $\mu = 0, 1, 2, \dots$, the solution ψ^D in (18.80) is replaced by

$$\psi^D = \begin{cases} -D_n(-z') - d'D_{-n-1}(-iz'), & q \leq 0, \\ D_n(z) + d'D_{-n-1}(iz), & q \geq 0, \end{cases} \quad (18.86)$$

where now

$$d' = -\frac{D_n(-\alpha q_0)}{D_n(-i\alpha q_0)}. \quad (18.87)$$

We note that above we used the parabolic cylinder functions with real argument. Then, the fact that for $\mu = 0, 1, 2, \dots$, $D_\mu(z)$ and $D_\mu(-z)$ are not linearly independent, forced us to use $D_\mu(-iz)$. In this context we observe that $D_\mu(z)$ and $D_\mu(-iz)$ are always linearly independent so that the dual solution (18.86) can be extended to arbitrary values of μ .

19 Generalizations

This section has been written in collaboration with Gaetano Bertoldi.

A property of the SE, which can be seen as a consequence of the EP, is that the reduced action associated to the state $\mathcal{W}^0$ is not constant. Actually, the structure of this solution is rather peculiar and turns out to be sufficient to fix the SE both in any dimension and for time-dependent potentials. This is not a surprise since basic features of the usual approach to QM, such as uncertainty relations, energy quantization, tunnel effect, Hilbert spaces etc. already arise in the one-dimensional stationary case. In this section we derive the quantum HJ equation for time-dependent potentials. Next, we will consider the higher dimensional case. Finally, we will derive the relativistic quantum HJ equation.

19.1 Time-dependent Schrödinger equation

Let us start by noticing that since for stationary states one has

$$\mathcal{S}(q, t) = \mathcal{S}_0(q) - Et, \quad (19.1)$$

it follows that we can rewrite the QSHJE (9.44) in the form

$$\frac{\partial \mathcal{S}}{\partial t} + \frac{1}{2m} \left(\frac{\partial \mathcal{S}(q)}{\partial q} \right)^2 + V(q) + \frac{\hbar^2}{4m} \{\mathcal{S}, q\} = 0. \quad (19.2)$$

This equation and Eq.(10.2) are equivalent to

$$i\hbar \frac{\partial \psi(q, t)}{\partial t} = \left(-\frac{\hbar^2}{2m} \frac{\partial^2}{\partial q^2} + V(q) \right) \psi(q, t), \quad (19.3)$$

with

$$\psi(q, t) = \frac{1}{\sqrt{\mathcal{S}'}} e^{\frac{i}{\hbar} \mathcal{S}}. \quad (19.4)$$

A peculiar property of Eq.(19.3) is that it is satisfied by all possible solutions of the SE irrespectively of the specific energy eigenvalue. Let us set²²

$$\Phi_k = \frac{1}{\sqrt{\mathcal{S}'_k}} e^{\frac{i}{\hbar} \mathcal{S}_k} = e^{-\frac{i}{\hbar} E_k t} \bar{\phi}_k, \quad (19.5)$$

where $\mathcal{S}_k = \mathcal{S}_{0,k} - E_k t$, and $\mathcal{S}_{0,k}$ is the reduced action corresponding to the eigenvalue E_k and

$$\phi_k = \frac{1}{\sqrt{\mathcal{S}'_{0,k}}} e^{-\frac{i}{\hbar} \mathcal{S}_{0,k}}. \quad (19.6)$$

Since

$$\left(-\frac{\hbar^2}{2m} \frac{\partial^2}{\partial q^2} + V(q) \right) \bar{\phi}_k = E_k \bar{\phi}_k, \quad (19.7)$$

²²Here we consider the case of discrete spectra, however similar arguments extend to the continuous ones.

we have that Eq.(19.3) is solved by

$$\Psi(q, t) = \sum_k c_k \Phi_k, \quad (19.8)$$

with c_k 's arbitrary constants. While $\bar{\Psi}$ satisfies the complex conjugated of Eq.(19.3), the function

$$\Psi^D(q, t) = \sum_k d_k \Phi_k^D, \quad (19.9)$$

where $\Phi_k^D = e^{-\frac{i}{\hbar} E_k t} \phi_k$, is still solution of (19.3). In particular, the most general solution of (19.3) is

$$\Phi = \sum_k (A_k \Phi_k^D + B_k \Phi_k). \quad (19.10)$$

An interesting aspect of Eq.(19.3) is that, like (19.2), the energy level does not appear: it is not specified, it appears, as in the classical HJ equation, as an integration constant. On the other hand, this is precisely the same role played by α and ℓ . As in the SE (10.2) there is no information about the initial conditions determining α and ℓ , in the SE (19.3) this lack of information is extended to the energy level. This aspect is evident in Eq.(19.3) where in order to determine $\mathcal{S}$ we need to know the initial conditions fixing the energy level E and the constant α and ℓ . Therefore, in the framework of the Quantum HJ Equation associated to Eq.(19.3) and that will be considered later on, the lack of information on the energy level is similar in nature to that associated to the integration constants α and ℓ . As a consequence, while a given solution of the SE (10.2) is not sufficient to specify $\mathcal{S}_0$, in the case of Eq.(19.3) the energy level E should be specified as one more integration constant. Fixing the value of E would be equivalent to specify which is the non-vanishing pair (A_k, B_k) in (19.10). Once this is done, one needs further conditions to completely fix $\mathcal{S}$. This analogy has the following natural consequence: as in the case of ψ_E , whose knowledge in general does not completely fix $\mathcal{S}_0$, the knowledge of Φ in (19.10) is not sufficient to completely fix $\mathcal{S}$.

Let us now consider the case of a system that at the time $t < t_1$ is described by $\mathcal{S}_{0,k}$, corresponding to the energy level E_k , associated to a given potential $V(q)$. Subsequently a perturbation, described by some time-dependent potential, is turned on in such a way that the system at the time $t > t_2 > t_1$ has a reduced action $\mathcal{S}_{0,n}$ corresponding to the energy level E_n . Although Φ_k and Φ_n satisfy the SE (19.7) with different energy level, they satisfy the same SE in the form of Eq.(19.3). In other words, (19.3) is the equation which remains form invariant for $t < t_1$ and $t > t_2$, and of which both Φ_k and Φ_n are solutions. As we have seen, this property is related to the linearity of Eq.(19.3). We now show that linearity and consistency imply that in the case of time-dependent potentials we have

$$i\hbar \frac{\partial \psi(q, t)}{\partial t} = \left(-\frac{\hbar^2}{2m} \frac{\partial^2}{\partial q^2} + V(q, t) \right) \psi(q, t). \quad (19.11)$$

To show this we first observe that to find the generalization of Eq.(19.3) in the case of time-dependent potentials the equation should have the following properties

- a) Linearity.
- b) In the time-independent case it should reduce to Eq.(19.3).

c) In the $\hbar \longrightarrow 0$ limit it should reduce to the CHJE (2.12).

These three conditions imply that the generalization of Eq.(19.3) to the case in which the potential depends on time has the form

$$\sum_{\alpha,\beta,\gamma,\delta} c_{\alpha\beta\gamma\delta} m^\alpha \hbar^\beta \frac{\partial^{\gamma+\delta} \psi}{\partial t^\gamma \partial q^\delta} = \left(-\frac{\hbar^2}{2m} \frac{\partial^2}{\partial q^2} + V(q, t) \right) \psi, \quad (19.12)$$

with $c_{\alpha\beta\gamma\delta}$ dimensionless constants. Even if the case of the time-dependent perturbation, considered previously, already shows that the only non-vanishing coefficient in (19.12) is $c_{0110} = i$, we show how (19.11) can be derived from (19.12) as a consequence of the points *a* and *b* above. The fact that in the stationary case Eq.(19.12) should correspond to (10.2), which is equivalent to (19.3), implies

$$\gamma \geq 1. \quad (19.13)$$

Let us now perform the dimensional analysis. A comparison between the left- and right-hand sides of Eq.(19.12) shows that the operator $m^\alpha \hbar^\beta \partial_t^\gamma \partial_q^\delta$ has the dimension of the energy, so that

$$\beta = 1 - \alpha, \quad \gamma = 1 + \alpha, \quad \delta = -2\alpha. \quad (19.14)$$

Eqs.(19.13) and (19.14) yield

$$\gamma \geq 1 \longrightarrow \alpha \geq 0 \longrightarrow \begin{cases} \beta \leq 1, \\ \delta \leq 0. \end{cases} \quad (19.15)$$

On the other hand, we have $\delta \geq 0$, so that by (19.14) and (19.15) one obtains

$$\alpha = 0, \quad \beta = 1, \quad \gamma = 1, \quad \delta = 0. \quad (19.16)$$

Therefore, the only surviving term in the left hand side of (19.12) is c_{0110} . Comparing with the stationary case we get $c_{0110} = i$, that is Eq.(19.12) reduces to (19.11). Making the identification

$$\psi(q, t) = R(q, t) e^{\frac{i}{\hbar} \mathcal{S}(q, t)}, \quad (19.17)$$

and using (19.11) we obtain the Quantum HJ Equation

$$\frac{\partial \mathcal{S}}{\partial t} + \frac{1}{2m} \left(\frac{\partial \mathcal{S}}{\partial q} \right)^2 + V(q, t) - \frac{\hbar^2}{2mR} \frac{\partial^2 R}{\partial q^2} = 0, \quad (19.18)$$

$$\frac{\partial R^2}{\partial t} + \frac{1}{m} \frac{\partial}{\partial q} \left(R^2 \frac{\partial \mathcal{S}}{\partial q} \right) = 0. \quad (19.19)$$

Note that in general the quantum potential $-\hbar^2 \partial_q^2 R / 2mR$, does not correspond to the one considered in the literature (*e.g.* the Bohmian one). The reason is that in general ψ in (19.17), solution of the SE, does not correspond to the wave-function. Thus, for example, in the stationary case, in which $\mathcal{S}(q, t) = \mathcal{S}_0(q) - Et$, we always have to satisfy the conditions (15.7), in particular $\mathcal{S}_0 \neq \text{const.}$

19.2 Higher dimension

Similarly to the case of the Heisenberg uncertainty relations

$$\Delta p_k \Delta q_k \geq \hbar/2, \quad (19.20)$$

in spite of being intrinsically one-dimensional, also our formulation essentially implies QM in higher dimension. In fact, it turns out that the EP implies the SE also in higher dimension. A detailed analysis of this argument is the subject of the forthcoming paper [16]. Here, we first outline the basic steps in [16] and then propose an alternative approach based on space compactification. Let us consider the D -dimensional CSHJE

$$\frac{1}{2m} \sum_{k=1}^D \left(\frac{\partial \mathcal{S}_0^{cl}(q)}{\partial q_k} \right)^2 + \mathcal{W}(q) = 0. \quad (19.21)$$

Also in this case, given another system with reduced action $\mathcal{S}_0^{clv}$, we set

$$\mathcal{S}_0^{clv}(q^v) = \mathcal{S}_0^{cl}(q). \quad (19.22)$$

Since $\mathcal{S}_0^{clv}(q^v)$ must satisfy the CSHJE

$$\frac{1}{2m} \sum_{k=1}^D \left(\frac{\partial \mathcal{S}_0^{clv}(q^v)}{\partial q_k^v} \right)^2 + \mathcal{W}^v(q^v) = 0, \quad (19.23)$$

it follows by (19.22) that

$$p_k \longrightarrow p_k^v = \frac{\partial \mathcal{S}_0^{clv}(q^v)}{\partial q_k^v} = \sum_{i=1}^D \frac{\partial q_i}{\partial q_k^v} \frac{\partial \mathcal{S}_0^{cl}(q)}{\partial q_i} = \sum_{i=1}^D J_{ki} p_i, \quad (19.24)$$

where J_{ki} denotes the Jacobian matrix

$$J_{ki} = \frac{\partial q_i}{\partial q_k^v}. \quad (19.25)$$

Let us set

$$(p^v|p) = \frac{\sum_k p_k^{v^2}}{\sum_k p_k^2} = \frac{p^t J^t J p}{p^t p}. \quad (19.26)$$

Note that in the one-dimensional case

$$(p^v|p) = \left(\frac{p^v}{p} \right)^2 = \left(\frac{\partial \mathcal{S}_0^{cl}}{\partial q^v} \frac{\partial q}{\partial \mathcal{S}_0^{cl}} \right)^2 = \left(\frac{\partial q^v}{\partial q} \right)^{-2}, \quad (19.27)$$

so that the Jacobian of the coordinate transformation is the ratio of momenta. By (19.21) we have

$$\mathcal{W}(q) \longrightarrow \mathcal{W}^v(q^v) = (p^v|p) \mathcal{W}(q), \quad (19.28)$$

that for the state $\mathcal{W}^0$ gives

$$\mathcal{W}^0(q^0) \longrightarrow \mathcal{W}^v(q^v) = (p^v|p^0) \mathcal{W}^0(q^0) = 0. \quad (19.29)$$

Thus we have that in any dimension the EP cannot be consistently implemented in CM. It is therefore clear that also in the higher dimensional case the implementation of the EP requires the deformation of the CSHJE. Therefore

$$\frac{1}{2m} \sum_{k=1}^D \left(\frac{\partial \mathcal{S}_0(q)}{\partial q_k} \right)^2 + \mathcal{W}(q) + Q(q) = 0. \quad (19.30)$$

The properties of $\mathcal{W} + Q$ under the VT (19.22) are determined by the transformed equation

$$\frac{1}{2m} \sum_{k=1}^D \left(\frac{\partial \mathcal{S}_0^{cl v}(q^v)}{\partial q_k^v} \right)^2 + \mathcal{W}^v(q^v) + Q^v(q^v) = 0, \quad (19.31)$$

which by (19.22) and (19.30) yields

$$\mathcal{W}^v(q^v) + Q^v(q^v) = (p^v|p) (\mathcal{W}(q) + Q(q)). \quad (19.32)$$

The only possibility to reach any state $\mathcal{W}^v$ from $\mathcal{W}^0$ is that it transforms with an inhomogeneous term. Namely, as $\mathcal{W}^0(q^0) \longrightarrow \mathcal{W}^v(q^v) \neq 0$, it follows by (19.32) that for each pair of states we have

$$\mathcal{W}^v(q^v) = (p^v|p^a) \mathcal{W}^a(q^a) + (q^a; q^v), \quad (19.33)$$

and

$$Q^v(q^v) = (p^v|p^a) Q^a(q^a) - (q^a; q^v). \quad (19.34)$$

Setting $\mathcal{W}^a(q^a) = \mathcal{W}^0(q^0)$ in Eq.(19.33) yields

$$\mathcal{W}^v(q^v) = (q^0; q^v), \quad (19.35)$$

so that, according to the EP, even in higher dimension all the states correspond to the inhomogeneous part in the transformation of the state $\mathcal{W}^0$ induced by v -maps. Comparing

$$\mathcal{W}^b(q^b) = (p^b|p^a) \mathcal{W}^a(q^a) + (q^a; q^b) = (q^0; q^b), \quad (19.36)$$

with the same formula with q^a and q^b interchanged, we have

$$(q^b; q^a) = -(p^a|p^b) (q^a; q^b), \quad (19.37)$$

in particular

$$(q; q) = 0. \quad (19.38)$$

More generally, comparing

$$\mathcal{W}^b(q^b) = (p^b|p^c) \mathcal{W}^c(q^c) + (q^c; q^b) = (p^b|p^a) \mathcal{W}^a(q^a) + (p^b|p^c) (q^a; q^c) + (q^c; q^b), \quad (19.39)$$

with (19.36) we obtain the basic cocycle condition

$$(q^a; q^c) = (p^c|p^b) [(q^a; q^b) - (q^c; q^b)], \quad (19.40)$$

which is a direct consequence of the EP. Eq.(19.40) is the higher dimensional generalization of Eq.(8.14). Similarly to the one-dimensional case, also Eq.(19.40) univocally leads to the higher dimensional SE. The details of the derivation will be given in [16]. Let us just mention that the higher dimensional version of the identity (9.7) is [16]

$$\alpha^2(\nabla\mathcal{S}_0) \cdot (\nabla\mathcal{S}_0) = \frac{\Delta(Re^{\alpha\mathcal{S}_0})}{Re^{\alpha\mathcal{S}_0}} - \frac{\Delta R}{R} - \alpha \left(2 \frac{\nabla R \cdot \nabla\mathcal{S}_0}{R} + \Delta\mathcal{S}_0 \right), \quad (19.41)$$

for any constant α . Therefore, if R satisfies the continuity equation

$$\nabla R^2 \cdot \nabla\mathcal{S}_0 + R^2 \Delta\mathcal{S}_0 = 0, \quad (19.42)$$

we have

$$(\nabla\mathcal{S}_0) \cdot (\nabla\mathcal{S}_0) = \hbar^2 \left(\frac{\Delta R}{R} - \frac{\Delta(Re^{\pm \frac{i\mathcal{S}_0}{\hbar}})}{Re^{\pm \frac{i\mathcal{S}_0}{\hbar}}} \right), \quad (19.43)$$

and the general form of the QHJE in D -dimensions is provided by (19.42) and

$$\frac{1}{2m}(\nabla\mathcal{S}_0) \cdot (\nabla\mathcal{S}_0) + \mathcal{W} + Q = 0, \quad (19.44)$$

where Q is the quantum potential

$$Q(q) = -\frac{\hbar^2}{2m} \frac{\Delta R}{R}. \quad (19.45)$$

We stress the important fact that in general, as in the case of the other Quantum HJ Equations we derived, Q does not correspond to the usual quantum potential. The reason for this is always the same. Namely, while in the standard approach Eqs.(19.44)(19.45) are derived from the Schrödinger equation by identifying $Re^{\frac{i\mathcal{S}_0}{\hbar}}$ with the wave-function, here the identification is between $Re^{\frac{i\mathcal{S}_0}{\hbar}}$ and a solution of the Schrödinger equation which in general does not correspond to the wave-function.

Here we show an alternative derivation with respect to [16] which is based on space compactification. As we will see, the approach is reminiscent of that in string theory and sets in a natural way a correspondence between potentials and space compactification. We will see that this approach poses a mathematical problem and will lead to the suggestion that potentials (forces) present in Nature have a geometrical origin, as they are connected with the properties of compactification. Let us start by observing that for potentials of the form

$$V(q) = \sum_{k=1}^D V_k(q_k), \quad (19.46)$$

the EP implies the decoupled quantum HJ equations

$$\frac{1}{2m} \left(\frac{\partial \mathcal{S}_{0,k}(q_k)}{\partial q_k} \right)^2 + \mathcal{W}_k(q_k) + \frac{\hbar^2}{4m} \{\mathcal{S}_{0,k}, q_k\} = 0, \quad (19.47)$$

$k = 1, \dots, D$, where $\mathcal{W}_k(q_k) \equiv V_k(q_k) - E_k$. Eq.(19.47) implies the Schrödinger equation

$$\left(-\frac{\hbar^2}{2m} \Delta_D(q) + \mathcal{W}(q) \right) \psi(q) = 0, \quad (19.48)$$

where

$$\mathcal{W}(q) = \sum_{k=1}^D \mathcal{W}_k(q_k), \quad (19.49)$$

and $\Delta_D(q)$ denotes the D -dimensional Laplacian

$$\Delta_D(q) = \sum_{k=1}^D \frac{\partial^2}{\partial q_k^2}. \quad (19.50)$$

Observe that

$$\psi(q) = \prod_{k=1}^D \psi_k(q_k), \quad (19.51)$$

where each ψ_k is solution of the one-dimensional SE

$$\left(-\frac{\hbar^2}{2m} \frac{\partial^2}{\partial q_k^2} + \mathcal{W}_k(q_k) \right) \psi_k(q_k) = 0. \quad (19.52)$$

Also in higher dimension the state $\mathcal{W}^0$ corresponds to the non-trivial solution

$$\mathcal{S}_0^0 = \frac{\hbar}{2i} \sum_{k=1}^D \ln \left(\frac{q_k^0 + i\bar{\ell}_{0,k}}{q_k^0 - i\ell_{0,k}} \right). \quad (19.53)$$

This guarantees that the Legendre transformation

$$\mathcal{S}_0 = \sum_{k=1}^D p_k \frac{\partial \mathcal{T}_0}{\partial p_k} - \mathcal{T}_0, \quad (19.54)$$

is defined for any physical system. As a consequence of the involutive nature of the Legendre transformation, we have that $\mathcal{S}_0$ - $\mathcal{T}_0$ duality extends to higher dimension.

A first remark for our construction is that Eq.(19.48), due to its structure, holds also for potentials which are more general than (19.46). To see this observe that the Laplacian is invariant under roto-translations of the coordinate (parity transformations can be included in the following construction)

$$\Delta_D(\tilde{q}) = \sum_{k=1}^D \frac{\partial^2}{\partial \tilde{q}_k^2} = \Delta_D(q) \quad (19.55)$$

where

$$\tilde{q}_k = \sum_{j=1}^D R_{kj} q_j + B_k, \quad (19.56)$$

$$R^t R = \mathbb{I}_D. \quad (19.57)$$

It follows that Eq.(19.52) is equivalent to

$$\left(-\frac{\hbar^2}{2m} \Delta_D(\tilde{q}) + \tilde{\mathcal{W}}(\tilde{q}) \right) \tilde{\psi}(\tilde{q}) = 0, \quad (19.58)$$

where $\tilde{\mathcal{W}}(\tilde{q}) = \mathcal{W}(q(\tilde{q}))$, and $\tilde{\psi}(\tilde{q}) = \psi(q(\tilde{q}))$. The observation is that in general both $\tilde{\mathcal{W}}(\tilde{q})$ and $\tilde{\psi}(\tilde{q})$ would not decompose as a sum and product respectively, namely

$$\tilde{\mathcal{W}}(\tilde{q}) = \sum_{k=1}^D \mathcal{W}_k(q_k) \neq \sum_{k=1}^D \tilde{\mathcal{W}}_k(\tilde{q}_k), \quad (19.59)$$

$$\tilde{\psi}(\tilde{q}) = \prod_{k=1}^D \psi_k(q_k) \neq \prod_{k=1}^D \tilde{\psi}_k(\tilde{q}_k). \quad (19.60)$$

Hence, removing the tilde from (19.58), we have that the EP implies at least the D -dimensional SE

$$\left(-\frac{\hbar^2}{2m} \Delta_D(q) + \mathcal{W}(q) \right) \psi(q) = 0, \quad (19.61)$$

where now $\mathcal{W}(q)$ is more general than (19.49).

19.3 Schrödinger equation and space compactification

The previous construction can be generalized to get a wider class of potentials. The point is to first consider the SE in $D + N$ dimensions. Eq.(19.47) implies

$$\left(-\frac{\hbar^2}{2m} \Delta_{D+N}(q) + \mathcal{W}(q) \right) \psi(q) = 0, \quad (19.62)$$

where

$$\mathcal{W}(q) = \sum_{k=1}^{D+N} \mathcal{W}_k(q_k). \quad (19.63)$$

Performing a rototranslation we have

$$\left(-\frac{\hbar^2}{2m} \Delta_{D+N}(\tilde{q}) + \tilde{\mathcal{W}}(\tilde{q}) \right) \tilde{\psi}(\tilde{q}) = 0, \quad (19.64)$$

which is equivalent to

$$\left(-\frac{\hbar^2}{2m} \Delta_D(\tilde{q}) + \tilde{\mathcal{W}}(\tilde{q}) - \frac{\hbar^2}{2m} \frac{\Delta_N(\tilde{q}) \tilde{\psi}(\tilde{q})}{\tilde{\psi}(\tilde{q})} \right) \tilde{\psi}(\tilde{q}) = 0, \quad (19.65)$$

where

$$\Delta_D(\tilde{q}) = \sum_{k=1}^D \frac{\partial^2}{\partial \tilde{q}_k^2}, \quad \Delta_N(\tilde{q}) = \sum_{k=D+1}^{D+N} \frac{\partial^2}{\partial \tilde{q}_k^2}. \quad (19.66)$$

Eq.(19.65) implies the D -dimensional SE

$$\left(-\frac{\hbar^2}{2m} \Delta_D(\tilde{q}) + \mathcal{W}_{eff}(\tilde{q}) \right) \tilde{\psi}(\tilde{q}) = 0, \quad (19.67)$$

where

$$\mathcal{W}_{eff}(\tilde{q}) = \tilde{\mathcal{W}}(\tilde{q}) - \frac{\hbar^2}{2m} \frac{\Delta_N(\tilde{q})\tilde{\psi}(\tilde{q})}{\tilde{\psi}(\tilde{q})}, \quad (19.68)$$

with $\tilde{q}_{D+1}, \dots, \tilde{q}_{D+N}$ seen as parameters for the potential.

The above approach seems to be sufficiently powerful to produce a wide range of potentials from compactification. It is then interesting to understand whether an arbitrary potential in D -dimensions can be obtained by a rototranslation and then compactifying N -dimensions. One should inductively define the compactification when $N \longrightarrow \infty$. There are many other questions concerning this approach. For example, it would be interesting to find in the case of some relevant potentials the associated minimal value of N such that it can be reproduced by compactification.

19.4 Time-dependent case

Let us consider a potential of the form

$$V(q, t) = \sum_{k=1}^D V_k(q_k, t). \quad (19.69)$$

Since

$$i\hbar \frac{\partial \psi_k(q_k, t)}{\partial t} = \left(-\frac{\hbar^2}{2m} \frac{\partial^2}{\partial q_k^2} + V_k(q_k, t) \right) \psi_k(q_k, t), \quad (19.70)$$

it follows that

$$i\hbar \frac{\partial \psi(q, t)}{\partial t} = \left(-\frac{\hbar^2}{2m} \Delta_D(q) + V(q, t) \right) \psi(q, t), \quad (19.71)$$

where

$$\psi(q, t) = \prod_{k=1}^D \psi_k(q_k, t). \quad (19.72)$$

Since space compactification does not affect the time component, we have that Eq.(19.71) holds for all the potentials obtained by the procedure described in the previous subsection.

19.5 Relativistic extension

A property of the EP is that it has a universal character. We already saw that it implies the cocycle condition also in higher dimension. This cocycle condition holds also in the relativistic case since it arises as consistency condition on the transformation properties of $\mathcal{W}$ and Q . Thus, in the relativistic case one arrives to the same cocycle condition (8.14) and to its higher dimensional version (19.40). In this sense the quantum correction has always the same structure. For example, in the one-dimensional relativistic case it is still given by the Schwarzian derivative of the reduced action times $\hbar^2/4m$. Since the classical stationary HJ equation has the form

$$\frac{1}{2m} \left(\frac{\partial \mathcal{S}_0^{cl}}{\partial q} \right)^2 + \mathcal{W}_{rel} = 0, \quad (19.73)$$

where

$$\mathcal{W}_{rel} \equiv \frac{1}{2mc^2}[m^2c^4 - (V - E)^2], \quad (19.74)$$

we immediately see that, according to the EP, its quantum version is

$$\frac{1}{2m} \left(\frac{\partial \mathcal{S}_0}{\partial q} \right)^2 + \mathcal{W}_{rel} + \frac{\hbar^2}{4m} \{\mathcal{S}_0, q\} = 0, \quad (19.75)$$

which has been considered also in [73]. Similarly to the non-relativistic case, we have

$$e^{\frac{2i}{\hbar} \mathcal{S}_0} = e^{i\alpha} \frac{w_{rel} + i\bar{\ell}}{w_{rel} - i\ell}, \quad (19.76)$$

where $w_{rel} = \phi^D/\phi$, with ϕ^D and ϕ denoting two real linearly independent solutions of the Klein-Gordon equation

$$-c^2 \hbar^2 \frac{\partial^2 \phi}{\partial q^2} + (m^2 c^4 - E^2 + 2EV - V^2) \phi = 0. \quad (19.77)$$

20 Conclusions

Let us make some concluding remarks concerning the main results of our investigation. We started by discussing our recent formulation of an EP from which the QSHJE was derived [1][2][12][13]. We focused on formulating the general theory by starting from basic concepts leading in a natural way to a new formulation of QM. As a result we obtained basic characteristics of QM from a first principle such as the postulated equivalence of states under spatial coordinate transformations. The initial sections were devoted to a critical examination of the rest frame and time parameterization as considered in CM. This analysis was done by a parallel investigation of a basic relationship between the Legendre transformation and second-order linear differential equations first observed in [6] in the framework of Seiberg-Witten theory [7]. We saw that the EP univocally leads to the QSHJE, which in turn implies the SE, which has a natural interpretation in terms of trajectories depending on initial conditions parameterized by the constant ℓ . These conditions are lost in the SE which we introduced as a mathematical tool to solve the QSHJE. Thus $\text{Re } \ell$ and $\text{Im } \ell$ can be seen as a sort of hidden variables depending on fundamental constants, notably on the Planck and Compton lengths. This suggests that fundamental interactions may in fact arise in the framework of the EP.

An alternative formulation of QM should reproduce the basic experimental facts, such as tunnel effect and energy quantization. In Bohm's theory [4] and stochastic quantization [19], the $L^2(\mathbb{R})$ condition for the wave-function in the case (17.6), which in fact implies tunnelling and energy quantization, still arises from a re-interpretation of the Copenhagen axioms. In our formulation we started with a criticism of the distinguished role of time and were then led to formulate the Equivalence Postulate in which spatial coordinate transformations are considered. We considered the transformations in the framework of HJ theory in which dynamics is described in terms of the functional relation between p and q , so that time parameterization does not appear. While in the Hamilton and Lagrange equations of motion time derivatives appear also in the stationary case, in HJ theory the time parameterization

is introduced only after one identifies p and $m\dot{q}$. In this way one can consider possible relations among different systems without introducing time parameterization, a concept that, as we saw, is related to the privileged nature of the rest frame. As a matter of fact, this property of HJ theory is in fact at the heart of our formulation of QM.

The above discussion illustrates in which sense our EP differs from that formulated by Einstein [3]. Loosely speaking, one can say that formulating Einstein's EP in a strong sense, that is for all possible potentials, and before introducing time parameterization, leads to QM. The basic fact is that implementation of the EP univocally leads to the QSHJE and therefore to the SE. Furthermore, it has been shown that the basic $L^2(\mathbb{R})$ condition is derived from the QSHJE itself. Thus, tunnelling and energy quantization directly follows from the EP and can be seen as basic tests of our formulation.

Our formulation has manifest p - q duality which traces back to the involutive nature of the Legendre transformation. A fact clearly expressed in terms of the Legendre brackets. This feature reflects in the appearance in the formalism of both ψ^D and ψ . In particular, while in the standard formulation of QM one usually considers a particular solution of the SE as the relevant quantity, here the relevant quantity is the ratio $w = \psi^D/\psi$ whose role is particularly transparent in considering the geometrical concept of trivializing map where, like in the case of $\mathcal{S}_0$, p and $\dot{q}$, ψ^D and ψ appear simultaneously.

Obtaining QM from a principle which is reminiscent of Einstein's EP [3], suggests that difficulties underlying quantization of gravity concern the way in which one usually considers the principles underlying the two theories. In particular, on general grounds, the fact that GR describes space-time as intrinsically related to matter, and that this is done in the framework of well-defined trajectories, indicates that the usual formulation of QM may have basic conceptual obstructions when gravitation is taken into account. The fact that QM arises from a simple principle, may in fact open the way to a reformulation of the problem of quantum gravity from a completely unexpected and somehow surprising perspective. In this context we note that the appearance of the Möbius symmetry and the role of the quantum potential may suggest that interactions present in Nature are deeply related to it. We saw that different initial conditions are related by a Möbius transformation. Furthermore, these conditions, seen as hidden variables, have an intrinsic dependence on fundamental constants such as the Planck length. A concrete signal that our formulation of QM is deeply connected to Gravitation.

Another distinguished feature of the present formulation is the fact that the quantum potential is never vanishing. This property is somehow reminiscent of the relativistic rest energy. This supports the old suspicion that special relativity itself and QM are in fact related. In the usual formulation this possible connection is suggested in considering the de Broglie wave-length. In this respect we note that in sect.16 we introduced a similar quantity expressed in terms of the conjugate momentum.

The above observations indicate that the problems one meets in formulating a theory in which the interactions appears to be unified, may in fact be connected to the usual scheme we have about basic concepts. A peculiarity of this picture is that it somehow reproduces the historical development of Theoretical Physics research. So, the standard picture is to think of QM as the basic framework in which interactions should be described. In this way we have that from one side there is the QM framework and on the other one there are the four distinguished fundamental interactions. The investigation we performed in the present paper may be an alternative starting point towards an effective unification

of fundamental interactions. However, in order to formulate it, we have to completely reconsider basic concepts such as the meaning of time, the role of trajectories and their connection with the concept of force at distance. In this context it is exciting to consider the possibility that the structure of interactions is in fact intimately related to the quantum potential that, as we have seen, plays the role of intrinsic energy. More precisely, it can be seen as a sort of self-energy of the particle whose structure depends on the external potential. In particular, this self-energy should provide the key to understand how particle interacts with external forces and can be considered as particle response to external perturbations. These observations would lead to replace the above scheme by a single theory.

It is not a fortuitous event that the EP we formulated originated from a critical analysis of the role of time in CM. This was done after noticing the peculiar nature of the rest frame. It is actually unavoidable that the classical concept of time loses its nature in the present formulation. This is somehow a parallel evolution of what happens in GR.

Another feature concerns the similarity of our formulation with the theory of uniformization of Riemann surfaces. It is even more frequent that this fascinating theory enters, in different contexts, in basic physical topics. We have seen that the concept of trivializing map is essentially the same as the one in Riemann surfaces theory. We believe that many of the geometrical concepts underlying our approach are related to the fact that the topic concerning complex structures of Riemann surfaces may be intimately related to the appearance of complex numbers in QM.

Acknowledgements. It is a pleasure to thank M. Appleby, D. Bellisai, M. Bochicchio, L. Bonora, R. Carroll, F. De Felice, G.F. Dell'Antonio, F. Guerra, F. Illuminati, A. Kholodenko, A. Kitaev, G. Marmo, R. Nobili, R. Onofrio, F. Paccanoni, P. Sergio, G. Travaglini, G. Vilasi and R. Zucchini for stimulating discussions. Special thanks are due to E.R. Floyd for interesting discussions on the QSHJE, to G. Bertoldi for contributions in some aspects of the present formulation of QM and to G. Bonelli, E. Gozzi, J.M. Isidro, P.A. Marchetti, P. Pasti and M. Tonin for basic discussions on the foundations of QM.

Work supported in part by DOE Grant No. DE-FG-0287ER40328 (AEF) and by the European Commission TMR programme ERBFMRX-CT96-0045 (MM).

References

- [1] A.E. Faraggi and M. Matone, Phys. Lett. **B450** (1999) 34, hep-th/9705108; Phys. Lett. **B437** (1998) 369, hep-th/9711028.
- [2] A.E. Faraggi and M. Matone, Phys. Lett. **B445** (1999) 357, hep-th/9809126.
- [3] A. Einstein, Annalen der Phys. **49** (1916) 769.
- [4] D. Bohm, Phys. Rev. **85** (1952) 166; ibidem 180.
- [5] M. Born, Rev. Mod. Phys. **21** (1949) 463.
- [6] M. Matone, Phys. Lett. **B357** (1995) 342, hep-th/9506102.
- [7] N. Seiberg and E. Witten, Nucl. Phys. **B426** (1994) 19, hep-th/9407087; Nucl. Phys. **B431** (1994) 484, hep-th/9408099.
- [8] A.E. Faraggi and M. Matone, Phys. Rev. Lett. **78** (1997) 163, hep-th/9606063.
- [9] R. Carroll, hep-th/9607219; hep-th/9610216; hep-th/9702138; hep-th/9705229; Nucl. Phys. **B502** (1997) 561; Lect. Notes Phys. **502**, Springer (Berlin, 1998), pp. 33 – 56.
- [10] I.V. Vanea, gr-qc/9801072.
- [11] G. Bonelli and M. Matone, Phys. Rev. Lett. **77** (1996) 4712, hep-th/9605090.
- [12] A.E. Faraggi and M. Matone, Phys. Lett. **249A** (1998) 180, hep-th/9801033.
- [13] A.E. Faraggi and M. Matone, Phys. Lett. **B445** (1998) 77, hep-th/9809125.
- [14] J.G. Muga, R. Sala and R.F. Snider, Phys. Scripta **47** (1993) 732.
- [15] E.R. Floyd, Phys. Rev. **D26** (1982) 1339.
- [16] G. Bertoldi, A.E. Faraggi and M. Matone, paper in preparation.
- [17] V.W. Guillemin and S. Sternberg, *Symplectic Techniques in Physics*, Cambridge Univ. Press (Cambridge, 1990).
B. Kostant, in “Lectures in Modern Analysis and Applications III”, Lect. Notes Math. **170** Springer (Berlin, 1970).
A.A. Kirillov, in “Dynamical Systems III”, Ed. by V.I. Arnold, Springer (Berlin, 1989) p. 137.
N.M.J. Woodhouse, *Geometric Quantization*, Clarendon Press (Oxford, 1992).
J. Sniatycki, *Geometric Quantization and Quantum Mechanics*, Springer-Verlag (New York, 1980).
J.-L. Brylinski, *Loop Spaces, Characteristic Classes and Geometric Quantization*, Birkhäuser (Basel, 1993).

- [18] J.R. Klauder and B.-S. Skagerstam, *Coherent States*, World Scientific (Singapore, 1985).
A.M. Perelomov, *Generalized Coherent States and Their Applications*, Springer (Berlin, 1986).
- [19] E. Nelson, *Quantum Fluctuations*, Princeton Univ. Press (Princeton, 1985).
F. Guerra, Phys. Rept. **77** (1981) 263.
- [20] E. Gozzi, Phys. Lett. **B158** (1985) 489; erratum, ibidem **B386** (1996) 495.
- [21] V. Periwai, Phys. Rev. Lett. **80** (1998) 4366, hep-th/9709200.
- [22] J.S. Anandan, *Classical and Quantum Physical Geometry*, in “Potential, Entanglement and Passion—at-a-Distance – Quantum Mechanics Studies for Abner Shimony”, Vol. 2. Ed. by R.S. Cohen, M. Horne and J. Stachel, Kluwer (Dordrecht, Holland 1997), pp. 31 – 52, gr-qc/9712015.
- [23] L. Viola and R. Onofrio, Phys. Rev. **D55** (1997) 455, quant-ph/9612039.
R. Onofrio and L. Viola, Mod. Phys. Lett. **A12** (1997) 1411, quant-ph/9706004.
- [24] M. Cini, Annals Phys. **273** (1999) 99, quant-ph/9807001; *Quantum Theory Without Waves: a Way of Eliminating Quantum Mechanical Paradoxes?*, Roma 1 preprint n.1103.
- [25] M. Gell-Mann and J.B. Hartle, *Quantum Mechanics in the Light of Quantum Cosmology*, in W.H. Zurek, “Complexity, Entropy, and the Physics of Information”, Reading, Addison-Wesley (MA, 1990), pp. 425 – 458.
- [26] N. Gisin and I.C. Percival, J. Phys. **A25** (1992) 5677.
I.C. Percival, Proc. Royal Soc. London **A447** (1994) 189.
- [27] S.L. Adler, *Quaternionic Quantum Mechanics and Quantum Fields*, Oxford Univ. Press (New York, 1995).
- [28] G.C. Ghirardi, A. Rimini and T. Weber, Phys. Rev. **D34** (1986) 470.
P. Pearle, Phys. Rev. **A39** (1989) 2277.
- [29] S. Kochen and E.P. Specker, Jour. Math. Mech. **17** (1967) 59.
- [30] J.S. Bell, *Speakable and Unspeakable in Quantum Mechanics*, Cambridge Univ. Press (Cambridge, 1987).
- [31] H.R. Brown, *Bell’s Other Theorem and its Connection with Non-Locality. Part 1.*, in “Bell’s Theorem and the Foundations of Modern Physics”, A. van de Merwe, F. Selleri and G. Tarozzi, Editors, World Scientific (Singapore, 1992).
- [32] C.J. Isham and J. Butterfield, quant-ph/9803055; Int. J. Theor. Phys. **38** (1999) 827, quant-ph/9808067.
- [33] V.I. Arnold, *Mathematical Methods of Classical Mechanics*, Springer-Verlag (New York, 1980).

- [34] R.C. Gunning, *Lectures on Riemann Surfaces*, Princeton Univ. Press (Princeton, 1966).
- [35] E. Cartan, Ann. Ec. Normale, **21** (1904) 153.
- [36] V.W. Guillemin and S. Sternberg, Bull. Amer. Math. Soc. **70** (1964) 16.
- [37] D.B. Fuks, *Cohomologies of Infinite-Dimensional Lie Algebras*, Consultants Bureau (New York, 1986).
- [38] G. Segal, Comm. Math. Phys. **80** (1981) 301.
- [39] V. Yu. Ovsienko, *Lagrange Schwarzian Derivative and Symplectic Sturm Theory*, CPT – 93/P.2890.
- [40] A. Messiah, *Quantum Mechanics*, Vol. 1, North-Holland (Amsterdam, 1961).
- [41] P.R. Holland, *The Quantum Theory of Motion*, Cambridge Univ. Press (Cambridge, 1993).
- [42] E.R. Floyd, Phys. Rev. **D25** (1982) 1547; **D29** (1984) 1842; Found. Phys. Lett. **9** (1996) 489, quant-ph/9707051; quant-ph/9708007; Int. J. Mod. Phys. **A14** (1999) 1111, quant-ph/9708026.
- [43] E.R. Floyd, Phys. Rev. **D34** (1986) 3246; Phys. Lett. **214A** (1996) 259.
- [44] A. Rosen, Am. J. Phys. **32** (1964) 377.
- [45] D.B. Berkowitz and P.D. Skiff, Am. J. Phys. **40** (1972) 1652.
- [46] J.O. Hirschfelder, A.C. Christoph and W.E. Palke, J. Chem. Phys. **61** (1974) 5435.
- [47] R. Schiller, Phys. Rev. **125** (1962) 1100; ibidem 1109.
A. Rosen, Am. J. Phys. **32** (1964) 597; Found. Phys. **16** (1986) 687.
- [48] A. Anderson, Annals Phys. **232** (1994) 292, hep-th/9305054.
- [49] P.A.M. Dirac, *The Principle of Quantum Mechanics*, 4th. ed., Oxford Univ. Press (Oxford, 1958).
- [50] H. Weyl, *The theory of Groups and Quantum Mechanics*, 2nd. ed., Dover (New York, 1950).
- [51] M. Born, W. Heisenberg and P. Jordan, Ztschr. f. Phys. **35** (1926) 557.
W. Heisenberg, Math. Ann. **95** (1926) 683.
P.A.M. Dirac, Proc. Roy. Soc. **A110** (1926) 561.
- [52] H.M. Farkas and I. Kra, *Riemann Surfaces*, 2nd. ed., Springer-Verlag (New York, 1992).
- [53] G.A. Jones and D. Singerman, *Complex Functions*, Cambridge Univ. Press (Cambridge, 1987).
- [54] M. Matone, Int. J. Mod. Phys. **A10** (1995) 289.

- [55] L. Bers, *On Moduli of Riemann Surfaces*, ETH, Zürich, 1964, (mimeographed); *Automorphic Forms and General Teichmüller Spaces*, Proceedings of the Conference on Complex Analysis, Minneapolis 1964, Springer-Verlag (New York, 1965).
- [56] A. Beurling and L.V. Ahlfors, *Acta Math.* **96** (1956) 124.
- [57] O. Pekonen, *Phys. Lett.* **B252** (1990) 555; *J. Geom. Phys.* **15** (1995) 227, hep-th/9310045.
- [58] M. Matone, *Phys. Rev.* **D53** (1996) 7354, hep-th/9506181.
- [59] C.G. Callan and F. Wilczek, *Nucl. Phys.* **B340** (1990) 366.
- [60] G. 't Hooft, *Dimensional Reduction in Quantum Gravity*, Essay dedicated to Abdus Salam. Published in Salamfest 1993; 0284 – 296, gr-qc/93100026.
C.R. Stephens, G. 't Hooft and B.F. Whiting, *Class. Quant. Grav.* **11** (1994) 621, gr-qc/9310006.
L. Susskind, *J. Math. Phys.* **36** (1995) 6377, hep-th/9409089.
- [61] J. Maldacena, *Adv. Theor. Math. Phys.* **2** (1998) 231, hep-th/9711200.
- [62] L. Susskind and E. Witten, hep-th/9805114.
- [63] L. Bers, *Universal Teichmüller Spaces*, in “Analytic methods in mathematical physics”, R.P. Gilbert, R.G. Newton, Editors, Gordon and Breach (New York, 1968).
- [64] H. Flanders, *J. Diff. Geom.* **4** (1970) 515.
- [65] V.P. Ermakov, *Univ. Izv. Kiev* **20** (1880) 1.
- [66] H.R. Lewis Jr., *Phys. Rev. Lett.* **18** (1967) 510; erratum, *ibidem* 636; *J. Math. Phys.* **9** (1968) 1976.
H.R. Lewis Jr. and W.B. Riesenfeld, *J. Math. Phys.* **10** (1969) 1458.
- [67] W.E. Milne, *Phys. Rev.* **D35** (1930) 863.
- [68] M. Lutzky, *Phys. Lett.* **68A** (1978) 3; **75A** (1979) 8; *J. Phys.* **A11** (1978) 249; **A12** (1979) 973.
P.G.L. Leach, *J. Math. Phys.* **18** (1977) 1902; **20** (1979) 96; **21** (1980) 300.
J.R. Ray and J.L. Reid, *Phys. Lett.* **71A** (1979) 317; *J. Math. Phys.* **20** (1979) 2054; **21** (1980) 1583.
H.J. Korsch, *Phys. Lett.* **74A** (1979) 294; **109A** (1985) 313.
H.J. Korsch and H. Laurent, *J. Phys.* **B14** (1981) 4213.
- [69] V.I. Arnold, *Ordinary Differential Equations*, MIT Press (1973).
- [70] V. Yu. Ovsienko, *Hook Law and Denogardus Great Number*, (Russian) *Kvant*, N.8, P.B – 16 (1989).
- [71] W. Thirring, *Quantum Mechanics of Atoms and Molecules*, Vol. 3, Springer-Verlag (New York, 1979).

- [72] W. Magnus and F. Oberhettinger, *Formulas and Theorems for the Functions of Mathematical Physics*, Chelsea (New York, 1954).
- [73] E.R. Floyd, Int. J. Theor. Phys. **27** (1988) 273.